



Received on 04 September, 2015; received in revised form, 15 February, 2016; accepted, 25 February, 2016; published 01 March, 2016

## SPECTROFLUORIMETRIC STUDY ON THE INTERACTION OF LOSARTAN POTASSIUM AND BOVINE SERUM ALBUMIN

Mingyan Sun, Ming Su and Hanwen Sun\*

College of Chemistry and Environmental Science, Hebei University, Key Laboratory of Analytical Science and Technology of Hebei Province, Baoding 071002, PR China

### Key words:

Bovine serum albumin,  
Losartan potassium, Fluorescence  
quenching, Binding constant,  
Three-dimensional fluorescence

### Correspondence to Author:

**Hanwen Sun**

College of Chemistry and  
Environmental Science, Hebei  
University, Key Laboratory of  
Analytical Science and Technology  
of Hebei Province, Baoding 071002,  
PR China.

**E-mail:** hanwen@hbu.edu.cn

**ABSTRACT:** The interaction of losartan potassium (LP) with bovine serum albumin (BSA) was studied by fluorescence quenching in combination with UV-Vis spectroscopic method under near physiological conditions. The fluorescence quenching rate constants and binding constants for BSA-LP system were determined at different temperatures. The fluorescence quenching of BSA by LP is due to static quenching and energy transfer. The results of thermodynamic parameters,  $\Delta H$  ( $-134.3 \text{ kJ mol}^{-1}$ ),  $\Delta S$  ( $-368 \text{ J mol}^{-1} \text{ K}^{-1}$ ) and  $\Delta G$  ( $-24.52$  to  $-20.83 \text{ kJ mol}^{-1}$ ), indicated that van der Waals interaction and hydrogen bonding played a major role for LP-BSA association. The competitive experiments demonstrated that the primary binding site of LP on BSA was located at site II in sub-domain IIIA of BSA. The distance between LP and a tryptophane unit was estimated to be 3.183 nm based on the Förster resonance energy transfer theory. The binding constant ( $K_a$ ) of BSA-LP at 298K was  $1.932 \times 10^4 \text{ L mol}^{-1}$ . Synchronous fluorescence and three-dimensional fluorescence studies showed that the presence of LP could change the conformation of BSA during the binding process.

**INTRODUCTION:** Serum albumins (SA) make a significant contribution to colloid osmotic blood pressure and aid in the transport, distribution and metabolism of many endogenous and exogenous ligands. Protein-drug binding greatly influences the absorption, distribution, metabolism and excretion properties of typical drugs<sup>1,2</sup>. Thus, it is necessary to study the interaction of drug with serum albumins at molecular level. Losartan potassium (LP) is a nonpeptide angiotensin II receptor antagonist used for the treatment of hypertension<sup>3</sup>.

It binds competitively and selectively to the angiotensin II subtype 1 (AT1) receptor, thereby blocking angiotensin II-induced physiological effects<sup>4</sup>. Following oral administration, losartan is well absorbed with a systemic bioavailability of approximately 33% and the mean peak plasma concentrations of losartan and its metabolite are reached in 1–2 h and in 3–4h, respectively<sup>5</sup>. The determination of losartan potassium in human urine<sup>6</sup> and rabbit plasma<sup>7</sup>, and human plasma<sup>8,9</sup> were carried out.

It was reported that high serum potassium levels after using losartan can reflect more severe renal disease<sup>10</sup>. Although the losartan determination and pharmacokinetic study were carried out<sup>7-9</sup>, the interaction of losartan with serum albumin was barely studied. To our knowledge, there is only a report for studying the interaction of LP at the

<b>QUICK RESPONSE CODE</b> 	<b>DOI:</b> 10.13040/IJPSR.0975-8232.7(3).1026-34
	Article can be accessed online on: <a href="http://www.ijpsr.com">www.ijpsr.com</a>
<b>DOI link:</b> <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.7(3).1026-34">http://dx.doi.org/10.13040/IJPSR.0975-8232.7(3).1026-34</a>	

binding sites of bovine serum albumin by equilibrium dialysis method in presence or absence of palmitic acid using ranitidine and diazepam as site-1 and site-2 specific probe, respectively<sup>11</sup>. However, no the information regarding energy transfer, binding forces and BSA conformation change for the interaction between LP and serum albumin was reported.

Human and bovine serum albumins exhibit similar binding chemistry due to the high percentage of sequence identities between the two proteins<sup>12</sup>. In this work, bovine serum albumin (BSA) was used because of its low cost and easy availability, and LP was used as a model drug. The interaction characteristic between LP and BSA under physiological conditions was examined by the fluorescence quenching in combination with the UV-vis spectroscopic method.

## Experimental:

### Drugs and reagents:

Commercially available bovine serum albumin (BSA, catalog no.A-7030, purity: 98%, M: 68,000) was purchased from Sigma Chemical Company. BSA stock solution ( $1.0 \times 10^{-4}$  M) was prepared by dissolving an appropriate amount of BSA with 0.1 M Tris-HCl (pH 7.4) buffer solution, and kept in the dark at 4 °C. BSA working solutions were prepared by diluting the stock solution with water. Losartan potassium (LP, purity: 98.7%) was purchased from the National Institutes for Food and Drug Control (Beijing, China).

A stock solution ( $1.0 \times 10^{-3}$  M) of LP was prepared in water, and stored in refrigerator at 4 °C. Tris-HCl buffer (pH 7.40) consists of Tris (0.1 M), HCl (0.1 M), and NaCl (0.5 M). NaCl solution was used to maintain the ion strength. All chemicals were of analytical reagent grade or better. Purified water was prepared by an XGJ- 30 highly pure water machine (Yongcheng purification Science & Technology Co. Ltd., Beijing, China). Equipment All fluorescence measurements were performed on an F-7000 Fluorescence spectrophotometer (Hitachi, Japan) which was equipped with a 1 cm quartz cell and thermostat bath. The spectrum data points were collected from 280 to 500 nm. The widths of the excitation and the emission slit were

both set at 5 nm. Fluorescence measurements were carried out at room temperatures.

The absorption spectra were performed on a TU-1900 double light Spectrophotometer (Beijing TAYASAF Science & Technology Co., Ltd., China) using a 1 cm quartz cell in the wavelength range of 200–500 nm. All pH measurements were performed with a pHS-3C pH meter (Shanghai, China).

### Determination of fluorescence intensity:

Five 10-mL clean and dried test tubes were taken, and 2 mL of 0.5 M NaCl, 2.0 mL Tris-HCl buffer (pH 7.40), 0.25 mL of  $4.0 \times 10^{-5}$  M BSA, and different volumes (0.5–2.5 mL) of LP standard solution of  $1.0 \times 10^{-4}$  M were added in each test tube, and diluted to 10 mL with water. The concentration of BSA was  $1.0 \times 10^{-6}$  M, and that of LP was 0.5, 1.0, 1.5, 2.0, and  $2.5 \times 10^{-5}$  M. Sixth test-tube containing only BSA solution at pH 7.4 was marked as “control”, and seventh test-tube containing only  $1 \times 10^{-5}$  M LP was used for the comparison. After mixing the solutions, these were allowed to stand for 15 min for maximum binding of LP to BSA. The fluorescence intensity after the correction of inner-filter effect was calculated by the equation<sup>13</sup>:  $F_{cor} = F_{obs} \exp(\frac{1}{2}A_{ex} + \frac{1}{2}A_{em})$ , where  $F_{obs}$  is fluorescence intensity measured before the correction of inner-filter effect,  $A_{ex}$  and  $A_{em}$  are absorbance of the test solution at excitation and emission wavelengths, respectively.


The corrected fluorescence intensity was used for studying on the interaction of LP and BSA. After corrected inner-filter effect, fluorescence intensity ( $F_0$ ) in the absence of quencher LP and the fluorescence intensity ( $F$ ) in the presence of quencher LP were measured at a wavelength of  $\lambda_{ex}=280$  nm and  $\lambda_{em}=340$  nm under temperature of 298, 303 and 308 K for estimating the interaction between LP and BSA.

## RESULTS AND DISCUSSION:

### Fluorescence quenching mechanism:

Fluorescence quenching refers to any process that decreases the fluorescence intensity of a sample. A variety of molecular interactions can result in fluorescence quenching of excited state fluorophores. These include molecular

rearrangements, energy transfer, ground state complex formation and collisional quenching. The quenching of fluorescence may be static or dynamic, and can be recognized by temperature dependence studies. The quenching rate constants are expected to decrease with increase in temperature for static quenching, while for dynamic quenching a reverse effect was observed<sup>14</sup>.




**FIG. 1: FLUORESCENCE SPECTRA OF BSA IN THE ABSENCE OR PRESENCE OF LP AT 280nm AFTER CORRECTED INTER-FILTER EFFECT**  
BSA,  $1.0 \times 10^{-6}$  M; LP ( $\times 10^{-5}$  M), a→f: 0, 0.5, 1.0, 1.5, 2.0 and 2.5; g: LP,  $1 \times 10^{-5}$  M

**Fig. 1** shows the fluorescence spectra of BSA in the absence and presence of LP after corrected inner-filter effect. No fluorescence of LP was observed at  $\lambda_{ex}=280$  nm. The fluorescence spectra of BSA show a broad band with maximum at  $\sim 340$  nm. It is observed that both fluorescence intensity of BSA decrease with increasing concentration of LP. A maximum fluorescence emission of BSA underwent spectral shift from 343.4.0 to 341.4 nm. It is suggested that BSA and LP formed a complex. The fluorescence quenching data are analyzed by the Stern–Volmer equation<sup>15</sup>:

$$F_0/F = 1 + k_q \tau_0 [Q] = k_{sv} [Q] \quad (1)$$

where  $F_0$  and  $F$  are the fluorescence intensity in the absence and presence of quencher, respectively.  $k_q$  is the quenching rate constant,  $\tau_0$  is the fluorescence life time of biopolymer BSA ( $\tau_0 = 10^{-8}$  s)<sup>16</sup>,  $k_{sv}$  and  $[Q]$  are the Stern–Volmer quenching constant and concentration of quencher, respectively. In this work, the Stern–Volmer plots of  $F_0/F$  vs concentration of LP were obtained (**Fig. 2**). The values estimated are given in **Table 1**.



**FIG. 2: STERN–VOLMER PLOTS OF  $F_0/F$  VS CONCENTRATION OF LP**

**Fig. 2 Table 1** The variation of  $F_0/F$  against LP concentration  $Q$  fits in the equation of  $y = c + mx$  with correlation coefficient ( $R$ ) of  $\geq 0.9945$ . Because the quenching rate constant ( $k_q$ ) decreased with increase in temperature and  $k_q$  values were greater than the maximum scatter collision quenching constants of various quenchers with the biomolecules ( $2.0 \times 10^{10}$  L mol<sup>-1</sup> s<sup>-1</sup>), indicating that the fluorescence quenching was mainly arisen from static quenching by complex formation instead of dynamic quenching<sup>14, 17</sup>.

**TABLE 1: QUENCHING PARAMETERS OF BSA–LOSARTAN POTASSIUM AT DIFFERENT TEMPERATURES**

T (K)	Equation	R	$K_{sv}$ (L mol <sup>-1</sup> )	$K_q$ (L mol <sup>-1</sup> ·s <sup>-1</sup> )
298	$F_0/F = 0.98 + 2.16 \times 10^4 [Q]$	0.9954	$2.16 \times 10^4$	$2.16 \times 10^{12}$
303	$F_0/F = 1.023 + 2.10 \times 10^4 [Q]$	0.9945	$2.10 \times 10^4$	$2.10 \times 10^{12}$
308	$F_0/F = 1.025 + 2.06 \times 10^4 [Q]$	0.9970	$2.06 \times 10^4$	$2.06 \times 10^{12}$

**Binding constant and binding site number:**

The binding of LP with BSA to form complex in the ground state is further understood on the basis of available binding site number and binding constant of the complex formation process.

Assuming the quenching belongs to a dynamic quenching. The following equation was used to calculate the binding constant and binding sites<sup>18,19</sup>:


$$\log [(F_0 - F)/F] = \log K_a + n \log [Q] \quad (2)$$

where  $K_a$  and  $n$  are the binding constant and binding site number, respectively. **Fig. 3** shows the plots of  $\log [(F_0 - F)/F]$  vs  $\log [Q]$ . Binding constant ( $K_a$ ) and the binding site number ( $n$ ) could be calculated from the intercept and slope, as shown in **Table 2**.

**TABLE 2: REGRESSION EQUATION, CORRELATION COEFFICIENT(R), BINDING CONSTANT ( $K_a$ ) AND THE NUMBER OF BINDING SITE ( $n$ ) BETWEEN LOSARTAN POTASSIUM AND BSA AT DIFFERENT TEMPERATURES**

T (K)	Equation	R	$K_a$ (L mol <sup>-1</sup> )	$n$
298	$\log[(F_0 - F)/F] = 4.286 + 0.998 \log[Q]$	0.9950	$1.932 \times 10^4$	0.998
303	$\log[(F_0 - F)/F] = 3.919 + 0.910 \log[Q]$	0.9960	$8.299 \times 10^3$	0.910
308	$\log[(F_0 - F)/F] = 3.521 + 0.829 \log[Q]$	0.9906	$3.319 \times 10^3$	0.829

**Fig. 3 Table 2** It is shown that the correlation coefficient of the regression equations for the curves of  $\log[(F_0 - F)/F]$  vs  $\log[Q]$  is approximately equal to 1. The binding constant decreased with the increase of temperature, showing that the quenching does not belong to a dynamic quenching, but belongs to a static quenching. The  $K_a$  values calculated for LP-BSA indicate a significant interaction between ligand and protein<sup>20,21</sup>.



**FIG. 3: THE PLOTS OF  $\log [(F_0 - F)/F]$  VS  $\log [Q]$  UNDER DIFFERENT TEMPERATURES**


**Interaction forces between LP with BSA:**

The interaction forces between drug and biomolecules include hydrogen bonds, van der Waals forces, electrostatic and hydrophobic interactions<sup>22</sup>. The temperature dependence of the

interaction of LP with BSA was investigated at 298, 303 and 308 K. The thermodynamic parameters can be evaluated from the Van't Hoff equation:

$$\ln K_a = -\Delta H/RT + \Delta S/R \quad (3)$$

where  $K_a$  is the binding constant at corresponding temperature  $T$ , and  $R$  is the gas constant. The enthalpy change ( $\Delta H$ ) and entropy change ( $\Delta S$ ) can be obtained from the slope and the ordinates at the origin of the Van't Hoff plot, respectively (**Fig. 4**).



**FIG. 4: VAN'T HOFF PLOT FOR THE INTERACTION OF LP WITH BSA AT pH 7.40.**

The free energy change,  $\Delta G$  is determined from the following relationship

$$\Delta G = \Delta H - T\Delta S \quad (4)$$

The values of  $\Delta G$ ,  $\Delta S$  and  $\Delta H$  are calculated and summarized in **Table 3**.

**TABLE 3: THERMODYNAMIC PARAMETERS OF THE INTERACTION BETWEEN LOSARTAN POTASSIUM AND BSA AT DIFFERENT TEMPERATURES**

T (K)	$K_a$ (L mol <sup>-1</sup> )	$\Delta H$ (kJ mol <sup>-1</sup> )	$\Delta S$ (J mol <sup>-1</sup> ·K <sup>-1</sup> )	$\Delta G$ (kJ mol <sup>-1</sup> )
298	1.932×10 <sup>4</sup>	-134.3	-368.4	-24.52
303	8.299×10 <sup>3</sup>			-22.67
308	3.319×10 <sup>3</sup>			-20.83

As can be seen in **Table 3**, a negative value of free energy ( $\Delta G$ ) supports the assertion that the binding process is spontaneous. The negative enthalpy ( $\Delta H$ ) and entropy ( $\Delta S$ ) values of the interaction of LP and BSA indicate that the binding is mainly enthalpy-driven and the entropy is unfavorable for it. van der Waals interactions and hydrogen bonding played major role in the reaction<sup>23</sup>.

#### Binding sites of LP on the BSA:

On the basis of the probe-displacement method, there are at least three relatively-high specific drug-

binding sites on the BSA molecules. These sites, commonly called warfarin, ibuprofen and digoxin-binding sites, are also denoted as Site I, Site II, and Site III, respectively<sup>24, 25</sup>. To further determine the binding site of LP, the competitive experiments were carried out at 298 K using warfarin, ibuprofen and digoxin as a Site I-, Site II- and Site III specific probe, respectively. Binding constant  $K_a$  and the binding site numbers  $n$  for the system in the presence of site specific probe are listed in **Table 4**.

**TABLE 4: BINDING CONSTANT ( $K_a$ ) AND THE NUMBER OF BINDING SITE ( $n$ ) BETWEEN LOSARTAN POTASSIUM AND BSA IN THE PRESENCE OF SITE-SPECIFIC PROBE AT 298 K.**


System	$K_a$ (L mol <sup>-1</sup> )	$n$	R
LP-BSA	1.932×10 <sup>4</sup>	0.998	0.9950
LP-BSA-warfarin	4.977×10 <sup>3</sup>	0.887	0.9948
LP-BSA-ibuprofen	2.924×10 <sup>3</sup>	0.809	0.9984
LP-BSA-digoxin	8.831×10 <sup>3</sup>	0.914	0.9922

In the presence of site specific probe, the binding constants for LP-BSA system decreased. In the case of ibuprofen probe, it is 2.924×10<sup>3</sup> L mol<sup>-1</sup>, obviously lower than that for the LP-BSA system without any specific probe, showing the competition of ibuprofen with LP at a same site. These results demonstrated that the primary binding site of LP at site II within subdomain IIIA of BSA. It is same with the result obtained with diazepam as site II specific probe<sup>11</sup>.

#### Energy transfer from BSA to LP:

Fluorescence resonance energy transfer is an important technique for investigating a variety of biological phenomena including energy transfer processes<sup>26</sup>. Here the donor and acceptor are BSA and LP, respectively. The fluorescence emission of BSA-LP solution at an excitation wavelength of 280 nm is only from BSA since LP is a non-fluorescence drug molecule. However, at this wavelength LP has weak absorption. It was observed that there was spectral overlap between

fluorescence emission of BSA and absorption spectra of LP (**Fig. 5**). It is suggested the possibility of fluorescence resonance energy transfer from BSA to LP molecules in solution.



**FIG. 5: SPECTRAL OVERLAP OF ABSORPTION OF LP WITH FLUORESCENCE OF BSA**

The region of integral overlap can be used to calculate the critical energy transfer distance ( $R_0$ ) between BSA (donor) and LP (acceptor) according to Förster's non-radiative energy transfer theory using Förster's equation<sup>20, 27</sup>. Based on this theory, the efficiency ( $E$ ) of energy transfer between donor (BSA) and acceptor (LP) can be calculated by Eq. 5:

$$E = R_0^6 / (R_0^6 + r^6) \quad (5)$$

where,  $r$  is the binding distance between donor and acceptor, and  $R_0$  is the critical binding distance when the efficiency ( $E$ ) of energy transfer is 50%, which can be calculated by Eq. (6):

$$R_0^6 = 8.8 \times 10^{-25} k^2 n^{-4} \Phi_D J \quad (6)$$

where, the  $k^2$  is the spatial orientation factor of the dipole;  $n$  is the refractive index of medium;  $\Phi_D$  is the quantum yield of the donor in the absence of acceptor; and  $J$  is the overlap integral of the emission spectrum of the donor and the absorption spectrum of the acceptor. The  $J$  value can be calculated by Eq. (7):

$$J = \int F(\lambda) \varepsilon(\lambda) \lambda^4 \Delta \lambda / \int F(\lambda) \Delta \lambda \quad (7)$$

where,  $F(\lambda)$  is the fluorescence intensity of the fluorescent donor,  $\varepsilon(\lambda)$  is the molar absorption coefficient of the acceptor. In the present case,  $k^2$ ,  $n$  and  $\Phi_D$  are 2/3, 1.336 and 0.118, respectively<sup>12</sup>. The efficiency ( $E$ ) of energy transfer can be determined by Eq. (8):

$$E = 1 - F/F_0 \quad (8)$$

where,  $F_0$  and  $F$  are the fluorescence intensities of BSA solutions in the absence and presence of LP, respectively. From the overlapping we found  $R_0 = 2.352$  nm from Eq. (7) using  $K^2 = 2/3$ ,  $n = 1.336$  and  $\Phi_D = 0.118$  (tryptophan residue) for the aqueous solution of BSA.  $J$  value could be calculated from Eq. 7 and the corresponding result was  $7.80 \times 10^{-15} \text{ cm}^3 \text{ L mol}^{-1}$ .  $E$  value calculated from Eq. 8 was 14.01%. At the same time, the binding distance ( $r = 3.23$  nm) between BSA and LP was obtained by Eq. 5. Apparently, it is less than 7.0 nm. This result indicates the non radiative energy transfer from BSA to LP occurs with high possibility<sup>28, 29</sup>. It also suggested that the bindings of LP to BSA molecules were formed through energy transfer, which quenched the fluorescence of BSA molecules, indicating the presence of static quenching interaction between BSA and LP<sup>30</sup>.

#### Change of BSA conformation:

Synchronous fluorescence is a kind of simple and sensitive method to measure the fluorescence quenching. It can provide the information of polarity change around the chromophore microenvironment.  $\Delta \lambda$ , representing the difference between excitation and emission wavelengths, is an important operating parameter. When  $\Delta \lambda$  is 15 nm, synchronous fluorescence is characteristic of tyrosine residue, while when  $\Delta \lambda$  is 60 nm, it provided the characteristic information of tryptophan residues<sup>29</sup>. The synchronous fluorescence spectra of tyrosine residue and tryptophan residues in BSA with addition of LP were observed.





FIG. 6: SYNCHRONOUS FLUORESCENCE SPECTRA OF BSA WITH DIFFERENT AMOUNT OF LP  
T = 298 K; BSA,  $1.0 \times 10^{-6}$  M; LP ( $10^{-5}$  M), a→f: 0, 0.5, 1.0, 1.5, 2.0, 2.5

**Fig. 6** shows that the fluorescence intensity of tyrosine increased lightly with increasing LP content, but no obviously change of maximum emission wavelength was observed, while the fluorescence intensity of tryptophan decreased regularly, with the change of maximum emission wavelength from 338.8 to 341.2 nm. The red shift of fluorescence spectrum suggests the polarity of the surrounding environment increasing<sup>31</sup>. The fluorescence intensity and the pitch of quenching for  $\Delta\lambda=60$  nm was much higher than those for  $\Delta\lambda=15$  nm. Such result means LP is closer to tryptophan residues than tyrosine residues, namely

binding sites mainly are focused on tryptophan moiety. Above result showed that LP could change the conformation of BSA because the microenvironment around tryptophan residue changed during the binding process.

The three-dimensional fluorescence spectrum is another powerful method for studying conformation change of BSA. In this work, the three-dimensional fluorescence spectra and the contour spectra of BSA and BSA-LP systems were observed, as shown in **Fig. 7**, and the characteristic parameters are summarized in **Table 5**.



**FIG. 7: THREE-DIMENSIONAL FLUORESCENCE SPECTRA (LEFT) AND CONTOUR SPECTRA (RIGHT) OF BSA AND BSA-LP SYSTEMS**

BSA,  $1 \times 10^{-6}$  M; LP,  $1 \times 10^{-5}$  M

**TABLE 5: THREE-DIMENSIONAL FLUORESCENCE SPECTRA CHARACTERISTIC PARAMETERS OF THE BSA AND BSA-LOSARTAN POTASSIUM SYSTEMS**

System	Parameters	Peak 1	Peak 2
BSA	Peak position ( $\lambda_{ex}/\lambda_{em}$ , nm)	275.0/340.0	225.0/340.0
	Relative intensity	163.5	154.5
	Stokes shift $\Delta\lambda$ (nm)	65	115
BSA-Losartan	Peak position ( $\lambda_{ex}/\lambda_{em}$ , nm)	280.0/340.0	225.0/340.0
	Relative intensity	130.2	88.57
	Stokes shift $\Delta\lambda$ (nm)	60	115

From **Fig. 7**, peak 1 ( $\lambda_{\text{ex}}/\lambda_{\text{em}}$ : 275.0/340.0 nm) reveals the spectral characteristic of tryptophan and tyrosine residues. After the addition of LP, the fluorescence intensity of BSA decreased from 163.5 to 130.2. This suggests a less polar environment of both residues and almost all the hydrophobic amino acid residues of BSA were buried in the hydrophobic pocket. Less polar environment means that the binding position between LP and BSA located within this hydrophobic pocket, the addition of LP changed the polarity of this hydrophobic microenvironment and the conformation of BSA<sup>32</sup>.

In **Fig. 7**, peak 2 ( $\lambda_{\text{ex}}/\lambda_{\text{em}}$ : 225.0/340.0 nm) reveals the fluorescence spectra behavior of polypeptide backbone structures, which is caused by the transition of  $\pi-\pi^*$  of BSA's characteristic polypeptide backbone structure C=O<sup>33</sup>. After the addition of LP, peak 2 decreased dramatically with increasing concentration of LP. The fluorescence intensity of BSA decreased from 154.5 to 88.47. After addition of LP in BSA, fingerprint lines in below part of contour spectra changed to be sparse. These revealed that the microenvironment and conformation of BSA were changed in the binding reaction. The interaction between LP and BSA induced the unfolding of the polypeptides chains of BSA and conformational change of BSA.

**CONCLUSION:** The results showed that losartan potassium can bind to bovine serum albumin by hydrogen bonding and Van der Waals forces. The primary binding for losartan potassium was located at site II in subdomain IIIA of BSA. The result observed by synchronous fluorescence and three-dimensional fluorescence spectra demonstrated that the presence of LP induced some micro environmental and conformational changes of BSA molecules. The results are of great importance in pharmacy, pharmacology and biochemistry, and are expected to provide important insight into the interactions of the physiologically important protein BSA with drugs.

**ACKNOWLEDGEMENT:** This work was supported by the Science Foundation Office of Hebei Province (B2014201171).

## REFERENCES:

1. Zhou B, Qi ZD, Xiao Q, et al: Interaction of loratadine with serum albumins studied by fluorescence quenching method. *J. Biochem. Biophys. Meth.*, 2007; 70: 743–747.
2. Kandagal PB, Shaikh SMT, Manjunatha DH, et al: Spectroscopic studies on the binding of bioactive phenothiazine compounds to human serum albumin. *J. Photochem. Photobiol., A* 2007; 189: 121–127.
3. Burnier M: Angiotensin II Type 1 Receptor Blockers. *Circulation*, 2001; 103: 904–912.
4. Sica DA, Gehr TW and Ghosh S: Clinical pharmacokinetics of losartan. *Clin. Pharmacokinet.*, 2005;44: 797–814.
5. Tamimi JJI, Salem II, Mahmood Alam S, et al: Comparative pharmacokinetics of two tablet formulations of Losartan: bioequivalence assessment. *Biopharm. Drug Dispos.*, 2005; 26: 205–210.
6. Cagigal E, Gonzalez L, Alonso R, et al: Experimental design methodologies to optimise the spectrofluorimetric determination of losartan and valsartan in human urine. *Talanta*, 2001; 54: 1121–1133.
7. Demirkaya-Miloglu F, Yaman M E and Kadioglu Y: A new spectrofluorimetric method for determination of losartan potassium in rabbit plasma and its application to pharmacokinetic study. *Lumin.*, 2015; 30: 53–59.
8. Salvadori MC, Moreira RF, Borges BC, et al: Simultaneous determination of losartan and hydrochlorothiazide in human plasma by LC/MS/MS with electrospray ionization and its application to pharmacokinetics. *Clin. Exp. Hypertens*, 2009; 31: 415–427.
10. Karra VK, Pilli NR, Inamadugu JK, et al: Simultaneous determination of losartan, losartan acid and amlodipine in human plasma by LC-MS/MS and its application to a human pharmacokinetic study. *Pharm. Methods*, 2012; 3: 18–25.
11. Gonçalves A R and El Nahas A M: High serum potassium levels after using losartan can reflect more severe renal disease. *Diabetologia*, 2011; 54: 2963–2964.
12. Ferdosi Kabir A, Nazim Uddin K, Nazmus Sadat AFM, et al: Interaction of palmitic acid with losartan potassium at the binding sites of bovine serum albumin. *ARS Pharm.*, 2010; 51: 28–36.
13. Shaikh SMT, Seetharamappa J, Kandagal PB, et al: Binding of the bioactive component isothipendyl hydrochloride with bovine serum albumin. *J. Mol. Struct.*, 2006; 786: 46–52.
14. Lakowicz JR: Principles of fluorescence spectroscopy, 3rd ed., Springer Science+Business Media, New York, 2006.
15. Lakowicz JR: Principles of Fluorescence Spectroscopy, 3rd edn., Plenum Press, New York, 2006, pp. 277–350.
16. Zhang LN, Wu FY and Liu AH: Study of the interaction between 2, 5-di-[2-(4-hydroxyphenyl) ethylene]-terephthalonitril and bovine serum albumin by fluorescence spectroscopy. *Spectrochim. Acta, A* 2011; 79: 97–103.
17. Lakowicz JR and Weber G: Quenching of fluorescence by oxygen. A probe for structural fluctuations in macromolecules. *Biochem.*, 1973; 12: 4161–4170.
18. Maurice RE and Camillo AG: Fluorescence quenching studies with proteins. *Anal. Biochem.*, 1981; 114: 199–212.
19. Johansson JS: Binding of the volatile anesthetic chloroform to albumin demonstrated using tryptophan fluorescence quenching. *J. Biol. Chem.*, 1997; 272: 17961–17965.

20. Sūkowska JA, Równicka J, Bojko B, et al: Interaction of anticancer drugs with human and bovine serum albumin. *J. Mol. Struct.*, 2003; 651: 133–140.
21. Hu YJ, Liu Y, Zhao RM, et al: Spectroscopic studies on the interaction between methylene blue and bovine serum albumin. *J. Photochem. Photobiol., A* 2006;179: 324–329.
22. Barbero N, Barni E, Barolo C, et al: A study of the interaction between fluorescein sodium salt and bovine serum albumin by steady-state fluorescence. *Dyes Pigments*, 2009; 80: 307–313.
23. Zhang YZ, Chen XX, Dai J, et al: Spectroscopic studies on the interaction of lanthanum(III) 2-oxo-propionic acid salicyloyl hydrazone complex with bovine serum albumin. *Lumin.*, 2008; 23: 150–156.
24. Ross DP and Subramanian S: Thermodynamics of protein association reactions: forces contributing to stability. *Biochem.*, 1981; 20: 3096–3102.
25. Kragh-Hansen U: Molecular aspects of ligand binding to serum albumin. *Pharmacol. Rev.*, 1981; 33: 17–53.
26. Dubois N, Lopicque F, Magdalou J, et al: Stereoselective binding of the glucuronide of ketoprofen enantiomers to human serum albumin. *Biochem. Pharmacol.*, 1994; 48: 1693–1699.
27. Mallick A, Haldar B and Chattopadhyay N: Spectroscopic investigation on the interaction of ICT probe 3-acetyl-4-oxo-6, 7-dihydro-12H indolo-[2,3-a] quinolizine with serum albumins. *J. Phys. Chem., B* 2005; 109:14683–14690.
28. Bhattar SL, Kolekar GB and Patil S.R: Fluorescence resonance energy transfer between perylene and riboflavin in micellar solution and analytical application on determination of vitamin B<sub>2</sub>. *J. Lumin.*, 2008; 128: 306–310.
29. He WY, Li Y, Xue CX, et al: Effect of Chinese medicine alpinetin on the structure of human serum albumin. *Bioorg. Med. Chem.*, 2005; 13: 1837–1845.
30. Guo M, Lue WJ, Li MH, et al: Study on the binding interaction between carnitine optical isomer and bovine serum albumin. *Eur. J. Med. Chem.*, 2008; 43: 2140–2148.
31. Hu YJ, Liu Y, Zhang LX, et al: Studies of interaction between colchicines and bovine serum albumin by fluorescence quenching method. *J. Mol. Struct.*, 2005; 750: 174–178.
32. Wang YQ, Zhang HM, Zhang GC, et al: Binding of brucine to human serum albumin. *J. Mol. Struct.*, 2007; 830: 840–845.
33. Guo XJ, Sun XD and Xu SK: Spectroscopic investigation of the interaction between riboflavin and bovine serum albumin. *J. Mol. Struct.*, 2009; 931: 55–59.
34. Xiao Q, Huang S, Liu Y, et al: Thermodynamics, conformation and active sites of the binding of Zn–Nd hetero-bimetallic schiff base to bovine serum albumin. *J. Fluoresc.*, 2009; 19: 317–326.

**How to cite this article:**

Sun M, Su M and Sun H: Spectrofluorimetric Study on the Interaction of Iosartan Potassium and Bovine Serum Albumin. *Int J Pharm Sci Res* 2016; 7(3): 1026-34. doi: 10.13040/IJPSR.0975-8232.7(3).1026-34.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)