



Received on 23 September, 2017; received in revised form, 08 December, 2017; accepted, 06 January, 2018; published 01 June, 2018

RECENT DEVELOPMENTS ON NATURAL TRANSDERMAL PENETRATION ENHANCERS

Shivani Kala* and Divya Juyal

Himalayan Institute of Pharmacy and Research Rajawala, Dehradun - 248007, Uttarakhand, India.

Keywords:

Penetration, Enhancers,
Bioavailability, Transdermal, Natural

Correspondence to Author:

Shivani Kala

Associate Professor,
Himalayan Institute of Pharmacy
and Research Rajawala, Near Atak
Fram Selaqui, Dehradun - 248007
Uttarakhand, India.


E-mail: shivani.kala88@gmail.com

ABSTRACT: The drug delivery by oral route is so far most convenient and accepted route of drug delivery but now a days considerable interest has been increased in delivery of drugs through skin to the systemic circulation and for local effect. However, the outer most layer of the human skin, stratum corneum; possess the formidable barrier to drug penetration thereby reducing bio-availability. Most of the drugs do not have an ability to penetrate the stratum corneum. Owing to the side effects of chemical penetration enhancers interest has been aroused among researchers to find out natural sources as penetration enhancers to improve the bio-availability of drugs so that they can be delivered through skin. The current review aims to screen various natural sources of penetration enhancers for transdermal delivery of different drugs as they have minimum chances of side effects and have improved efficacy compared to chemical enhancers. Various natural derivatives have been proposed that could show potential penetration enhancing capacity with varied mechanism of penetration. Though a large number of natural sources of penetration enhancers have been discussed but still a lot of studies and research need to be done to explore the sources of natural penetration enhancers for effective delivery of drug *via* transdermal route.

INTRODUCTION: The transdermal route always provide innovative research area in drug delivery, and many drug delivery products are under clinical evaluation related to the transdermal or dermal system. The success of a dermatological drug to be used for systemic drug delivery depends on the ability of the drug to penetrate through skin in sufficient quantities to achieve the desired therapeutic effect¹. Owing to some complexities skin have a barrier for penetration through it.

To enhance the transport of drug through skin various techniques are applied called as permeation enhancement techniques and agents utilized in it are Penetration Enhancers² also called sorption promoters or accelerants) which penetrate into skin to reversibly decrease the barrier resistance³.

Brief Anatomy of the Skin: Skin is a complex biological structure consisting of many layers of a thickness of 2.97 ± 0.28 mm. Its function is to safeguard the major internal parts of the body from the external effects, regulation of temperature, sensation and water balance. A normal adult human body skin cover around two square meters surface area and gets about one-third of the blood circulating in the body. **Fig. 1** shows the structure of skin.

<p>QUICK RESPONSE CODE</p> 	<p>DOI: 10.13040/IJPSR.0975-8232.9(6).2190-96</p>
<p>Article can be accessed online on: www.ijpsr.com</p>	
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.9(6).2190-96</p>	

The skin is considered to have four distinct layers of tissue⁴.

1. Non-viable epidermis (stratum corneum)
2. Viable epidermis
3. Viable dermis
4. Subcutaneous connective tissue (hypodermis)




FIG. 1: STRUCTURE OF SKIN⁵

Advantages and Disadvantages of Penetration Enhancers:

Various advantages of penetration enhancers include⁶

- They help to provide penetration rate of drug sufficiently high for therapeutic efficiency
- It facilitates the absorption of non-absorbable drugs through skin.
- It can improve the transdermal absorption of topical preparation completely.
- It determines the penetration rate of transdermal drug delivery system.
- The terpenes like limonene in propylene glycol solution are effective penetration enhancer for cytotoxic drugs.

Various disadvantages of penetration enhancers are⁷

- The effective concentration varies from drug to drug.
- The use of various penetration enhancers with various concentrations vary
- Physicochemical properties of enhancers are also affecting the side effects in the body.

Mechanism of Action of Penetration Enhancers:

Skin Absorption Promoters may act by one or more of three potential mechanisms according to the lipid-protein-partitioning theory.

Firstly, penetration enhancers can alter the inter-cellular lipid structure between the corneocytes to increase diffusivity.

Secondly, they can modify intracellular protein domains within the horny layer.

Thirdly, they may increase the partitioning of the drug into the skin tissue⁸. Fig. 2 demonstrates the possible mechanism of drug penetration.

The three main functions of penetration enhancers include:⁹

Lipid Disruption: They change the structure of stratum corneum lipid organization and make it permeable for the drug *e.g.* Azone, terpenes, fatty acids, Dimethyl sulfoxide (DMSO) and alcohol.

Protein Modification: They interact with keratin and corneocytes and open up the dense protein structure and make it more Permeable *e.g.* Ionic surfactants.

Partitioning Promotion: Many solvents change the solution properties of horny layer and increase the partition of a drug, co-enhancer and co-solvents.

Group of Substances That Could Show Potential Absorption Promoter Activity: Table 1 gives the example of various potential absorption promoters.


FIG. 2: PERMEATION PATHWAYS OF SKIN¹⁰

TABLE 1: GIVES THE EXAMPLE OF VARIOUS POTENTIAL ABSORPTION PROMOTERS

Type of Penetrator	Mode of Action	Example
Physical Penetration enhancers ¹¹	Rate control over the release and transdermal permeation of drugs	1. Iontophoresis 2. Sonophoresis 3. Phonophoresis 4. Magnetophoresis 5. Electroporation 6. Thermophoresis 7. Radiofrequency 8. Needleless injection
Chemical enhancers ¹²	They act by three mechanisms 1. By disruption of highly ordered structure of stratum corneum lipid. 2. By interaction with intercellular protein. 3. By improved partition of the drug or solvent into stratum corneum.	Sulphoxides and similar chemicals-dimethyl sulphoxide(DMSO), Dimethyl formamide (DMF), dimethyl acetamide (DMAC) 2. Azones 3. Pyrrolidones 4. Fattyacids–Lauric acid, Myristic acid and capric acid 5. oxazolindiones (4 decyclozolidine-2-one) 6. Amine and Amides – Urea
Essential oils and terpenes ¹³	It may increase one or more of following effects 1. Partition coefficient 2. Diffusion coefficient 3. Lipid Extraction 4. Drug Solubility 5. Macroscopic Barrier Perturbation 6. Molecular Orientation of Terpenes Molecule with Lipid Bilayer	Essential oils: Niaouli Oil, Eucalyptus oil, Neem oil, Peppermint oil, etc. Terpenes: Geraniol, nerol, Menthol, Carveol, Camphor, etc.
Fixed Oils/ Fatty Acids ¹⁴	Oleic acid might increase the permeability via a mechanism involving perturbation of stratum corneum lipid bilayer and lacunae formation as penetration enhancing effect. (1) lipids could alter the permeability of the stratum corneum by having a direct effect and/or (2) solubility of the drug could be increased	Fish oil, cod liver oil, Fatty Acids from Algae, Phospholipids, oleic acid
Polysaccharides ¹⁵	Chitosan interact with negative charges in the skin to improve drug diffusion into the deeper layers of the skin. <i>A. vera</i> gel had a higher permeation enhancement which was explained by the fact that a drug with a larger molecular weight effectively blocks the permeation routes allowing increased possibility for the drug to interact with the enhancing factor and complex with it prior to being transported across the skin.	Chitosan, Aloe vera Gel,
Miscellaneous ¹⁶	Capsaicin reduces the diffusional resistance of the intercellular domains by inserting itself into the lipid bilayers within the intercellular channels thereby disrupting their stacking. moderate improvement in the permeability of the stratum corneum can be attributed to the restricted insertion of Vitamin E in the ceramide-rich bilayer structure.	Capsaicin, Vitamin E

Non-ionic Surfactant ¹⁷	the surfactant interacted with skin to destructure lipids and thus increase permeability; however the ability of surfactant to influence skin permeation was dependent on the physicochemical properties of permeant.	Tweens and brij
Drug Vehicle based ¹⁸	Based on drug section, vesicles and particles, prodrug, chemical potential of drug and eutectic system	Niosomes, liposome, aquasomes, Transfersomes, nanoparticles

Natural Penetration Enhancers: Various natural sources have been investigated for potential skin penetration enhancement activity. Some of the natural sources have been discussed below:

***Aloe vera*:**¹⁹ *Aloe barbadensis* Miller (*Aloe vera* Linn.) is the most widely used both commercially and for its therapeutic properties. This plant is having various medicinal, cosmetic and nutraceutical purposes *Aloe vera* gel is the viscous, transparent and colourless mucilaginous gel obtained from the parenchymatous cells in the fresh leaves of *Aloe vera*.

Penetration enhancers work possibly by increasing the solubility of the drug within the SC by altering the partitioning of the drug into the SC and/or by influencing the diffusion of the drug across the SC by disrupting the ordered nature of the skin lipids. It can soak into all the layers of the skin and this may be helpful in increasing the penetration of certain drug molecules across the skin, as lignins can penetrate the toughened areas of the skin.

Histopathological examination of *Aloe vera* oil treated SC suggested additional mechanisms for permeation *i.e.* transient reduction in barrier resistance of SC and intracellular transport by dekeratinization of corneocytes which may be attributed to the presence of triglycerides as constituents of *Aloe vera* oil. It is feasible to deliver therapeutically effective dose of LP *via* transdermal route using *Aloe vera* oil as penetration enhancer²⁰.

Essential Oils: Essential oils are volatile, odoriferous substances found in the flowers, fruit, leaves and roots of certain plants. The skin penetration enhancing effect of several isolated essential oils has been thoroughly investigated²¹. Essential oils and their constituents have been widely investigated as safe and suitable skin penetration enhancers for both hydrophilic and hydrophobic drugs but the mechanism of their action is not fully understood. The predominant compounds within oils are terpenes, terpenoids, phenyl propanoids, as well as minor amounts of miscellaneous volatile organic compounds.

TABLE 2: NATURAL OILS AS PENETRATION ENHANCER

Ingredient	Chemical Composition	Drug Used	Ref
Clove oil	Eugenol, Caryophyllene, Eugenyl acetate, Caryophyllene oxide	Ibuprofen	22
Angelica oil	α -Pinene, trans- β -Ocimene, Ligustilide, Eugenol	Ibuprofen	23
chuanxiong oil	(+)- α -Pinene, γ -Terpinene, Terpinolene, Methyl 4-ethylbenzoate	Flurbiprofen	24
Eucalyptus Oil	1,8-cineole	ketoconazole	25
Peppermint Oil	L-Menthol	5-fluorouracil	26
Cyperus Oil	Cyperene, Dehydrofukinone, α -Cyperone	Ibuprofen	27
Cinnamon Oil	trans-cinnamaldehyde, (E)-Cinnamyl acetate, linalool	nifedipine	28
Turpentine Oil	α -pinene, β -pinene, camphene, longifolene, A3-carene, limonene and β -caryophyllene	Labetalol Hydrochloride	29
Cardamom Oil	β -pinene, α -phellandrene, limonene, p-cymene, terpinolene, 1,8-cineole, γ -terpinene, linalool, linalyl acetate, α -terpineol acetate, citronellol, nerol, terpinen-4-oil, α -terpineol, geraniol, methyl eugenol and trans-nerolidol.	Piroxicam	30
Mint oil	Menthone, Menthol, α -terpineol, Eucarvone	Geniposide	31
rose oil	citronellol, geraniol, nerol, linalool, phenyl ethyl alcohol, farnesol	Losartan	32
tea tree oil	terpinen-4-ol, γ -terpinene, 1,8-cineole, α -terpinene, α -terpineol, p-cymene, and α -pinene	Losartan	33
Tulsi oil	Methyl eugenol, carvacrol, caryophyllene	celecoxib	34
Fennel oil	fenchone (ketone), trans-anethole (phenolic ester), α -pinene (monoterpene), linalol (alcohol), and methyl chavicol (phenol).	trazodone hydrochloride	35
Niaouli Oil	1,8-cineole (oxide) and limonene (monoterpene), α -pinene	estradiol	36

Black cumin oil	(monoterpene), β -pinene (monoterpene) and viridiflorol thymoquinone, thymohydroquinone, dithymoquinone, thymol, carvacrol, nigellimine-N-oxide, nigellidine and alpha-hederin.	carvedilol	37
Lemon grass oil	myrcene, citronellal, geranyl acetate, nerol, geraniol, neral and traces of limonene and citral.	hydrocortisone acetate	38
Rosemary oil	1,8-cineol, α -pinene, camphor, camphene, (E)-caryophyllene	diclofenac	39
Alpinia oxyphylla oil	1H-cycloprop[e]azulene, octahydro-1,8-dimethyl-7-(2-methylethenyl)-naphthalene, α -panasinsen, germacrene B, humulene 6,7-epoxide, cis- α -copaene-8-ol and nootkatone	indomethacin	40

Essential Fatty Acids: Fatty acids have been shown to accelerate skin permeation of drugs probably modifying intercellular lipid packing in the stratum corneum lipid domains and so reduce the barrier function. They can form lipophilic complexes with compounds. Although fatty acids can be used to enhance the permeation of both lipophilic and hydrophilic APIs, the flux of polar APIs is improved to a larger degree ⁴¹. Unsaturated fatty acids (UFAs) with a C18 chain length (e.g. oleic

and linoleic acids), increase penetration-enhancing effects with linolenic acid as the most potent, followed by oleic acid, palmitoleic acid, linoleic acid and arachidonic acid ⁴². Some of the Fatty acids that has been studied as penetration enhancers are listed below in **Table 3**.

Vegetable oils: Various vegetable oils have been investigated for their penetration enhancement activity as shown in **Table 4**.

TABLE 3: FATTY ACIDS AS PENETRATION ENHANCERS

Ingredient	Chemical Composition	Drug used	Ref
Evening primrose oil (EPO),	linoleic acid (70%-80%), γ -linolenic acid (8% - 12%), and oleic acid (6%-11%)	Flurbiprofen	43
Vitamin F fish oil	linoleic (35.5%), linolenic (30.5%), and oleic acid (21.4%).	Flurbiprofen	44
Botryococcus braunii (Algae)	ω -3 fatty acids eicosapentaenoic acid (EPA), and docosahexaenoic acid (DHA)	atenolol	45
soya bean	unsaturated fatty acids	flurbiprofen	46
egg yolk	phospholipids Phosphatidylcholine and 70% phosphatidylethanolamine	diclofenac	47
	levo- α -phosphatidylcholine	naproxen	48

Table 4: VEGETABLE OIL AS PENETRATION ENHANCERS

Vegetable oil	Experiment	Ref
Sunflower oil, Sesame oil and jojoba oil	Studies revealed that this oil in concentration of 10 % w/w of polymer weight exhibited enhancing effects on losartan potassium release from transdermal patches, with the cumulative amount of drug release	49
Almond oil	It was used as penetration enhancer in various concentrations to significantly enhance the penetration of drug from transdermal gels and patch across synthetic membrane/rabbit skin but was most significant when used in 3% concentration.	50
Olive oil	In a study on preparation and evaluation of olanzepine transdermal patches, the formulation containing 10% olive oil has enhanced permeation through rat skin.	51
Corn oil	Corn oil containing unsaturated fatty acids was found to be promising natural permeation enhancer for transdermal delivery of olanzapine with no skin irritation. The therapeutic effectiveness of the optimized transdermal system was confirmed by tranquilizing activity in rotarod and grip mice model.	52
Coconut oil and pistachio oil	Permeation of clotrimazole gel was enhanced by various combinations of coconut oil, pistachio oil and sodium lauryl sulphate that exhibited a more pronounced and promising effect through rat skin.	53

Miscellaneous:

Capsaicin: Capsaicin (trans-8-methyl-N-vinilyl-6-nonenamide) is an alkaloid derived from hot chili peppers, belonging to the genus Capsicum of the Solanaceae family It contain a ring at one end of a

long alkyl chain, with a log P value of 3.31. The penetration enhancing effects of capsaicin on naproxen was investigated with in vitro experiments employing full-thickness, female human skin and an *ex-vivo* perfused rabbit ear model. It was found that

capsaicin enhanced the permeation of naproxen through full-thickness skin approximately 2-fold⁵⁴.

Vitamin E: (α -tocopherol) enhanced the permeability coefficient of radio labeled hydrocortisone with an average enhancement factor of 1.81 through excised human cadaver skin. There was a reduction of lag time in skin samples treated with Vitamin E. This increase in the permeability of the stratum corneum can be due to the restricted insertion of Vitamin E in the ceramide-rich bilayer structure⁵⁵.

CONCLUSION: Transdermal drug delivery is effective over oral delivery of drugs and research is being carried out to overcome the skin barrier function. The penetration enhancer or sorption promoter functions by altering the barrier characteristics of skin membranes or by increasing the drug solubility inside the skin. Current review aims to summarize the natural sources that have been reported as promising transdermal penetration enhancers.

It can be concluded that research or exploration is beneficial for scaling up of the system of natural permeation enhancers and execute the manufacture of the dosage forms with natural permeation enhancers will play a prominent role in progress of useful and efficient transdermal products in future.

ACKNOWLEDGEMENT: Authors are highly grateful towards the management of the institute for providing a healthy environment for working.

CONFLICT OF INTEREST: Nil

REFERENCES:

1. Mathur *et al.*: Enhancers in transdermal drug delivery system, Asian Journal of Pharmaceutics. 2010; 173-183.
2. Masarrat B and Ahmed G: Chemical permeation enhancement through skin. International Journal of Advanced Research 2015; 3(8): 644-651.
3. Williams AC and Barry BW: Penetration enhancers. Adv Drug Deliv Rev. 2004; 56(5): 603-18.
4. Kanikkannan N, Kandimalla K, Lamba SS and Singh M: Structures activity relationship of chemical penetration enhancers in transdermal drug delivery. Current Medicinal Chemistry 1999; 6: 593-608.
5. Pavani P and Bhavani G: Formulation and evaluation studies on transdermal dosage forms of diclofenac sodium. World Journal of Pharmacy and Pharmaceutical Sciences, 4(3): 1043-1063.
6. Vyas S and Khar RK: Controlled drug delivery - concept and advances. Vallab Prakash. 2002; 418-22.
7. Vyas S and Khar RK: Controlled drug delivery - concept and advances. Vallab Prakash. 2002; 418-22.
8. Fox TL: Transdermal drug delivery enhancement by compounds of natural origin molecules. 2011; 10507-10540.
9. Singla V and Saini S: Penetration Enhancers: A novel strategy for enhancing transdermal drug delivery. International Research Journal of Pharmacy 2011; 2(12): 32-36.
10. Pavan P and Bhavani G: Formulation and evaluation studies on transdermal dosage forms of diclofenac sodium. World Journal of Pharmacy and Pharmaceutical Sciences. 4(3): 1043-1063.
11. Mathur V and Satrawala Y: Physical and chemical penetration enhancers in transdermal drug delivery system. Asian Journal of Pharmaceutics. 2010; 173-183.
12. Bavaskar K and Jain A: The impact of penetration enhancers on transdermal drug delivery system: Physical and chemical approach. International Journal of Pharma Research and Review. 2015; 4(7): 14-24.
13. Saini S: Recent development in Penetration Enhancers and techniques in transdermal drug delivery system. Journal of Advanced Pharmacy Education and Research. 2014; 4(1): 31-41.
14. Raut VS and Nemade SL: Chemical penetration enhancers: for transdermal drug delivery systems. International Journal of Pharmacy Review and Research. 2014; 4(1): 33-40.
15. Patel H: Transdermal drug delivery system as prominent dosage forms for the highly lipophilic drugs, International Journal of Pharmaceutical Research and Bio-science. 2012; 1(3): 42-65.
16. Patel H: Transdermal drug delivery system as prominent dosage forms for the highly lipophilic drugs, International Journal of Pharmaceutical Research and Bio-science. 2012; 1(3): 42-65.
17. Dhameecha D, Rathi AA; Drug vehicle based approaches of Penetration enhancement, International Journal of Pharma Sci. 2009, 1: 24-46
18. Dhameecha D and Rathi AA: Drug vehicle based approaches of Penetration enhancement, International Journal of Pharma Sci. 2009, 1: 24-46.
19. Sharma K and Mittal A: *Aloe vera* as penetration enhancer. International Journal of Drug Development and Research. 2015; 7 (1): 31-43.
20. Vashisth I and Ahad A: Investigating the potential of essential oils as penetration enhancer for transdermal losartan delivery: Effectiveness and mechanism of action. Asian journal of pharmaceutical sci. 2014; 9: 260- 267.
21. Fox LT and Gerber M: Transdermal drug delivery enhancement by compounds of natural origin, Molecules. 2011; 16: 10507-10540.
22. Chen J, Jiang QD, *et al.*: Potential of essential oils as penetration enhancers for transdermal administration of ibuprofen to treat dysmenorrhoea, Molecules 2015; 20: 18219-18236.
23. Fox LT and Gerber M: Transdermal drug delivery enhancement by compounds of natural origin, Molecules 2011; 16: 10507-10540.
24. Zhang LC and Jin-Hong HU: *In-vivo* and *in-vitro* evaluation of essential oils from Ligusticum chuanxiiong HORT on the transdermal delivery of flurbiprofen in rabbits, Biol. Pharm. Bull. 2006; 29(6): 1217-1222.
25. Rajan R and Vasudevan DT: Effect of permeation enhancers on the penetration mechanism of transfersomal gel of ketoconazole J Adv Pharm Technol Res. 2012; 3(2): 112-116.

26. Bao YYX: Enhancing effect of essential oils on the penetration of 5-fluorouracil through rat skin. 1996; 31(3): 214-21.
27. Jiang Q, Wu Y, Zhang H, Liu P, Yao J, Yao P, Chen J and Duan J: Development of essential oils as skin permeation enhancers: penetration enhancement effect and mechanism of action, *Pharmaceutical Biology* 2017; 55(1): 1592-1600.
28. Thacharodi D and Rao KP: Transdermal absorption of nifedipine from microemulsions of lipophilic skin penetration enhancers, *International Journal of Pharmaceutics*, 1994; 111(3): 235-240.
29. Zafar S and Ali A: Transdermal drug delivery of labetalol hydrochloride: Feasibility and effect of penetration enhancers, *Journal of Pharmacy and Bioallied Sciences*, 2010; 2(4): 321-324.
30. Huang YB, Wu PC, Ko HM and Tsai YH: Effect of pretreatment by cardamom oil on *in-vitro* percutaneous penetration of piroxicam gel. *Int J Pharm*, 1996; 131: 137-141.
31. Want J: Percutaneous penetration enhancement effect of essential oil of mint (*Mentha haplocalyx* Briq.) on Chinese herbal components with different lipophilicity. *Journal of Traditional Chinese Medical Sci.* 2014; 1(2): 109-119.
32. Vashisth I: Investigating the potential of essential oils as penetration enhancer for transdermal losartan delivery: Effectiveness and mechanism of action. *Asian journal of pharmaceutical sciences.* 2014;9: 260-267.
33. Indu Vashisth: Investigating the potential of essential oils as penetration enhancer for transdermal losartan delivery: Effectiveness and mechanism of action. *Asian Journal Of Pharmaceutical Sciences.* 2014; 9: 260-267.
34. Shamsheer AA and Charoo NA: Tulsi oil as a potential penetration enhancer for celecoxib transdermal gel formulations. 2014; 19(1): 21-30.
35. Das MK, Bhattacharya A and Ghosal SK: Effect of different terpene-containing essential oils on percutaneous absorption of trazodone hydrochloride through mouse epidermis. *Drug Delivery*, 2006; 13(6): 425-431.
36. Monti D, Chetoni P, Burgalassi S, Najarro M, Fabrizio Saettoni M and Boldrini: Effect of different terpene-containing essential oils on permeation of estradiol through hairless mouse skin. *Int. J. Pharm* 2002; 237: 209-214.
37. Aljabre SHM: Dermatological effects of *Nigella sativa*. *Journal of Dermatology and Dermatologic Surgery.* 2015; 19(2): 92-98.
38. Parney S and Dhurke RK: Effect of natural penetration enhancers on dermal delivery of hydrocortisone acetate. *Journal of Pharma. Investigation* 2014; 44(5): 365- 380.
39. Akbari J, Saeedi M and Farzin D: Transdermal absorption enhancing effect of the essential oil of *Rosmarinus officinalis* on percutaneous absorption of Na diclofenac from topical gel, *Pharmaceutical Biology*, 2015; 53(10): 1442-1447.
40. Fang JY, Leu Y and Hwang TL: Development of sesquiterpenes from *Alpinia oxyphylla* as novel skin permeation enhancers. *Eur. J. Pharm. Sci.* 2003; 19: 253-262.
41. Williams AC: *Transdermal and topical drug delivery.* London: Pharmaceutical Press; Edition 1st, 2003; 242.
42. Viljoen JM, Cowley A, Du Preez J, Gerber M and Du Plessis J: Penetration enhancing effects of selected natural oils utilized in topical dosage forms. *Drug Dev Ind Pharm.* 2015; 41(12): 2045-2054.
43. Zyl LV and Preez JD: Essential fatty acids as transdermal penetration enhancers. *Journal of Pharmaceutical Sciences* 2016; 105: 188-193.
44. Zyl LV and Preez JD: Essential fatty acids as transdermal penetration enhancers. *Journal of Pharmaceutical Sciences* 2016; 105: 188-193.
45. Puglia C and Bonina F: Effect of polyunsaturated fatty acids and some conventional penetration enhancers on transdermal delivery of atenolol, *Drug Delivery*, 2008; 15(2): 107-112.
46. Fang JY, Chiu HC, Wu JT and Chiang YR: Fatty acids in *Botryococcus braunii* accelerate topical delivery of flurbiprofen into and across skin. *Int. J. Pharm.* 2004; 276: 163-173.
47. Nashihata T, Kotera K, Nakano Y and Yamazaki M: Rat percutaneous transport of diclofenac and influence of hydrogenated soya phospholipids. *Chem. Pharm. Bull.* 1987; 35: 3807-3812.
48. Valjakka-Koskela R, Kirjavainen M and Monkkonen J: Enhancement of percutaneous absorption of naproxen by phospholipids. *Int. J. Pharm.* 1998; 175: 225-230.
49. Amit Kumar *et al.*: Comparison of vegetable and volatile oils as skin permeation enhancers for transdermal delivery of losartan potassium, *Der Pharmacia Lettre*, 2014; 6(1): 199-213
50. Hussain A, Khan GM and Shah SU: Development of a novel ketoprofen transdermal patch: effect of almond oil as penetration enhancers on *in-vitro* and *ex-vivo* penetration of ketoprofen through rabbit skin. *Pak Journal of Pharma Sciences*, 2012; 25(1): 227-232.
51. Sharma S, Aggarwal G and Dhawan S: Design and evaluation of olanzapine transdermal patches containing vegetable oils as permeation enhancers. *Scholars Research Library. Der Pharmacia Lettre*, 2010; 2(6): 84-98.
52. Aggarwal G, Dhawan S, HariKumar SL: Natural oils as skin permeation enhancers for transdermal delivery of olanzapine: *In-vitro* and *In-vivo* Evaluation. *Current Drug Delivery*. 2012; 9(2): 172-181.
53. Ubaid M, *et al.*: Formulation and *in-vitro* evaluation of carbopol 934-based modified clotrimazole gel for topical application. *Anais da Academia Brasileira de Ciências.* 2016; 88(4): 2303-2317.
54. Degim IT, Uslu A, Hadgraft J, Atay T, Akay C and Cevheroglu S: The effects of Azone® and capsaicin on the permeation of naproxen through human skin. *Int. J. Pharm* 1999; 179: 21-25.
55. Trivedi JS, Kril SL and For JJ: Vitamin E as a human skin penetration enhancer. *Eur. J. Pharm. Sci* 1995; 3: 241-243.

How to cite this article:

Kala S and Juyal D: Recent developments on natural transdermal penetration enhancers. *Int J Pharm Sci Res* 2018; 9(6): 2190-96. doi: 10.13040/IJPSR.0975-8232.9(6).2190-96.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)