



Received on 17 September, 2017; received in revised form, 29 November, 2017; accepted, 06 January, 2018; published 01 June, 2018

MORPHOLOGICAL CHANGES IN HUMAN RBCs DUE TO EXPOSURE TO SELECTED MEDICINAL PLANT LEAF EXTRACTS *IN-VITRO*

Meena Yadav*, G. Chaturvedi and N. Jha

Department of Zoology, Maitreyi College, Chanakyapuri, New Delhi - 110021, Delhi, India.

Keywords:

RBCs, Leaf extracts,
Tris-HCl, Blood groups,
Coagulation, Shrinking, Aggregation

Correspondence to Author:

Dr. Meena Yadav

Assistant Professor,
Department of Zoology,
Maitreyi College, University of
Delhi, Chanakyapuri, New Delhi
- 110021, Delhi, India.


E-mail: drmeena.yadav@gmail.com

ABSTRACT: The aim of this experiment was to find out the *in-vitro* effects of the extracts of certain plants, considered to have medicinal value, on the erythrocyte model. The plants, selected for the study, were Tulsi (*Ocimum tenuiflorum*), Neem (*Azadirachta indica*) and Amla (*Phyllanthus emblica*). Three different concentrations of leaf extracts were prepared using tris-HCl, namely 0.3 mg/ml, 0.15 mg/ml and 0.075 mg/ml and their effects on the RBCs in blood samples were studied using optical microscope. Various morphological changes like haemagglutination, shrinking, swelling, distortions, granulation of the membrane and bursting of the cells were observed. The degree of occurrence of a certain morphological change on a particular blood type varied with a variation in its concentration. Also, the same concentration of the extract showed different morphological changes with different blood groups indicating that the blood type is also important in the effects shown by the cell membranes. The crude leaf extracts of medicinal plants significantly affected the integrity and thus, the morphology of the RBCs.

INTRODUCTION: Plants are one of the most important sources of medicines in India. Today a large number of drugs in use are derived from plants like Tulsi (*Ocimum tenuiflorum*; Family-Lamiaceae), Neem (*Azadirachta indica*; Family-Meliaceae) and Amla (*Phyllanthus emblica*; Family- Phyllanthaceae). Scientists are targeting medicinal plants for research and development of alternative drugs, especially *Azadirachta indica*. It is one of the most promising medicinal plants whose extracts can act as anti-oxidant, anti-inflammatory, anti-bacterial, anti-fungal and anti-ulcer agents^{1, 2}. Several medicinal properties have also been attributed to *Ocimum sanctum* Linn.

Different parts of the plant like leaves, flowers, stems, roots, seeds etc. are known to possess therapeutic potentials and have been used, by traditional medical practitioners as expectorant, analgesic, anti-cancer, anti-asthmatic, anti-emetic, diaphoretic, anti-diabetic, anti-fertility, hypotensive, hepatoprotective, hypolipidmic and antistress agents^{3, 4}. Traditional Ayurvedic medicines use *Phyllanthus emblica* for treating diseases like scurvy, common cold, cancer and heart disease. *Phyllanthus emblica* is reported to have hypolipidemic, hypoglycemic activities. It also possesses good antimicrobial, anti-inflammatory and anticancer properties. *Phyllanthus emblica* has strong antioxidant property due to flavonoids and several gallic acid derivatives including epigallocatechingallate and tannins⁵.

The experiment was performed using the human RBCs due to their high density in blood (allowing observation of a large number of the cells in a single sampling) and lack of organellar structures

<p>QUICK RESPONSE CODE</p> 	<p>DOI: 10.13040/IJPSR.0975-8232.9(6).2405-10</p>
<p>Article can be accessed online on: www.ijpsr.com</p>	
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.9(6).2405-10</p>	

which would ensure that the changes occurring within the structure are those of the membrane alone. The human erythrocytes have a diameter of about 7-8 μm , they are biconcave in shape and lack cellular organelles like nucleus, mitochondria etc. and hence cannot replicate and divide on their own. The erythrocyte model has been widely used as it presents a direct indication of toxicity of injectable formulations as well as general indication of membrane toxicity. Another advantage of erythrocyte model is that blood is readily available and the cells are easy to isolate from the blood and its membrane resembles other cell membranes⁶. Thus, the suitability of the RBCs as a model and the inquisitiveness to Study the effects of medical plant extracts on them, led us to design the current experiment to study the various effects these extracts can have on the plasma membrane which can influence the homeostasis of the RBCs.

METHODOLOGY: The leaves of Tulsi (*Ocimum tenuiflorum*), Neem (*Azadirachta indica*) and Amla (*Phyllanthus emblica*), commonly found plants known to have medicinal values, were collected from the Maitreyi College premises and washed thoroughly with distilled water. The leaves were dried and the extract was prepared with tris-HCl (pH 6.8) with three different concentrations - 0.3 mg/ml, 0.15 mg/ml and 0.075 mg/ml. The different blood samples i.e. blood groups A⁺, B⁺, AB⁺, O⁺, A⁻ and B⁻, were incubated with equal volume of various extracts, individually, for 5 minutes and the observations were made. For examining the

changes in the morphology, the slides were observed using Nikon Eclipse E100 microscope at 400X. The same procedure was repeated with remaining medicinal plant extracts. The study was carried out from August, 2016 to April, 2017.

RESULTS: Various characteristic morphological changes were shown by the RBCs when treated with different medicinal plant leaf extracts having varied concentrations as shown in detail in **Table 1**, **2** and **3**. Treatment with *Ocimum tenuiflorum* extract showed maximum effect on blood group O⁺ followed by B⁺. The morphological changes included shrinking, granular surface and darkening of cell membrane. While these characteristics were shown by majority of the blood groups in all the strengths of extract, with varying degrees, some unique morphological changes were also observed like rupturing of cells (B⁺), aggregation (O⁺), distortion (A⁺, B⁺ and AB⁺: only in 0.3 mg/ml extract) and polygonal cells (AB⁺: 0.15 mg/ml).

For *Azadirachta indica* leaf extract (**Table 2**), blood group B⁺ was the most affected as it showed haemagglutination, distortion, shrinkage and granular surface while O⁺ was the least affected. Further, while haemagglutination was seen in all the blood groups studied, swelling was observed only in blood groups A⁻ and B⁻. As we decreased the concentration of the extract, the morphological changes were also lesser as compared to 0.3 mg/ml extract.

TABLE 1: MORPHOLOGICAL CHANGES IN HUMAN RBCs WITH VARIOUS STRENGTHS OF OCIMUM TENUIFLORUM LEAF EXTRACT

<i>Ocimum Tenuiflorum</i> leaf extract (0.3 mg/ml)	A ⁺	B ⁺	AB ⁺	O ⁺	A ⁻	B ⁻
Granular Surface		+	+	+	+	
Darkening cell membrane			+			
Shrinking	+	+		+	Few	Few
Swelling	+					
Smooth surface	+	+			+	+
<i>Ocimum Tenuiflorum</i> leaf extract (0.15 mg/ml)						
Granular surface		+		+	Rare	
Darkening of the cell membrane				+		
Shrinking	+	+	+	+	+	+
Swelling				+		
Smooth surface	+	+	+	+	+	+
<i>Ocimum Tenuiflorum</i> leaf extract (0.075 mg/ml)						
Granular surface		+	+	+	+	+
Darkening of cell membrane				+		
Shrinking		+	+	+	+	
Swelling	+					
Smooth surface	+	+	+		+	+

TABLE 2: MORPHOLOGICAL CHANGES IN HUMAN RBCs WITH VARIOUS STRENGTHS OF AZADIRACHTA INDICA LEAF EXTRACT

<i>Azadirachta indica</i> leaf extract (0.3 mg/ml)	A ⁺	B ⁺	AB ⁺	O ⁺	A ⁻	B ⁻
Haemagglutination (highly packed)	+	+	+	+	+	Weak
Granular surface		+				
Shrinking		+	+		+	
Swelling					+	+
Distortion	+	+				+
<i>Azadirachta indica</i> leaf extract (0.15 mg/ml)						
Haemagglutination			Lesser	Lesser		Lesser
Granular surface	+	+	+		+	
Shrinking	+	+	+		+	
Swelling						+
Distortions	+	+				+
<i>Azadirachta indica</i> leaf extract (0.075 mg/ml)						
Haemagglutination		Lesser	Lesser	Lesser		Lesser
Granular surface		+	+		+	
Shrinking	Lesser	Lesser	Lesser			
Swelling					+	
Distortions	Lesser	Lesser				Lesser

TABLE 3: MORPHOLOGICAL CHANGES IN HUMAN RBCs WITH VARIOUS STRENGTHS OF PHYLLANTHUS EMBLICA LEAF EXTRACT

<i>Phyllanthus emblica</i> leaf extract (0.3 mg/ml)	A ⁺	B ⁺	AB ⁺	O ⁺	A ⁻	B ⁻
Haemagglutination	+	+	+	+	+	+
Granular surface		+				
Shrinking					+	
Swelling				+		
Deformities		Major				+
Hemolysis				+		
Clumps of cells turning dark brown			+			
Polygonal cells	+					
<i>Phyllanthus emblica</i> leaf extract (0.15 mg/ml)						
Haemagglutination	Lesser	Lesser		Lesser		
Granular surface		+				
Shrinking			+			
Distortion	+	+		+		+
Brownish colour of cells				+		
Smooth surface					+	
<i>Phyllanthus emblica</i> leaf extract (0.075 mg/ml)						
Haemagglutination	Lesser		Lesser	Lesser		Lesser
Granular surface		+		+		
Shrinking			Faster than 50 %			
Swelling				+		
Distortions (less than 50%)	Lesser	Lesser				

Other than these changes, some other changes in morphology included change of color of aggregates from yellow to brown (AB⁺ and O⁺: 0.3 mg/ml and 0.15 mg/ml) and polygonal shape of RBCs (AB⁺: 0.075 mg/ml). Further, with *Phyllanthus emblica* leaf extract, the blood group O⁺ was the most affected while A⁻ and B⁻ were least affected (**Table 3**). Haemagglutination (**Fig. 1**) was a common feature in all the blood groups studied while granular surface of RBCs (**Fig. 2**) was seen only in blood group B⁺.

Shrinking was seen in blood groups A⁻ and AB⁺, swelling in only O⁺ and major distortions in blood groups B⁺ and B⁻.

The other changes observed are: aggregates turned brown (AB⁺ and O⁺: 0.3 and 0.15 mg/ml), smooth surface (A⁻: 0.15 and 0.075 mg/ml), bursting of cells (O⁺: 0.3 mg/ml) and polygonal cells (A⁺: 0.3 mg/ml). The respective control experiments showed no effects on the membrane of the erythrocytes.




FIG. 1: A) AND B) HAEMAGGLUTINATION (x40)




FIG. 2: GRANULAR CELL SURFACE (x40)

DISCUSSION: Our study revealed that varying concentrations of the leaf extracts, of selected medicinal plants, had varying effects on the morphology of the erythrocyte membrane. Darkening of the RBC membrane was observed in the blood group B⁺ when treated with *Ocimum tenuiflorum* extract (0.3 mg/ml). 0.15 mg/ml and 0.075 mg/ml concentrations of the same extract showed endosmosis leading to swelling and exosmosis leading to shrinking of the RBCs in blood group B⁺. Haemagglutination (Fig. 1) was evident in blood groups AB⁺, O⁺ and A⁺ when treated with different concentrations of the *Phyllanthus emblica* extract. With decrease in concentrations of *Phyllanthus emblica* extract, clump formation decreased.

The results show significant changes in the morphology of the RBCs when exposed to various strengths of medicinal plants studied. The homeostasis and integrity of the cell membrane depends on the immediate environmental conditions in which the cell is placed including the tonicity of the medium and other substances which may interfere with the cell membrane. It was interesting

to know that the extracts of plants, considered to have medicinal value, significantly influenced the integrity of the cell membrane, thereby affecting

the RBCs on the whole *in-vitro*. The RBCs in different blood groups also showed differences in the morphological changes. This observation may be attributed to the molecules on the cell surface of RBCs which may be directly or indirectly involved in the alteration in the functioning of the ion channels or other mechanisms involved in maintaining the osmotic balance. Similar results were observed when RBCs were exposed to extract of *Ricinus communis* L., where the extract proved to be toxic to cells and modified the cell membrane structure⁷ and exposure to extract of *Combretum zeyheri* caused hemolysis in sheep erythrocytes⁸.

Another study showed that treatment of blood cells with extract of *Paullinia cupana* results in the altered morphology of RBCs⁹. The aqueous extract of *Matricaria recutita* also alters the RBC morphology¹⁰. Similarly the interaction between the drugs and plasma membrane has been studied by different techniques previously^{11, 12}. Thus, the morphological analysis of the RBC membrane is an important method available to study how the natural products affect the RBC membrane^{9, 13, 14, 15} and it is the standard protocol to see the effect of natural or synthetic drugs on the cell membrane¹⁶.

The above explained changes in the morphology of the RBCs like breaking of cell membrane,

fragmentation, hemolysis and hemagglutination, may be attributed to the various chemical components present in the medicinal plant extracts studied as the RBCs when placed in the tris-HCl and even the physiological saline don't show any changes in the nature of the plasma membrane¹⁷. Thus, these changes may be due to the interaction of the plant extract components with the ion transport channels or osmotic transport system present in the plasma membrane of RBCs.

Further, the study by Dkhil et al.,¹⁸ has shown that the leaf extract of *Azadirachta indica* contains high amount of flavonoids like rutin, quercetin, flavono glycosides, polyphenols and tannins which have antioxidant and anti-inflammatory properties. However, the protective activity of *Azadirachta indica* extract is dose dependent with the most effective dose being 1000 and 2000 mg/kg for renal damage in mice, produced by *Plasmodium berghei*². A similar study has reported that human erythrocytes undergo hydrolysis in presence of extracts from *Daphne gnidium*, *Tamarixa phylla* and *Morettia canescens*¹⁹. It has been reported that the methanol extract from the leaves of *Daphne gnidium* has antileukemic, antibacterial, antimycotic activities²⁰ and *Tamarixa phylla* also has antimicrobial as well as phytochemical activities and are traditionally used to evacuate coagulated blood and treat hematomas^{19, 21}.

The RBC membrane fluidity, permeability and its functions depend on the levels of cholesterol and phospholipid. In medical conditions where the ratio between cholesterol and phospholipid is altered leading to malfunctioning of the RBC membrane, the application of *P. atlantica* extract helps in normalizing the RBC membrane structure²². The papaya leaf extracts have the capability of reducing the hemolysis *in-vitro* and hence, are a potent alternative for treating the diseases which result in altering the membrane structure²³. Thus, the medicinal plant extracts have been traditionally used in medicine all over the world but for the use of these extracts, dose should be precisely formulated, else they can prove toxic to the cells as has been seen in several studies^{24, 25, 26, 27}.

CONCLUSION: To summarize, the study supports the belief that the plant extracts of

Azadirachta indica, *Phyllanthus emblica* and *Ocimum tenuiflorum* do have potent influence on the RBC plasma membrane, may be by influencing ion channels or by altering the integrity of the cell membrane. Since, they have been known since ancient times to have medicinal value, these plant extracts may be used to treat medical conditions or minor health issues by modulating the concentrations of the active components in these plant extracts. For this, the knowledge of the actual mediators in the plant extracts needs to be elucidated and the correct dose needs to be formulated to maximize the use of these plant extracts, for medicinal purposes, by more and more people.

ACKNOWLEDGEMENT: The authors are thankful to our Principal and also to the funding agency Department of Biotechnology, for providing necessary support under its Star College Scheme.

CONFLICT OF INTEREST: Nil

REFERENCES:

1. Alzohairy MA: Therapeutics role of *Azadirachta indica* (Neem) and their active constituents in disease prevention and treatment. Evid Based Complement Alternat Med 2016. doi: 10.1155/2016/7382506
2. Somsak V, Chachiyo S, Jaihan U and Nakinchat S: Protective effect of aqueous crude extract of neem (*Azadirachta indica*) leaves on *Plasmodium berghei*-induced renal damage in mice. Journal of Tropical Medicine 2015; Article ID 961205, 5.
3. Cohen MM: Tulsi *Ocimum sanctum*: A herb for all reasons. J Ayurveda Integr Med 2014; 5(4): 251-9.
4. Prakash P and Gupta N: Therapeutic uses of *Ocimum sanctum* linn (Tulsi) with a note on eugenol and its pharmacological actions: a short review. Indian J Physiol Pharmacol 2005; 49(2): 125-31.
5. Ashadevi S, Benny B and Sherin MS: Protection of DNA and erythrocytes from free radical induced oxidative damage by methanolic extract of amla (*Phyllanthus emblica*). Int. J. Pharm. Sci. Rev. Res. 2014; 28(2): 38-42.
6. Noudeh GD, Sharififar F, Khatib M, Behravan E and Afzadi MA: Study of aqueous extract of three medicinal plants on cell membrane-permeabilizing and their surface properties. African J Biotech 2009; 9(1): 110-16.
7. Mousinho KC, Correia MBL, da Silva JO, Magnata SSLP, de Souza IA and Catanho MTJA: Effect of the extract of *Ricinus communis* L. on the osmotic fragility, labeling of red blood cells with Technetium-99m and morphology of the cells. Braz. Arch. Biol. Technol. 2008; 51(6). <http://dx.doi.org/10.1590/S1516-89132008000600008>
8. Mapfunde S, Sithole S and Mukanganyama: *In-vitro* toxicity determination of antifungal constituents from *Combretum zeyheri*. BMC Complementary Alternative Med, 2016; 16: 162. <https://doi.org/10.1186/s12906-016-1150-9>

9. de Oliveira JF, Avila AS, Braga AC, de Oliveira MB, Boasquevisque EM, Jales RL, Cardoso VN and Bernardo-Filho M: Effect of extract of medicinal plants on the labeling of blood elements with Technetium-99m and on the morphology of red blood cells: I-a study with *Paullinia cupana*. *Filoterapia* 2002; 73(4): 305-12.
10. Garcia-pinto AB, Santos-Filho SD, Carvalho JJ, Pereira MJS, Fonseca AS and Bernardo-Filho M: *In-vitro* and *in-vivo* studies of an aqueous extract of *Matricaria recutita* (German chamomile) on the radiolabeling of blood constituents, on the morphology of red blood cells and on the biodistribution of the radiopharmaceutical sodium pertechnetate. *Pharmacogn Mag* 2013; 9(1): S49-S56.
11. Mineo H, Kasai K, Makihara R and Yuuki T: Effects of benzoic acid and its analogues on osmotic fragility in guinea pig erythrocytes *in-vitro*. *J Membr Sci and Technol* 2016; 6: 4. DOI: 10.4172/2155-9589.1000161.
12. Laboni FR, Afsari F, Howlader MSI, Labu ZK and Julie AS: Thrombolytic and membrane stabilizing activities of ethanolic extract of local medicinal plant *Murraya paniculata*. (Family: Rutaceae). *J Pharmacogn and Phytoch* 2015; 4(2): 17-20.
13. Sinha A, Chu TTT, Dao M and Chandramoandas R: Single-cell evaluation of red blood cell bio-mechanical and nano-structural alterations upon chemically induced oxidative stress. *Sci Rep* 2015; 5: 9768. doi: 10.1038/srep09768.
14. Tomaiuolo G: Biomechanical properties of red blood cells in health and disease towards microfluidics. *Biomicrofluidics* 2014; 8(5): 051501. doi: 10.1063/1.4895755.
15. Braga ACS, Gomes ML, Santos JS, Oliveira JF, Machado EF, Oliveira MP, Filho SDS and Bernardo-Filho M: Alteration of the labeling of blood constituents with technetium-99m and the morphology of red blood cells by *Baccharis trimera* extract. *African Journal of Pharmacy and Pharmacology* 2012; 6(4): 228-34.
16. Presta GA, Fonseca AS and Bernardo-Filho M: *A Chrysobalanus icaco* extract alters the plasmid topology and the effects of stannous chloride on the DNA of plasmids. *Rev Bras Farmacogn* 2007; 17: 331-35.
17. Meena Y, Romsha S, Sneha Y, Yadav S and Dhiman S: Effect of various solute concentrations on the membrane permeability of RBCs in human blood. *Journal of Cell and Tissue Research* 2015; 15(2): 4943-7.
18. Dkhil MA, Al-Quraishy S, Abdel Moneim AE and Delic D: Protective effect of *Azadirachta indica* extract against *Eimeria papillata* induced coccidiosis. *Parasitology Research* 2013; 112(1): 101-6.
19. Zohra M and Fawzia A: Hemolytic activity of different herbal extracts used in Algeria. *International Journal of Pharma Sciences and Research (IJPSR)* 2014; 5(8): 495-500.
20. Deiana M, Rosa A, Casu V, Cottiglia F, Bonsignore L and Dessi MA: Chemical composition and antioxidant activity of extracts from *D. gnidium*. *JAOCS* 2003; 80: 65-70.
21. Alrumman AA: Phytochemical and antimicrobial properties of *Tamarix phylla* L. leaves growing naturally in the Abha region, Saudi Arabia. *Arabian J Sci and Engg* 2016; 41(6): 2123-29.
22. Tolooei M and Mirzaei A: Effects of *Pistacia atlantica* extract on erythrocyte membrane rigidity, oxidative stress, and hepatotoxicity induced by CCl4 in rats. *Global Journal of Health sciences* 2015; 7(7): 32-8.
23. Ranasinghe P, Ranasinghe P, Abeysekera WPK, Premakumara GAS, Perera YS, Gurugama P and Gunatilake SB: *In-vitro* erythrocyte membrane stabilization properties of *Carica papaya* L. leaf extracts. *Pharmacognosy Res.* 2012; 4(4): 196-202.
24. Lin H, Guo X, Zhang S, Dial SL, Guo L, Manjanatha MG, Moore MM and Mei N: Mechanistic evaluation of *Ginkgo biloba* leaf extract-induced genotoxicity in L5178Y cells. *Toxicological Sciences* 2014; 139(2): 338-49.
25. Adu-Amoah L, Agyare C, Kisseih E, Ayande PG and Mensah KB: Toxicity assessment of *Erythrophleum ivorense* and *Parquetina nigrescens*. *Toxicology reports* 2014; 1: 411-20.
26. Ayodele AA, Adetayo ON, Osita SO, Ali HB, Oladimeji Ida and Amin AH: Red blood cell membrane stabilizing potentials and phytochemical assessment of water extract thylakoids of *Sennatoria Linn* in natural population of Northeastern Nigeria. *International Journal of Herbal Medicine* 2015; 2(6): 3-6.
27. Furman AEF, Henneberg R, Hermann PB, Leonart MSS and Nascimento AJ: *Ginkgo biloba* extract (EGB 761) attenuates oxidative stress induction in erythrocytes of sickle cell disease patients. *Braz. J. Pharm. Sci* 2012; 48(4).
<http://dx.doi.org/10.1590/S1984-82502012000400009>

How to cite this article:

Yadav M, Chaturvedi G and Jha N: Morphological changes in human RBCs due to exposure to selected medicinal plant leaf extracts *in-vitro*. *Int J Pharm Sci Res* 2018; 9(6): 2405-10. doi: 10.13040/IJPSR.0975-8232.9(6).2405-10.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)