



Received on 28 September, 2017; received in revised form, 01 June, 2018; accepted, 07 June, 2018; published 01 July, 2018

BACTERICIDAL SCREENING AND ESTIMATION OF FREE RADICAL SCAVENGING ACTIVITY OF FRUTICOSE LICHEN *CLADONIA SCABRIUSCULA* DELISE (NYL.) FROM GOPESHWAR, UTTARAKHAND, INDIA

Anand Pandey ^{*1}, Afifa Qidwai ¹, Madhu Pandey ¹, Rajesh Kumar ², Shashi Kant Shukla ² and Anupam Dikshit ¹

Biological Product Lab, Department of Botany ¹, Centre of Rural Technology and Development ², University of Allahabad, Allahabad - 211002, Uttar Pradesh, India.

Keywords:

Mycobiont,
Lichen, *Cladonia*,
CLSI, FRSA, Antioxidant, etc.

Correspondence to Author:

Dr. Anand Pandey

CSIR- Research Associate,
Biological Product Lab, Department
of Botany, University of Allahabad,
Allahabad - 211002, Uttar Pradesh,
India.

E-mail: deep.7890@gmail.com

ABSTRACT: Lichens came as a consortium of two distinct and dissimilar components- the mycobiont and the phycobionts, which were defined as a composite organism. Lichen vegetation in India is dominated by more than 2500 species. Lichen secondary metabolites specially the phenolics, depsides and depsidones are unique in their origin and are found nowhere else. Much attention has been paid on the biological roles of lichen secondary substances in the recent past; which have been found to possess a lot of bioactivities, such as anti-tumor, antibacterial, antifungal, antiviral, anti-inflammatory and antioxidant activities. The present study aims on the bactericidal activity of lichen, *Cladonia scabriuscula*, collected from the Gopeshwar, Uttarakhand in the Western Himalayas. The 50% v/v ethanolic extract (CSE) extracted through cold extraction method was tested for its bactericidal potential against prevalent water borne pathogen and its antioxidant potential through CLSI protocol. The CSE was found active against *E. coli* and *S. typhimurium* with MIC 2.48 and 1.02 mg/ml respectively; while the FRSA IC₅₀ was 0.625 mg/ml.

INTRODUCTION: In the course of evolution on Earth, the lichen came as a result of mutual inter-relationship of algal and fungal cells. Thus, lichens can be defined as composite organism consisting of two distinct and dissimilar components- the mycobiont and the phycobionts. There are more than 20,000 known lichen taxa. Since the fungal constituent usually dominates the association, lichens traditionally have been considered as a lifestyle of fungi ¹.

Erik Acharius, who was the last student of Swedish naturalist, Carolus Linnaeus (1707- 1778), and is acclaimed “The father of Lichenology”, a Physician by profession, studied the specimens collected by Linnaeus and created many new genera and documented his work in Lichenographiae suecicae prodromus (1798); Methodus lichenum (1803); Lichenographia universalis (1810) and Synopsis methodica Lichenum (1814).

Lichens synthesize numerous metabolites, the “lichen substances,” which comprise amino acid derivatives, sugar alcohols, aliphatic acids, macrocyclic lactones, mono - cyclic aromatic compounds, quinones, chromones, xanthones, dibenzofuranes, depsides, depsidones, depsones, terpenoids, steroids, carotenoids and diphenyl ethers ^{2,3}.

<p>QUICK RESPONSE CODE</p>	<p>DOI: 10.13040/IJPSR.0975-8232.9(7).2847-51</p> <hr/> <p>Article can be accessed online on: www.ijpsr.com</p>
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.9(7).2847-51</p>	

In the recent past, much attention has been paid on the biological roles of lichen secondary substances; which have been found to have a lot of bioactivities, such as anti-tumor, antibacterial, antifungal, antiviral, anti-inflammatory and also antioxidant activities⁴. In the folklore of many European countries, lichens were used as a remedy for pulmonary tuberculosis and in the treatment of wounds and skin disorders. Burkholder reported for the first time the presence of antibiotic substances in lichens⁵. Several lichen metabolites were found to be active against Gram-positive organisms⁶. The anti-mycobacterial activity of lichen compounds was reported against non-tubercular species of *Mycobacterium*⁷. Lichen extracts have cytotoxic activity in different degrees. The aqueous extract of *P. polydactyla* and the ethanol extract of the *R. farinacea* exhibited potent antibacterial activities⁸.

On the other hand, the global environment is day by day getting polluted and unhygienic for humans, by anthropogenic activities. The vital life resource, water has become so polluted, that getting potable water from natural sources is now literally not possible. Waterborne bacterial pathogens, such as *Escherichia coli*, *Vibrio cholerae*, *Salmonella typhimurium*, *Klebsiella pneumoniae*, *Shigella dysenteriae*, etc. have possessed threat to the life of many people residing in the villages, slums, chawls, etc⁹. Reports on the antibacterial activity of the medicinal plants have flooded the scientific arena^{10, 11, 12, 13, 14}. Current study focuses on the antibacterial activity of lichen, *Cladonia scabriuscula* against prevalent water borne pathogen and its antioxidant potential.

MATERIALS AND METHODS:

a. Collection of Lichens and Extraction of Metabolites: The lichen *Cladonia scabriuscula*

(Delise) Nyl. is widely distributed in all temperate regions of Uttarakhand, Sikkim, Tamil Nadu, Nagaland and West Bengal. The specimen for the current study was collected from Gopeshwar, Uttarakhand; on the way to Badrinath, at an altitude of 1500 - 1700 meters. The specimen was identified by Dr. D.K. Upreti, Chief Scientist, Lichenology Laboratory, NBRI, Lucknow. The voucher specimen for LWG Herbarium was LWG-014583. The thallus primarily is squamulose, 1-3 mm X 1-2 mm dimensions. Podetia up to 3-8 cm tall, 1-2 mm in diameter. Colour of the thallus is greyish to greenish white, corticated podetia with scaling off to form squamules of the thallus. Apothecia are present, brown coloured, on lateral branches and at the apices **Fig. 1a**.

Fumeroprotetraric acid is the main compound present in the thallus. The extraction of the metabolite was done by soaking 5 gm of lichen thallus in 50 ml 50% v/v Ethanol (CSE), overnight. The filtrate obtained by Whatman no. 1 filter paper was condensed in Rotary Evaporator (BUCHI). Further, the crude extract dried, weighed and kept at 4 °C in refrigerator. Percentage yield was obtained as follows:

$$\% \text{ age yield} = (\text{weight of the lichen thallus}) / (\text{weight of the crude extract}) \times 100$$

b. Procurement and Maintenance of Bacterial Cultures: All the water-borne bacterial pathogens viz., *E. coli* (MTCC- 723), *V. cholera* (MTCC-3906), *S. typhimurium* (MTCC- 3231) and *K. pneumoniae* (MTCC- 4032) were procured from microbial type culture collection (MTCC, IMTECH), Chandigarh **Fig. 1b**. The culture slants were maintained in Muller Hinton Agar medium at 35 ± 2 °C.




FIG. 1A: LICHEN CLADONIA SCABRIUSCULA; B: SLANTS OF BACTERIAL CULTURES- K. PNEUMONIAE, S. TYPHIMURIUM, V. CHOLERAEE AND E. COLI

c. Antimicrobial Assay: The susceptibility of the water borne pathogen against the CSE was performed by globally accepted NCCLS, now CLSI (Clinical and Laboratory Standards Institute), recommended broth micro-dilution method¹⁵. Briefly, in a 96 well microtitre plate, a series of dilution was designed for CSE was made in 4th to 11th well in duplicate, to assess the inhibitory action against the pathogen. To minimize, the effect of broth medium and experimental drug, a media control well in 2nd column and 3rd column wells respectively. A negative and positive growth control well was also assigned to 1st and 12th column wells, as well **Fig. 2**. Results were obtained in form of 50% age Inhibitory Concentration (IC₅₀) and Minimum Inhibitory Concentration (MIC). Stock solution (50mg/ml) of CSE extract, was prepared in DMSO. In brief, the initial bacterial inoculum suspension was prepared as per 0.5 McFarland standards (corresponding to a CFU of 1.5×10⁷ cell/ml). The MIC and IC₅₀ mg/ml values were obtained spectrophotometrically by (SpectraMax Plus384, Molecular Devices Corporation, USA) at 492 nm, after an incubation of 24 h at 35 ± 2 °C.




FIG. 2: SCHEMATIC REPRESENTATION OF CLSI-BROTH MICRODILUTION BACTERIAL SUSCEPTIBILITY PROTOCOL

d. Free Radical Scavenging Activity: DPPH assay- The lichen extract was also tested for its free radical scavenging activity (FRSA) by DPPH assay protocol¹⁶. DPPH (2, 2- Diphenylpicryl-1- picryl-

hydrazyl) produces free radical in the solution when dissolved in methanol. The reducing power of the lichen extract is thus examined by this assay. The CET from *Cladonia scabriuscula*, was tested with Ascorbic acid (Vitamin C), taken as Standard. Result was expressed in form of IC₅₀ obtained from the percent inhibition of free radical as FRSA.

RESULTS: Results were expressed in form of IC₅₀ and MIC (mg/ml), based on the turbidimetric results obtained in form of optical density and statistically analyzed by SOFTMAX PRO software- Spectramax Plus 384, Molecular Devices, USA. The CSE, was found active against *Escherichia coli*, *Salmonella typhimurium*, *Vibrio cholerae*, *Klebsiella pneumoniae* was depicted in the form of IC₅₀ and MIC values **Table 1** and **Fig. 3**.




FIG. 3: PERCENT INHIBITION (MIC AND IC₅₀) OF WATER BORNE BACTERIAL PATHOGEN BY CSE (©SOFTMAX PRO)

Appreciable degree of inhibition was found active against *E. coli* and *S. typhimurium* with IC₅₀; 0.35 and 0.29 mg/ml and MIC; 2.48 and 1.02 mg/ml respectively. While only IC₅₀ was obtained against *V. cholerae* and *K. pneumoniae* viz., 0.59 and 0.61 mg/ml respectively.

The FRSA in form of IC₅₀ for CSE was found at 0.625 mg/ml while the Standard- Ascorbic acid has an IC₅₀ at 0.02 mg/ml **Table 2; Fig. 4**.

TABLE 1: ANTIBACTERIAL ACTIVITY OF CSE (CLADONIA SCABRIUSCULA 50% v/v ETHANOL) EXTRACT

Name of the lichen	Activity against tested pathogens	Antibacterial activity (mg/ml)	
		IC ₅₀	MIC
50% EtOH <i>Cladonia scabriuscula</i>	<i>Escherichia coli</i>	0.348	2.476
	<i>Vibrio cholerae</i>	0.587	Range
	<i>Klebsiella pneumoniae</i>	0.612	Range
	<i>Salmonella typhimurium</i>	0.287	1.015

TABLE 2: PERCENT DPPH RADICAL REDUCTION AT DIFFERENT CONCENTRATION (mg/ml)

Name of lichen	Percent DPPH radical reduction at different concentration (mg/ml)								
	0.02	0.039	0.078	0.157	0.316	0.625	1.25	2.5	
Std. (Ascorbic acid)	91.08	91.22	91.48	91.43	91.56	91.46	91.77	91.22	
<i>Cladonia scabriuscula</i>	-12.65	-11.45	7.67	27.67	47.97	74.56	87.87	89.27	




FIG. 4: FREE RADICAL SCAVENGING ACTIVITY (FRSA) FOR CLADONIA SCABRIUSCULA BY DPPH ASSAY

DISCUSSION: With the ever changing world, the microorganisms are also evolving day by day. As the time passes on, pathogenic microbes develop resistance against several classes of antibiotic. At this stage several medicinal plants have shown promising results in the treatment of water borne diseases like diarrhea, dysentery, etc. caused by bacteria. Above results clearly indicate that, CSE was highly active against *E. coli* and *S. typhimurium*. Ethanol helps to maximize the extraction of effective phenol acids, here fumaroprotectoric acid. Variability in the MIC may be attributed to the phylogenetic position of the various pathogens¹⁷. The herbal extracts have no side effects as compared to the synthetic drugs and thus are preferably a better alternate. Lichen metabolites are unique and have high degree of antimicrobial potential as compared to other herbals.

Lichen *Cladia aggregata* was tested for its chemical characterization and antimicrobial activity of barbatic acid and was found active against *S. aureus*¹⁸. Effect of several herbal plant based medicine are in practice for the development of many herbal drug in a market for consumption purposes. With the range of these views and several properties of the *Tridax procumbens* L. as well as *Lippia nodiflora* L. were tested and showed antibacterial activity against both gram-positive and gram-negative bacteria^{9,10}.

The phenolic compounds and their derivatives in lichen have been proved to be detrimental for pathogenic microbial fauna. These substances generally acidify the microbial cell wall and consequently, cause cytoplasm membrane rupture, inactivate or immobilize the enzymes, and interfere with physiological functions such as electrons

transport, oxidative phosphorylation etc.^{19, 20, 21}. In a study, on antimicrobial activity of atranorin, it was elucidated that it showed high bactericidal activity values ranging from 5 µg to 70.7 µg/ml and this value also varied from genus to genus, as in case of *Cladonia foliacea* (15.6µg to 500µg/ml)²². While, another finding based on the cultured lichens, showed mycobiont and photobiont culture of *U. ghattensis* and *A. awasthii* have high antioxidative and antibacterial potential²³. In the current study also, the lichen CSE has an FRSA count of 0.625 mg/ml (IC₅₀), which is a value addition to its bioprospective attributes. Reports on concentration dependant antioxidative activity of lichen metabolites; salazinic, sekikaic and usnic acid. Salazinic acid showed 17.2 to 20.5, sekikaic acid 13.7 to 17.4, and usnic acid 18.8 to 25.5 as IC₅₀ (µg/ml), have also been observed, which indicated lichens as comparatively better natural antioxidant and a good replacement of synthetics²⁴.

Many path-breaking studies have come recently, focusing on the drug loading capacity of herbal compounds derived from lichens as well their capacity to fabricate nanoparticles. These novel antibacterial compounds have potentiality to not only kill the bacteria but also can be used as an effective drug loader or carrier^{25, 26, 27, 28, 29}.

CONCLUSION: It is clearly evident from the above results that fruticose lichen, *Cladonia scabriuscula* 50% v/v ethanolic extract (CSE) has marked antibacterial potential against a wide range of water borne bacterial pathogens, which possess threat to health of human beings in diverse localities, where potable water is still a big problem. Its antioxidant potential is a value addition to its bioprospective values. Moreover, bioprospection should not be limited to mere exploration of the novel antimicrobial agents but should lead formulation development. Thus, present finding strongly supports CSE as a possible alternate of synthetic therapeutic agents after organoleptic analysis, currently in progress.

ACKNOWLEDGEMENT: Thanks are due to the Head of Department of Botany, University of Allahabad for laboratory facilities, to Dr. D.K. Upreti, Chief Scientist, NBRI for lichen identification and CSIR, for financial assistance.

CONFLICT OF INTEREST: This work is a sole outcome of experiments done at Biological Product lab., and authors have no conflict of interest for this work.

REFERENCES:

- Feuerer T and Hawksworth DL: Biodiversity of lichens, including a world-wide analysis of checklist data based on Takhtajan's floristic regions. *Bio Conserv* 2007; 16: 85-98.
- Clix JA, Whitton AA, and Sargent MV: Recent progress in the chemistry of lichen substances. *Prog. Chem. Org. Nat. Prod.* 1984; 45: 207-211.
- Fiedler P, Gambaro V, Garbarino JA and Quihot W: Epiphorellic acids 1 and 2, two diaryl ethers from the lichen *C. epiphorella*. *Phytochemistry* 1986; 25: 461-465.
- Oksanen I: Ecological and biotechnological aspects of lichens. *Appl Microbiol Biotechnol* 2006; 73: 723-734.
- Burkholder PR, Evans AW, McVeigh I and Thornton HK: Antibiotic activity of Lichens. *Proceedings on National Academic Science, USA* 1944; 30(9): 250-255.
- Lauterwein M, Oethinger M, Belsner K, Peters T, and Marre R: *In vitro* activities of the lichen secondary metabolites vulpinic acid, (+)-usnic acid and (-)-usnic acid against aerobic and anaerobic microorganisms. *Antimicrob Agents Chemother* 1995; 39: 2541-2543.
- Ingoldsdottir K, Chung GA, Skulason VG, Gissurarson SR and Vilhelmsdottir M: Antimycobacterial activity of lichen metabolites *in vitro*. *Europ J Pharm Sci*; 1998; 6: 141-144.
- Karagöz A, Arda N, Gören N, Nagata K, and Kuru A: Antiviral activity of *Sanicula europaea* L. extracts on multiplication of human parainfluenza virus type. *Phytotherapy Research* 1999; 13(5): 436-438.
- Pandey M, Pandey A, Shukla SK, Kumar R, Pathak A, Mishra RK and Dikshit A: A comparative analysis of *in vitro* growth inhibition of water-borne bacteria with bioactive plant *Lippia nodiflora* L. and Camphor. *Desalination and Water Treatment* 2016a; 1-7.
- Pandey M, Pandey A, Kumar R, Pathak A and Dikshit A: A comparative antimicrobial analysis of *T. procumbens* L. various extracts on waterborne bacterial pathogens. *International Current Pharmaceutical J* 2016b; 5(3): 22-26.
- Kumar A, Mishra RK, Srivastava S, Tiwari AK, Pandey A, Shukla AC and Dikshit A: Antibacterial activity of *Trachyspermum ammi* L. essential oil with special reference to phylogenetic analysis of water borne bacterial pathogens. *Adv. in Env. Biology* 2011; 5(6): 1271-1278.
- Pandey M, Qidwai A, Pandey A, Shukla SK, Kumar R, Pathak A and Dikshit A: Investigation for *in vitro* antibacterial activity of *Citrus sinensis* L. against *Propionibacterium acnes*. *International Journal of Current Advanced Research* 2017; 6(8): 2341-2345.
- Qidwai A, Pandey M, Shukla SK, Kumar R, Pandey A and Dikshit A: Antibacterial activity of *Mentha piperita* and *Citrus limetta* against *Propionibacterium acnes* (Anaerobic Bacteria). *International Journal of Pharma-ceutical Sciences and Research* 2016; 7(7): 2917-2924.
- Kumar R, Shukla SK, Pandey M, Pandey A, Pathak A and Dikshit A: Synthesis and antimicrobial effects of colloidal Gold nanoparticles against prevalent water borne bacterial pathogens. *Cogent Chemistry* 2016; 1192522: 1-9.
- NCCLS: Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically (5th Ed.), Approved standard, M7-A5. Pennsylvania: Wayne 2003.
- Brand-Williams W, Cuvelier ME and Berset C: Use of a free radical method method to evaluate antioxidant activity. *Lebensm.-Wiss. u.-Technol.* 1995; 28: 25-30.
- Pandey A, Mishra RK, Tiwari AK, Kumar A, Bajaj AK, Dikshit A: Management of cosmetic embarrassment caused by *Malassezia spp.* with fruticose lichen *Cladia* using phylogenetic approach. *Hindawi Biomed Research International* 2013; 8.
- Martins MCB, Lima MJG, Silva FP, Ximenes EA, Silva NH, and Pereira EC: *Cladia aggregata* (lichen) from Brazilian Northeast: Chemical characterization and antimicrobial activity. *Brazilian Archives of Biology and Technology* 2010; 53: 115-122.
- Randhir R, Lin YT and Shetty K: Stimulation of phenolic, antioxidant and antimicrobial activities in dark germinated mung bean sprouts in response to peptide and phyto-chemical elicitors. *Process Biochem.* 2004; 39: 637-646.
- Vattem DA, Lin YT, Lable RG and Shetty K: Phenolic antioxidant mobilization in cranberry pomace by solid-state bioprocessing using food fungus *Lentinusedodes* and effect on antimicrobial activity select food borne pathogens. *Innovative Food Sci Emerg Tech* 2004; 5: 81-91.
- Mueller K: Pharmaceutically relevant metabolites from lichens. *Applied Microbio. and Biotech.* 2001; 56: 9-16.
- Yilmaz M, Türk AO, Tay T and Kivanc M: The antimicrobial activity of extracts of the lichen *Cladonia foliacea* and its (-)-usnic acid, atranorin and fumaro-protetraric acid constituents. *Z Nat.* 2004; 59c: 249-254.
- Behera BC, Verma N, Sonone A and Makhija U: Antioxidant and antibacterial properties of some cultured lichens. *Bio-resource Technology* 2008; 99: 776-784.
- Verma N, Behera BC and Sharma BO: Glucosidase inhibitory and radical scavenging properties of lichen metabolites salazinic acid, sekikaic acid and usnic acid. *Hacettepe J. Biol. and Chem.* 2012; 40 (1): 7-21.
- Mishra RK, Mishra V, Pandey A, Tiwari AK, Pandey H, Sharma S, Pandey AC and Dikshit A: Exploration of anti-*Malassezia* potential of *Nyctanthes arbor-tristis* L. and their application to combat the infection caused by *Mala s1* a novel allergen. *BMC Complementary and Alternative Medicine* 2016; 16(114): 1-14.
- Mishra T, Shukla S, Meena S, Singh R, Pal M, Upreti DK, and Datta D: Isolation and identification of cytotoxic compounds from a fruticose lichen *Roccella montagnei*, and its *in-silico* docking study against CDK-10. *Revista Brasileira de Farmacognosia* 2017; 27(6) 724-728
- Singh BN, Upreti DK, Gupta VK, Dai XF and Jiang Y: Endolichenic Fungi: A Hidden Reservoir of Next Generation Biopharmaceuticals. *Trends in Biotechnology* 2017; 35(9): 808-813.
- Singh BN, Prateeksha, Upreti DK, Singh BR, Defoirdt T and Gupta VK: Bactericidal, quorum quenching and anti-biofilm nanofactories: a new niche for nanotechnologists. *Critical Reviews in Biotechnology* 2017; 37(4): 525-540.
- Prateeksha, Paliya BS, Bajpai R, Jadaun V, Kumar J, Kumar S, Upreti DK, Singh BR, Nayaka S, Joshi Y and Singh BN: The genus *Usnea*: a potent phytomedicine with multifarious ethnobotany, phytochemistry and pharmacology. *RSC Advances* 2016; 26: 25.

How to cite this article:

Pandey A, Qidwai A, Pandey M, Kumar R, Shukla SK and Dikshit A: Bactericidal screening and estimation of free radical scavenging activity of fruticose lichen *Cladonia scabriuscula* delise (nyl.) from Gopeshwar, Uttarakhand, India. *Int J Pharm Sci Res* 2018; 9(7): 2847-51. doi: 10.13040/IJPSR.0975-8232.9(7).2847-51.