



Received on 12 March, 2017; received in revised form, 13 October, 2017; accepted, 20 October, 2017; published 01 November, 2017

LETHAL EFFECTS OF *PISTACIA ATLANTICA* LEAVES EXTRACT AGAINST PROTO-SCOLECES OF HYDATID CYSTS

Mohammad Reza Aflatoonian^{1, 2}, Mojgan Saki³, Farnaz Kheirandish⁴, Hossein Mahmoudvand³ and Mohammad Reza Nazer^{*4}

Bam University of Medical Sciences¹, Bam, Iran.

Research Center for Tropical and Infectious Diseases², Kerman University of Medical Sciences, Kerman, Iran.

Student Research Committee³, Razi Herbal Medicines Research Center⁴, Lorestan University of Medical Sciences, Khorramabad, Iran.

Keywords:

Pistacia, Protoscoleces,
Cystic echinococcosis, *In-vitro*

Correspondence to Author:

Dr. Mohammad Reza Nazer

Razi Herbal Medicines Research
Center, Lorestan University of
Medical Sciences, Khorramabad,
Iran.


E-mail: Dr_nazer1@yahoo.com

ABSTRACT: Surgery is one of the best options to treat cystic echinococcosis (CE, hydatid cyst). To date, surgeon used different protoscolicidal drugs to eliminate of protoscoleces during CE surgery, but most of these agents have some serious side effects. The current study aims to explore the *in-vitro* protoscolicidal effects of *Pistacia atlantica* leave extract against hydatid cyst proto-scolices. Proto-scolices were collected from livers of infected sheep and were exposed with different concentrations of *P. atlantica* extract (8.75 - 70 mg/ml) for 10 - 60 min. Finally, to explore the viability of proto-scolices eosin exclusion examination was applied. The major components of leave extract of *P. atlantica* in gas chromatography/ mass spectrometry were trans-caryophyllene (15.18%), α -amorphene (8.1%), and neoalloocimene (6.21%), respectively. The obtained results revealed that *P. atlantica* leave extract had remarkable protoscolicidal effects against proto-scoleces of hydatid cysts; so that killed completely (100%) all of proto-scoleces at the dose of 70 mg/ml after 10 min of treatment. Also, *P. atlantica* leave extract at the concentration of 35 mg/ml was eliminated 100% of proto-scoleces after 20 min of incubation. These findings indicated that *P. atlantica* leave extract had remarkable protoscolicidal properties versus *hydatid cyst* proto-scoleces which indicated the ability of *P. atlantica* as a natural resource to produce of novel proto-scolicidal drug for CE surgery.

INTRODUCTION: Historically, the use of medicinal plants to treat illness and keep public health is highly prevalent around the world. Plant materials and their derivates show a central role in the field of novel drugs investigation for treatment or prevention of a wide range of diseases such as infectious ones¹.

One of these interesting plants is *Pistacia atlantica* Desf. (Anacardiaceae), which commonly observed in various parts of the world such as Iran. In folk medicine, some parts of the *P. atlantica* have been extensively applied to treat several disease and illness such as diarrhea, asthma, anemia and infectious diseases². Reviews have also reported anti-inflammatory, anticancer, antitumor, antialergic and antimicrobial properties of *P. atlantica*².

Cystic echinococcosis (CE) or hydatidosis consider as one of the main parasitic diseases in human and animals which caused by a small dog tapeworm, *Echinococcus granulosus*³.

ICK RESPONSE CODE	DOI: 10.13040/IJPSR.0975-8232.8(11).4653-58
	Article can be accessed online on: www.ijpsr.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.8(11).4653-58	

Normally, humans acquire infection *via* consumption of vegetables, food or water contaminated with *E. granulosus* eggs. CE in human because of its effects on vital organs including liver and lung, and livestock due to considerable economic problems in stock breeding has attracted World Health Organization (WHO) attention to itself⁴. From last decades, surgery is the most common choice in CE treatment. Surgeons during open cyst removal use some chemical scolicalidal agents to decrease the risk linkage of proto-scolecex⁵.

However, it has been proven that these common proto-scolicalidal agents have some side effects such as necrosis of liver^{6, 7}. Therefore, search for a proper protoscolicalidal drug with potent efficacy and lowest side effects is necessary for surgeons. The present investigation aims to evaluate the protoscolicalidal effects *P. atlantica* on protoscolecex of hydatid cyst on *in - vitro*.

MATERIALS AND METHODS:

Plant Materials: Leaves of *P. atlantica* were collected from mountains of Lorestan province, West of Iran, in September 2014. The materials were determined by a botanist of the Botany Department of Lorestan University, Lorestan, Iran. A voucher sample also has been recorded in the herbarium of Research Center for Agriculture Sciences, Khorramabad, Iran (LF 2522).

Preparation of Extract: One hundred g of powdered dried leaves were isolated using percolation method with methanol (80%) for 72 h. in 21 °C. The extract was crossing through filter paper (Whatman No. 3, Sigma, Germany) to delete artifacts. Finally, the extract was concentrated at 50 °C using a rotary evaporator (Heidolph, Germany) and kept at -20 °C, until use^{8,9}.

Gas Chromatography/Mass Spectrometry (GC/MS) Analysis of Extracts: Chemical compounds of *P. atlantica* extract were identified by extracting them using a solid phase micro extraction (SPME) technique. Initially, the samples were first ground into a fine powder using a household mill. Two grams of the sample was weighed and transferred to a 20 ml vial. Then the vial containing the sample was transferred to the ultrasonic device to extract the volatile substances. The temperature of the ultrasonic device was set on 50 °C for 15 min. In

the next step, the SPME fiber was placed on the upper space of the sample for 40 min to extract the volatile compounds. Immediately after the extraction, the SPME fiber was injected into the GC-Mass device for desorbing and identifying the compositions of sample. Desorbing was performed in the GC column for 2 min.

The SPME fiber holder for manual use and the 100 µm Polydimethyl siloxane (PDMS) fibers were obtained from Supelco (Bellefonte, PA, USA)¹⁰. GC/MS was carried out by an Agilent 6890N coupled with a HP-5MS column (30m × 0.25 mm, film thickness 0.25 mm). The column temperature was maintained at 40 °C for 12 min and programmed to 180 °C at a rate of 10 °C per min, and kept constant at 180 °C for 4 min. Injector and interface temperatures were 250 and 280 °C, respectively. The flow rate of Helium as carrier gas was (1 mL/min C.F). The percentages were calculated by electronic integration of FID peak areas without the use of response factors correction. Linear retention indices for all components were determined by co-injection of the samples with a solution containing homologous series of C8-C24 *n*-alkanes.

Identification of Components: The constituents of the extracts were determined in comparison with their standards Wiley 2001 library data of the GC/MS system or compared with those of reported in the previous studies¹¹.

Collection of Proto-scolecex: *E. granulosus* proto-scolecexes were obtained from the livers of naturally infected livestock slaughtered at Kerman abattoir, Iran. The cyst fluid was aseptically aspirated and was left to set for 30 min. After washing of proto-scolecexes for two times with PBS (pH 7.2) solution the number of protoscolecex/ml was adjusted as 2×10^3 protoscolecexes in 0.9 % NaCl solution with at least 90 % viability rate¹⁶.

Proto-scolicalidal Effects: To determine the lethal effects *P. atlantica* on hydatid cyst protoscolecexes, various doses of *P. atlantica* extract were applied for 10, 20, 30 and 60 min. The protocol was performed according to the method described by Mahmoudvand *et al.*, (2014). Initially, 0.5 ml of the protoscolecexes (2×10^3 /ml) and 0.5 ml of different doses of tested extract was added to each test tube. Test tubes were slowly mixed and then kept at

37°C for 10 - 60 min. After the time of incubation, the supernatant was discarded and 50 µl of 0.1 % eosin stain was added to protoscolecocytes and smeared on a glass slide and tested under a light microscope¹⁷. The percentages of dead proto-scolecocytes were counted by counting 100 proto-scolecocytes^{18, 19}. Moreover, normal saline and hypertonic saline of 20% were applied as negative and positive controls.

Evaluation of Viability of Protoscolecocytes: In this study, eosin exclusion assay (eosin solution 0.1%) was applied to determine the viability of protoscolecocytes of hydatid cysts¹⁴. The live proto-scolecocytes are colorless and displayed characteristic muscular

movements and flame cell activity after exposure to the stain, whereas dead proto-scolecocytes are red color.

Statistical Analysis: Statistical analysis was performed using SPSS software (17.0). Moreover, *t*-test was applied to determine the differences between groups. $P < 0.05$ was also considered statistically significant¹⁵.

RESULTS:

GC/MS Analysis of Extracts: Table 1 shows the identified compounds of leave extract of *P. atlantica* using GC/MS.

TABLE 1: LEAVE EXTRACT COMPOSITION OF *P. ATLANTICA* IDENTIFIED BY GC/MS

S. no.	Compound	Retention time	% of components
1	α-Pinene	6.67	3.345
2	β-Pinene	7.89	0.801
3	β-Myrcene	8.30	2.097
4	Tetradecane	10.36	0.480
5	γ-Terpinene	10.48	0.715
6	Nonanal	12.03	0.554
7	L-camphor	13.44	0.134
8	Menthone	13.76	1.298
9	Borneol	14.30	0.662
10	L-Menthol	14.54	1.725
11	Dodecane	15.19	0.369
12	Bornyl acetate	18.18	0.725
13	Thymol	19.04	0.180
14	Dodecane	19.41	0.633
15	Camphene	20.30	0.837
15	Aromadendrene	20.98	3.137
17	Copaene	21.12	1.818
18	α-Cadinene	21.55	0.417
19	Tetradecane	21.77	1.113
20	α-Gurjunene	22.15	1.826
21	trans-Caryophyllene	22.48	15.18
22	Calarene	22.70	1.290
23	Germacrene D	22.91	1.149
24	Nealloocimene	23.05	6.21
25	β-Selinene	23.15	1.493
26	α-Humulene	23.47	3.000
27	Heptacosane	23.56	1.100
28	Valencene	23.65	1.537
29	α-Amorphene	24.11	8.1
30	Dodecane	24.48	2.451
31	Isodene	24.61	2.989
32	α-Muurolene	24.76	2.803
33	β-Cadinene	25.16	3.669
34	delta.-Cadinene	25.38	5.915
35	β-Cadinene	25.69	0.607
36	α-Muurolene	25.77	1.143
37	α-calacorene	25.95	1.118
38	Nerolidol	26.47	1.422
39	Caryophyllene oxide	27.00	1.620
40	1,2-Dihydro-2,2,3-trimethyl-1-quinoxalinol-4-dioxide	28.35	3.436
41	Eudesmol	28.79	1.718
42	Tritetracontane	30.68	0.901
43	Tetracosane	32.75	1.699
	Total		85.7

The major components of leave extract of *P. atlantica* were trans-caryophyllene (15.18%), α -amorphenone (8.1%), and neoalloocimene (6.21%), respectively.

Protoscolicidal Effects: As shown in Table 2 *P. atlantica* leave extract demonstrated protoscolicidal activity against hydatid cysts protoscoleces. The current finding revealed that *P. atlantica* leave extract had remarkable protoscolicidal effects against protoscoleces of hydatid cysts; so that killed completely (100%) all of proto-scoleces at the dose of 70 mg/ml after 10 min of treatment. Also, *P. atlantica* leave extract at the concentration of 35 mg/ml was eliminated 100% of protoscoleces

after 20 min of incubation. The protoscolicidal effect of *P. atlantica* leave extract at the concentration of 17.5 mg/ml was 14.3, 37.6, 63.3 and 100 % after 10, 20, 30 and 60 min of incubation, respectively.

The mean dead of proto-scoleces in the normal saline control group was 5.3 % after 60 min of treatment, while the proto-scolicidal effects of 20% hypertonic saline as the positive controls was 100% after and 10 min of application. These findings exhibited that the scolicedal activity of *P. atlantica* leaves extract was significant higher ($p < 0.05$) in comparison with the control group at all exposure times.

TABLE 2: SCOLICIDAL EFFECTS OF *P. ATLANTICA* LEAVE EXTRACT AGAINST PROTO-SCOLECES OF HYDATID CYST AT VARIOUS CONCENTRATIONS FOLLOWING VARIOUS EXPOSURE TIMES

Concentration (mg/mL)	Exposure time (min)	Mean of mortality rate (%)
70	10	100
	20	100
	30	100
	60	100
35	10	59.6
	20	100
	30	100
	60	100
17.5	10	14.3
	20	37.6
	30	63.3
	60	100
8.75	10	6.6
	20	29.3
	30	52.3
	60	93.3
Normal saline	10	1.3
	20	2.6
	30	4.3
	60	7.1
2.0 % Hypertonic saline	10	100
	20	100
	30	100
	60	100

DISCUSSION: Medicinal plants and their pure compounds show proper opportunities for new drug searches in order to the having many benefits such as higher efficacy and availability as well as lower toxicity¹⁶. Here, we found that *P. atlantica* leave extract had remarkable protoscolicidal effects against proto-scoleces of hydatid cysts; so that killed completely (100%) all of protoscoleces at the dose of 70 mg/ml after 10 min of treatment. Also, *P. atlantica* leave extract at the concentration of 35 mg/ml was eliminated 100 % of protoscoleces after 20 min of incubation. This scolicedal effects of *P.*

atlantica is comparable with scolicedal activity of hypertonic saline, silver nitrate and mannitol, cetrimide, ethyl alcohol (95%), H₂O₂ and 10% providone iodine, albendazole, chlorhexidine gluconate, Selenium NPs and some plant extracts¹⁷⁻²⁴. Reviews have reported antibacterial, antiviral, antifungal and antiparasitic effects of *P. atlantica*³.

We found that the major components of leave extract of *P. atlantica* were trans-caryophyllene (15.18%), α -amorphenone (8.1%), and neoalloocimene (6.21%), respectively. However, according

to the previous investigations chemical composition of extracts depend on species, climate, and time of collection along with growth stage, thereby altering the biological activities studied²⁵.

Previous studies also revealed the presence of terpenoids, phenols, flavonoids, fatty acids and sterols in phytochemical screening of the crude extract *P. atlantica*³. In several investigations biological especially antimicrobial activities of these components have been proven²⁴⁻²⁷. Thus, we can suggest that these components are responsible for the scolicalidal activity of *P. atlantica*; however their exact action mechanism is not clearly understood. Regarding the some mechanism of antimicrobial terpenoids compounds such as monoterpenes, investigations have demonstrated that these compounds diffuse into pathogen and destroy cell membrane. Other studies also reported that anti-microbial effects of these components interfering with vital intracellular sites when cross the cell membranes and penetrate into the cell^{28, 29}.

CONCLUSION: These findings indicated that *P. atlantica* leave extract had remarkable protoscolicalidal properties versus hydatid cyst protoscolices which indicated the ability of *P. atlantica* as a natural resource to produce of novel protoscolicalidal drug for CE surgery.

ACKNOWLEDGEMENT: Nil.

CONFLICT OF INTEREST: The author declares that there is no conflict of interests in this study.

REFERENCES:

1. Mahmoudvand H, Saedi Dezaki E, Ezatpour B, Sharifi I, Kheirandish F, and Rashidipour M: *In-vitro* and *In-vivo* antileishmanial activities of *Pistacia vera* essential Oil. *Planta Medica* 2016; 82(4): 279-284.
2. Bozorgi M, Memariani Z, Mobli M, Salehi Surmaghi MH, Shams-Ardekani MR and Rahimi R: Five *Pistacia* species (*P. vera*, *P. atlantica*, *P. terebinthus*, *P. khinjuk*, and *P. lentiscus*): a review of their traditional uses, phytochemistry and pharmacology. *Sci World J.* 2013; 15: 219815.
3. World Health Organization (WHO): informal working group on echinococcosis. *Bull WHO* 1996; 74: 231-42
4. Eckert J and Deplazes P: Biological, epidemiological and clinical aspects of echinococcosis, a zoonosis of increasing concern. *Clin Microbiol Rev* 2004;17: 107-135
5. Brunetti E, Kern P and Vuitton DA: Writing Panel for the WHO-IWGE expert consensus for the diagnosis and treatment of cystic and alveolar echinococcosis in humans. *Acta. Trop.* 2010; 114(1): 1-16.

6. Sahin M, Eryilmaz R and Bulbuloglu E: The effect of scolicalidal agents on liver and biliary tree (experimental study). *J Invest Surg* 2004; 17(6): 323-6.
7. Rajabi MA: Fatal reactions and methaemoglobinaemia after silver nitrate irrigation of hydatid cyst. *Surg Pract* 2009; 13: 2-7.
8. Ezatpour B, Saedi Dezaki E, Mahmoudvand H, Azadpour M and Ezzatkah F: *In-vitro* and *in-vivo* antileishmanial effects of *Pistacia khinjuk* against *Leishmania tropica* and *Leishmania major*. *Evid Based Complement Alternat Med.* 2015; 149707.
9. Mahmoudvand H, Tavakoli Oliaei R, Mirbadie SR, Kheirandish F, Tavakoli Kareshk A, Ezatpour B, Mahmoudvand H. Efficacy and Safety of Bunium Persicum (Boiss) to Inactivate Protoscolices during Hydatid Cyst Operations. *Surg Infect (Larchmt)*. 2016; 17(6): 713-719.
10. Ghiasvand AR, Nasseri M, Farsizaeh S, Meshkatsaladat MH, Sadeghi-Sarabi R, Shadabi S and Borzoei M: Chemical characterization of cultivated *Tagetes minuta* L. by use of ultrasound-assisted head space SPME and GC-MS. *Chromatographia*. 2011; 73: 1031-1035.
11. Adams RP: Identification of essential oil components by gas chromatography/mass spectroscopy; Illinois, USA: Carol Stream, IL: Allured Publishing Corporation 2004.
12. Jahanbakhsh S, Azadpour M, Tavakoli Kareshk A, Keyhani A and Mahmoudvand H: *Zataria multiflora* Bioss: lethal effects of methanolic extract against protoscolices of *Echinococcus granulosus*. *J Parasit Dis DOI* 10.1007/s12639-015-0670-4.
13. Mahmoudvand H, Fallahi S, Mahmoudvand H, Shakibaie M, Harandi MF and Dezaki ES: Efficacy of *Myrtus communis* L. to inactivate the hydatid cyst protoscolices. *J Invest Surg.* 2015; 18: 1-7.
14. Smyth JD and Barrett NJ: Procedures for testing the viability of human hydatid cysts following surgical removal, especially after chemotherapy. *Trans R Soc Trop Med Hyg* 1980; 74: 649-652.
15. Mahmoudvand H, Mirbadie SR, Ghasemi Kia M, Badparva E, Shamsadini Lori S, Fasihi Harandi M. Efficacy of *Pistacia khinjuk* Fruits on Viability of Hydatid Cyst Protoscolices and Its Acute Toxicity in Mice Model. *Iran J Parasitol.* 2016; 11(3):383-388.
16. Rocha LG, Almeida JR, Macedo RO and Barbosa-Filho JM: A review of natural products with antileishmanial activity. *Phytomedicine* 2005; 12: 514-535.
17. Landa García JI, Alonso E, Gonzalez-Uriarte J and Rodriguez Romano D: Evaluation of scolicalidal agents in an experimental hydatid disease model. *Eur Surg Res* 1997; 29: 202-208.
18. Paksoy Y, Odev K, Sahin M, Arslan A and Koç O: Percutaneous treatment of hydatid cysts: comparison of direct injection of albendazole and hypertonic saline solution. *AJR Am J Roentgenol* 2005; 185: 727-734.
19. Puryan K, Karadayi K, Topcu O, Canbay E, Sumer Z, Turan M, Karayalcin K and Sen M: Chlorhexidine gluconate: an ideal scolicalidal agent in the treatment of intraperitoneal hydatidosis? *World J Surg* 2005; 29: 227-230.
20. Kilicoglu B, Kismet K, Koru O, Tanyuksel M, Oruc MT, Sorkun K and Akkus MA: The scolicalidal effects of honey. *Adv Ther* 2006; 23: 1077-1083.
21. Mahmoudvand H, Fasihi Harandi M, Shakibaie M, Aflatoonian MR, Makki MS and Jahanbakhsh S: Scolicalidal effects of biogenic selenium nanoparticles against protoscolices of hydatid cysts. *Int J Surg.* 2014; 12: 399-403.

22. Mahmoudvand H, Asadi A, Harandi MF, Sharififar F, Jahanbakhsh S and Dezaki ES: *In vitro* lethal effects of various extracts of *Nigella sativa* seed on hydatid cyst protoscoleces. Iran J Basic Med Sci. 2014; 17(12):1001-6.
23. Mahmoudvand H, Saedi Dezaki E, Kheirandish F, Ezatpour B, Jahanbakhsh S and Fasihi Harandi M: Scolicidal effects of black cumin seed (*Nigella sativa*) essential oil on hydatid cysts. Korean J Parasitol. 2014; 52(6): 653-659.
24. MahmoudvandH, Sharififar F, Saedi Dezaki E, EzatpourB, Jahanbakhsh S and Fasihi Harandi M: Protoscolecidal effect of *Berberis vulgaris* root extract and its main compound, berberine in cystic echinococcosis. Iranian J Parasitol. 2014; 9(4): 26-34.
25. Saedi Dezaki E, Mahmoudvand H, Sharififar F, Fallahi S, Monzote L and Ezatkah F: Chemical composition along with anti-leishmanial and cytotoxic activity of *Zataria multiflora*. Pharm Biol. 2015; 8: 1-7.
26. Sokovic M and van Griensven LG: Antimicrobial activity of essential oils and their components against the three major pathogens of the cultivated button mushroom, *Agaricus bisporus*. Eur J Plant Pathol 2006; 116: 211-224.
27. Abbaszadeh S, Sharifzadeh A, Shokri H, et al., Antifungal efficacy of thymol, carvacrol, eugenol and menthol as alternative agents to control the growth of food-relevant fungi. J Myco Med 2014; 24 (2): 51-56.
28. Sikkema J, De Bont DA and Poolman B: Mechanisms of membranetoxicity of hydrocarbons. Microbiol Mol Biol Rev 1995; 59: 201-222.
29. Cristani M, D'Arrigo M, Mandalari G, Castelli F, Sarpietro MG, Micieli D, Venuti V, Bisignano G, Saija A and Trombetta D: Interaction of four monoterpenes contained in essential oils with model membranes: implications for their antibacterial activity. J Agric Food Chem 2007; 55(15): 6300-6308.

How to cite this article:

Aflatoonian MR, Saki M, Kheirandish F, Mahmoudvand H and Nazer MR: Lethal effects of *Pistacia atlantica* leaves extract against protoscoleces of hydatid cysts. Int J Pharm Sci Res 2017; 8(11): 4653-58.doi: 10.13040/IJPSR.0975-8232.8(11).4653-58.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)