



Received on 16 May, 2017; received in revised form, 21 July, 2017; accepted, 25 July, 2017; published 01 February, 2018

THERAPEUTIC POTENTIAL OF CUCURBITS AGAINST RADIATION INDUCED INFLAMMATION AND DAMAGE IN MICE GUT

Sonal Chauhan, Dhara Sharma and Harish Chandra Goel*

Amity Centre for Radiation Biology, Amity University Uttar Pradesh (AUUP), Sector - 125, Noida - 202303, Uttar Pradesh, India.

Keywords:

Radiation damage, Gut-inflammation, Cytokines, Gut-microbiota, Villi, Dietary-cucurbits

Correspondence to Author:

Harish Chandra Goel

Amity Centre for Radiation Biology, Amity University Uttar Pradesh (AUUP), Sector - 125, Noida - 202303, Uttar Pradesh, India.

E-mail: hcgoel@amity.edu

ABSTRACT: Whole body or local irradiation to gut is known to induce significant damage to gut epithelium and crypt stem cells. A large population of microbes in the gut remains unaffected. We hypothesize that the modifications in diet and enteric microbial population may provide a non-toxic mechanism to counter radiation induced health deficit, especially in the gut and therefore investigated the effect of dietary cucurbits in this context. Radio-protective effects of *Lagenaria* (Ls), *Cucurbita* (Cp) and *Luffa* (Lc) were evaluated in mice for cytokines (in jejunum tissues) through ELISA and intestinal villi microscopically. Irradiation of mice increased pro-inflammatory biomarkers NAP-3, IFN- γ and IL-17 and decreased anti-inflammatory biomarkers IL-10, IL-4 and TGF- β . Post-irradiation treatment with cucurbit extract (Ls, Cp or Lc) decreased pro-inflammatory cytokines NAP-3, IFN- γ and IL-17 and increased anti-inflammatory cytokines IL-10, IL-4 and TGF- β in mice. Treatment with cucurbits restored structural integrity of villi in irradiated mice. Various antioxidants and anti-inflammatory molecules present in cucurbits, their positive effects on probiotics and host tissues and microbial fermentation of dietary fibers may mitigate radiation induced damage. The cucurbits, as non-toxic anti-inflammatory agents and nutrition enhancer can be helpful in irradiated and inflamed systems.

INTRODUCTION: Whole body exposure to ionizing radiations results in production of reactive oxygen species, which initiate a series of signalling cascades and lead to the genesis of several pro-inflammatory responses in various tissues and systems leading HP and GI syndrome, depending on the dose received^{1, 2}. Radiation induced inflammation in the gut is one of the major events during radiotherapy and radiation accidents.

The gut mucosa, under normal conditions maintains a quiescent state against a diverse array of dietary antigens and luminal microbes. However exposure to ionizing radiation leads to gut inflammation and stem cell necrosis^{3, 4}. The inflammatory manifestations of the gut are therapeutically managed in palliative and symptomatic manner through analgesic, anti-emetic and anti-diarrheal agents, though most of them induce adverse effects⁵.

In view of this, the search for newer, less toxic and more effective anti-inflammatory agents becomes inevitable. In fact, a large number of agents of natural and synthetic origin have been attempted against gastro-intestinal (GI) inflammation⁶. The

<p>QUICK RESPONSE CODE</p>	<p>DOI: 10.13040/IJPSR.0975-8232.9(2).550-62</p> <hr/> <p>Article can be accessed online on: www.ijpsr.com</p> <hr/> <p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.9(2).550-62</p>
-----------------------------------	---

human colonic microbiota is very large, about 10 - 100 times the total number of human cells in an individual, comprises a consortium of a large number of bacterial species (450 - 500), and out of this only about 45 - 50 spp. are present in an individual⁷. The triangular interaction between host, the enteric microbiota and the diet, leads to the generation of metabolome which remains central to all the organs and systems of the body for nutrition, growth and the repair / recovery. The modification of metabolomic contents can be manipulated through diet and exploited for alleviation of the radiation induced health deficits. The recent literature extensively reveals about the influence of diet on host and microbiota and their subsequent implications in inflammatory and immune responses^{8,9}.

The inflammatory response is coordinated by a wide range of signalling molecules and pathways that constitute complex regulatory networks like NF κ B and MAPK signalling pathways. Nuclear Factor-kappaB (NF- κ B) is the gene transcription factor, ubiquitously present in every cell and has a key role in enhancing immune and inflammatory responses¹⁰. NF κ B upregulate expression of genes for pro-inflammatory cytokines, chemokines, enzymes, receptors, apoptotic regulators, growth factors and adhesion molecules that play a key role in the inflammatory processes and regulate several aspects of innate and acquired immune responses¹⁰. A large number of the immune-competent cells are located in the intestinal epithelium, Peyer's patches, lamina propria and isolated lymphoid follicles. Within the lymphocyte subsets that populate mucosa and gut-associated lymphoid tissues (GALT), T helper cell populations (Th-1, Th-2, and Th-17) and a wide variety of regulatory T cells (T_{regs}) such as Th₃ (primarily secrete TGF- β) and Tr₁ (secrete high levels of IL-10) are noteworthy¹¹. Th-17 cells have been identified as major mediators of pathology in the inflammatory disease models. The balance between Th-17 and T_{reg} populations is important for maintaining a healthy immune status.

A number of medicinal plants evaluated for their radio-protective efficacy have shown protective effects against the damaging effects of ionizing radiation. An urge to consume cucurbit diet during dysbiosis may be an indication of signalling by gut-

brain axis and this led to the selection of cucurbit plants for this study. Cucurbits were investigated for their effect on various inflammatory biomolecules involved in intestinal inflammation. Since ancient times, the cucurbits have been used as prebiotics for treating pathological conditions of GI tract^{12,13}. A wide range of chemical compounds including sterols, terpenoids (cucurbitacins), flavonoids, bryonolic acid and saponins have been isolated from various cucurbit plants and have been reported to exhibit analgesic, anti-inflammatory, immunomodulatory and antioxidant^{12,14,15}.

To counter or mitigate the inflammatory action in GI tract several prebiotics have been demonstrated to suppress pro-inflammatory mediators like tumor necrosis factor- α (TNF- α), interferon- γ (IFN- γ) and IL-8^{9,16}, which are found in high levels in intestinal mucosa of patients suffering of inflammatory bowel syndrome (IBS). Prebiotics also modulate immune responses by down-regulating T helper population but little information is available on the direct effect of prebiotics on various populations of T-helper cells like Th₁, Th₂, Th₁₇ and T_{regs} and warranted investigation. The damage to GI system is also reflected in the shortening of intestinal villi through destruction of the stem cells (crypt cells). The intestinal villus is a self-renewing organ and the mucosa overlying it is replaced entirely every 3 - 5 days. Stem cells present at the base of each villus are responsible for self-renewal of intestinal villi. During radiotherapy of GI tumors, the highly radiosensitive stem cells of the crypts get damaged and subsequently, inhibition of stem cell proliferation leading to shortening of villi height.

Cucurbits (*Lagenaria siceraria*, *Luffa cylindrica* and *Cucurbita pepo*) have already been shown to display potent anti-inflammatory and radio-protective activities *in-vitro*¹⁷. *Lagenaria* extract has been reported to improve the tight junction integrity and length of the villi in the mice which were subjected to radiation exposure (10 Gy) *in-vivo*¹⁸.

In view of the sparse information regarding the effect of cucurbits as prebiotics on modulation of inflammatory response, we investigated the effect of cucurbit extracts (Ls, Cp, and Lc) on various inflammatory (NAP-3, IFN- γ and IL-17) and anti-

inflammatory (IL-10, TGF- β and IL-10) cytokines and intestinal villi morphology (height and width).

MATERIALS AND METHODS:

Reagents: Various chemicals and reagents were procured from different sources: Formaldehyde, Ethanol, Xylene, Paraffin, Canada balsam from Merck (Mumbai); Hematoxylin and Eosin from M/S Fisher Scientific, Mumbai. Protease inhibitor cocktail from M/s Sigma Aldrich (USA). Bradford reagent from Bio-Rad Laboratories (USA). ELISA kits for NAP-3, IL-17, IFN- γ , IL-10, IL-4, and TGF- β from Cloud-Clone Corp., Houston.

Preparation of Plant Extracts: Fresh fruits of cucurbits, *Lagenaria siceraria* (Ls), *Luffa cylindrica* (Lc), *Cucurbita pepo* (Cp) were procured from local market, identified by the botanist at Department of Biosciences, Jamia Millia Islamia, New Delhi and were thoroughly washed with distilled water. Known quantity of a plant material was homogenized in distilled water and kept for 24 h at 25 °C and was filtered thereafter through a fine strainer having a spread of muslin cloth. The filtrate was further passed through membrane filter of 0.22 μ , lyophilized and stored at 4 °C in air tight bottles.

Animals: Balb/c female mice (14 weeks old) weighing about 25g were bred, housed and treated under specific pathogen free laboratory environment (25 \pm 2 °C, photoperiod - 12 h) and were given standard animal diet (Lipton, India) and tap water *ad libitum*. The experimental protocol was approved by institutional animal ethics committee of Amity University, Noida.

Irradiation: The Tele-therapy cobalt machine (Bhabhataron II) at 'Institute of Nuclear Medicine and Allied Sciences' Delhi, was used for whole body gamma irradiation (6 Gy) of experimental animals. The dose rate during these experiments ranged between 1.278 to 1.250 Gy / Minute and each mouse was irradiated individually.

Experimental Design: 66 mice were randomly assorted to 11 groups, each containing 6 animals. Cucurbit extract at the rate of 2 mg/kg body weight was administered to each mouse orally through a catheter. For sample collection, the animals were sacrificed by cervical dislocation on 8th day after irradiation. The animals were grouped as under:

- **Group 1 (n = 6):** Control, no treatment.
- **Group 2 (n = 6):** Irradiated control, received 6Gy (single dose) whole body gamma irradiation.
- **Group 3, 4, 5 (n = 6 in each group):** Post-irradiation cucurbit treatment (Protocol A), each mouse received 6 Gy WBI and one hour later received a single dose of cucurbit extract (Ls, Cp or Lc) for 7 post irradiation days.
- **Group 6, 7, 8 (n = 6 in each group):** Pre-irradiation cucurbit treatment (Protocol B), each mouse received a single dose of a cucurbit extract (Ls, Cp or Lc) for 7 days and irradiated on 8th day.
- **Group 9, 10, 11 (n = 6 in each group):** Pre + Post-irradiation cucurbit treatment (Protocol C), each mouse received cucurbit extract (Ls, Cp or Lc) for 7 days before and 7 days after irradiation.

Cytokine Analysis: The jejunal section (2cm long) about 3cm away from duodenal end was carefully taken out from each mouse, and was washed with cold PBS, pH 7.4, and fresh frozen immediately in liquid nitrogen for analysis of different cytokines (NAP-3, IFN- γ , IL-10, IL-4, IL-17, TGF- β). Samples were prepared following the method described by Liu *et al.*,¹⁹ Briefly, frozen tissues were homogenized in lysis buffer containing protease inhibitors. The homogenates were then centrifuged at 12,000g for 10 min at 4 °C after incubation on ice for 30 min. The supernatant from each sample was collected and stored at - 20 °C. In each sample total protein concentration was estimated using Bradford Reagent. Cytokines level in each sample was estimated through ELISA following manufacturer's instructions.

Histological Study: About 2cm long pieces of jejunum were taken out from each mouse and were fixed in 10% neutral formalin (pH 7.0–7.6) for 24 hours. The tissue was processed for dehydration and paraffin block making following standard procedure. Microtomy was done to get 5 μ thick sections which were stained using Haematoxylin and Eosin (H and E) following standard method²⁰. The sections were studied under the light microscope (40X) for various villar parameters

(height and width). Data for 10 villi per section was collected.

Statistical Analysis: The data from animals of each group was pooled and has been presented as mean \pm SE. Student's t test was applied to estimate the statistical significance difference between different experimental groups at $P < 0.05$.

RESULTS AND DISCUSSION: Exposure to ionizing radiation is known to generate oxidative stress, a pro-inflammatory response and modulation of the expression of pro- and anti-inflammatory cytokines. The inflammatory manifestations in the gut have been reported to be mediated *via* pro-inflammatory cytokines like TNF- α , IFN- γ , IL-17, IL-6 and NAP-3^{21, 22}. In this context, the immunosuppressive responses are mediated by anti-inflammatory cytokines such as IL-10^{23, 24}, TGF- β ²⁵ and IL-4²⁶.

Effects of Cucurbits on Expression of NAP-3 *in vivo*: Daily administration of cucurbit extract (Ls, Cp or Lc) orally to mice for 7 days post irradiation (Protocol-A) significantly decreased the expression of NAP-3 to 91, 89 and 83% of irradiated control, 100% (**Fig. 1**). Pre-treatment of mice with Ls for 7 days before irradiation (Protocol-B) also decreased NAP-3 expression significantly, 85% of irradiated control; while Cp, or Lc remained insignificant at decreasing the NAP-3, 92 and 91% of irradiated control respectively. Mice under protocol C received cucurbit extracts Ls, Lc or Cp for 7 days before and 7 days after irradiation and NAP-3 expression decreased to 82, 81 and 87% of irradiated control (100%) respectively. Cp when administered under protocol C (Pre- and post-irradiation combined treatment) rendered maximum decrease in NAP-3 when compared to control.




FIG. 1: RADIO-MODIFYING EFFECT OF CUCURBITS, *LAGENARIA SICERARIA* (LS), *CUCURBITA PEPO* (CP) AND *LUFFA CYLINDRICA* (LC) ON THE EXPRESSION OF NAP-3 IN THE HOMOGENATE OF JEJUNUM OF BALB/C MICE

Each animal in protocol A (Post -R) received a single whole body dose of 6 Gy and cucurbit extract was given each day for 7 days; Protocol B (Pre-R) received cucurbit extract each day for 7 days and 6 Gy dose on 8th day; Protocol C (Pre and Post -R) received a cucurbit extract each day for 7 days, and 6 Gy dose on 8th day and cucurbit extract each day for 7 days. The changes in NAP-3 were evaluated with respect to the irradiated control, considered as 100%. Each bar represented the Mean \pm SE of three experiments; asterisk above a bar indicated significance of difference with respect irradiated control (* $p < 0.05$, ** $p < 0.01$).

Effects of Cucurbits on Expression of IFN- γ *in vivo*: Daily administration of cucurbit extract Ls, Cp or Lc to mice orally for 7 days (post irradiation, Protocol-A) significantly decreased the expression of IFN- γ to 63, 64 and 60% respectively when compared with irradiated control, considered as 100% (**Fig. 2**). Pre-treatment of mice with cucurbit extracts Ls, Cp, or Lc for 7 days before irradiation (Protocol-B) also significantly decreased IFN- γ

expression to 72, 68 and 64% respectively as compared to irradiated control. Mice under protocol C received cucurbit extracts Ls, Lc or Cp for 7 days before and 7 days after irradiation and IFN- γ levels were 67, 64 and 64% of the control value. Post irradiation treatment of mice with cucurbit extract (Lc) rendered maximum decrease in IFN- γ levels as compared to control.




FIG. 2: RADIO-MODIFYING EFFECT OF CUCURBITS, *LAGENARIA SICERARIA* (LS), *CUCURBITA PEPO* (CP) AND *LUFFA CYLINDRICA* (LC) ON THE EXPRESSION OF IFN- γ IN THE HOMOGENATE OF JEJUNUM OF BALB/C MICE

Each animal in protocol A (Post -R) received a single whole body dose of 6 Gy and cucurbit extract was given each day for 7 days; Protocol B (Pre-R) received cucurbit extract each day for 7 days and 6 Gy dose on 8th day; Protocol C (Pre and Post -R) received a cucurbit extract each day for 7 days, and 6 Gy dose on 8th day and cucurbit extract each day for 7 days. The changes in IFN- γ were evaluated with respect to the irradiated control, considered as 100%. Each bar represented the Mean \pm SE of three experiments; asterisk above a bar indicated significance of difference with respect irradiated control (* p <0.05, ** p <0.01).

Effects of Cucurbits on Expression of IL-17 *in-vivo*: Daily administration of cucurbit extract Ls, Cp or Lc to mice orally for 7 days (post irradiation, Protocol-A) significantly decreased the expression of IL-17 to 54, 53 and 54% respectively when compared to irradiated control, 100% (Fig. 3). Pre-treatment of mice with cucurbit extracts Ls, Cp, or Lc for 7 days before irradiation (Protocol-B) also significantly decreased IL-17 expression to 73, 57

and 52% respectively as compared to irradiated control. Mice under protocol C received cucurbit extracts Ls, Lc or Cp for 7 days before and 7 days after irradiation and IL-17 levels were 65, 67 and 55% respectively with respect to irradiated control. Among the three protocols, post irradiation treatment (Protocol-A) of mice with cucurbit extract (Ls, Cp or Lc) rendered maximum decrease in IL-17 as compared to control.




FIG. 3: RADIO-MODIFYING EFFECT OF CUCURBITS, *LAGENARIA SICERARIA* (LS), *CUCURBITA PEPO* (CP) AND *LUFFA CYLINDRICA* (LC) ON THE EXPRESSION OF IL-17 IN THE HOMOGENATE OF JEJUNUM OF BALB/C MICE

Each animal in protocol A (Post -R) received a single whole body dose of 6 Gy and cucurbit extract was given each day for 7 days; Protocol B (Pre-R) received cucurbit extract each day for 7 days and 6 Gy dose on 8th day; Protocol C (Pre and Post -R) received a cucurbit extract each day for 7 days, and 6 Gy dose on 8th day and cucurbit extract each day for 7 days. The changes in IL-17 were evaluated with respect to the irradiated control, considered as 100%. Each bar represented the Mean \pm SE of three experiments; asterisk above a bar indicated significance of difference with respect irradiated control (* p <0.05, ** p <0.01, *** p <0.001).

Effects of Cucurbits on Expression of IL-10 *in-vivo*:

Daily administration of a cucurbit extract (Ls, Cp or Lc) to mice orally for 7 days post irradiation (protocol A), significantly increased the expression of IL-10 to 124, 114 and 132% respectively vs irradiated control considered as 100% (Fig. 4). Treatment of mice with cucurbit extract (Ls) for 7 days before irradiation (Protocol B) significantly increased IL-10 expression to 119%; however treatment of mice with Cp or Lc for 7 days before

irradiation insignificantly modified IL-10 levels to 106 and 103% respectively vs irradiated control. Mice under protocol C received cucurbit extracts Ls, Lc or Cp for 7 days before and 7 days after irradiation and the IL-10 expression increased to 114, 109 and 127% respectively with respect to irradiated control. Lc when administered under protocol A (post-irradiation treatment) rendered maximum increase in IL-10 when compared to control.




FIG. 4: RADIO-MODIFYING EFFECT OF CUCURBITS, *LAGENARIA SICERARIA* (LS), *CUCURBITA PEPO* (CP) AND *LUFFA CYLINDRICA* (LC) ON THE EXPRESSION OF IL-10 IN THE HOMOGENATE OF JEJUNUM OF BALB/C MICE

Each animal in protocol A (Post -R) received a single whole body dose of 6 Gy and cucurbit extract was given each day for 7 days; Protocol B (Pre-R) received cucurbit extract each day for 7 days and 6 Gy dose on 8th day; Protocol C (Pre and Post -R) received a cucurbit extract each day for 7 days, and 6 Gy dose on 8th day and cucurbit extract each day for 7 days. The changes in IL-10 were evaluated with respect to the irradiated control, considered as 100 %. Each bar represented the Mean ± SE of three experiments; asterisk above a bar indicated significance of difference with respect irradiated control (*p<0.05, **p<0.01).




FIG. 5: RADIO-MODIFYING EFFECT OF CUCURBITS, *LAGENARIA SICERARIA* (LS), *CUCURBITA PEPO* (CP) AND *LUFFA CYLINDRICA* (LC) ON THE EXPRESSION OF IL-4 IN THE HOMOGENATE OF JEJUNUM OF BALB/C MICE

Each animal in protocol A (Post -R) received a single whole body dose of 6 Gy and cucurbit extract was given each day for 7 days; Protocol B (Pre-R) received cucurbit extract each day for 7 days and 6 Gy dose on 8th day; Protocol C (Pre and Post -R) received a cucurbit extract each day for 7 days, and 6 Gy dose on 8th day and cucurbit extract each day for 7 days. The changes in IL-4 were evaluated with respect to the irradiated control, considered as 100%. Each bar represented the Mean ± SE of three experiments; asterisk above a bar indicated significance of difference with respect irradiated control (*p<0.05, **p<0.01).

Effects of Cucurbits on Expression of IL-4 *in-vivo*: Daily administration of a cucurbit extract (Ls, Cp or Lc) to mice orally for 7 days post irradiation (protocol A), significantly increased the expression of IL-4 to 144, 153 and 174% respectively vs irradiated control considered as 100% (Fig. 5). Treatment of mice with cucurbit extracts Ls, Cp or Lc for 7 days before irradiation (Protocol B) significantly increased IL-4 expression to 138, 121 and 146% respectively vs irradiated control. Mice under protocol C received cucurbit extracts Ls, Lc or Cp for 7 days before and 7 days after irradiation and IL-4 expression increased to 127, 141 and 139 % respectively with respect to irradiated control. Lc when administered under protocol A (post-irradiation treatment) rendered maximum increase in IL-4 when compared to control.

Effects of Cucurbits on Expression of TGF- β *in-vivo*: Daily administration of Ls orally to mice for 7 days post irradiation (protocol A) significantly

increased the expression of TGF- β to 121% with respect to irradiated control (100%); while the increase in TGF- β expression remained in significant (113%) with Cp or Lc (Fig. 6). Similarly treatment of mice with Ls for 7 days before irradiation (Protocol B) significantly increased TGF- β expression to 126% in comparison to irradiated control; while the increase in TGF- β expression remained insignificant (111% and 112%) with Cp or Lc respectively. Pre-treatment of mice with Ls or Cp for 7 days followed by irradiation after 24 h and again treatment with Ls or Cp for 7 days (Protocol C) rendered significant increase in TGF- β expression (125 and 117% respectively) with respect to irradiated control while Lc remained insignificant at increasing TGF- β (113%). Lc when administered under protocol A (post-irradiation treatment) rendered maximum increase in IL-10 when compared to control.




FIG. 6: RADIO-MODIFYING EFFECT OF CUCURBITS, *LAGENARIA SICERARIA* (LS), *CUCURBITA PEPO* (CP) AND *LUFFA CYLINDRICA* (LC) ON THE EXPRESSION OF TGF- β IN THE HOMOGENATE OF JEJUNUM OF BALB/C MICE.

Each animal in protocol A (Post -R) received a single whole body dose of 6 Gy and cucurbit extract was given each day for 7 days; Protocol B (Pre-R) received cucurbit extract each day for 7 days and 6 Gy dose on 8th day; Protocol C (Pre and Post -R) received a cucurbit extract each day for 7 days, and 6 Gy dose on 8th day and cucurbit extract each day for 7 days. The changes in TGF- β were evaluated with respect to the irradiated control, considered as 100%. Each bar represented the Mean \pm SE of three experiments; asterisk above a bar indicated significance of difference with respect irradiated control (* p <0.05, ** p <0.01).

In the present study, irradiation of mice rendered an increase of pro-inflammatory biomarkers NAP-3, IFN- γ and IL-17 (Fig. 1, 2 and 3 respectively) and concurrent decrease of anti-inflammatory biomarkers IL-10, IL-4 and TGF- β (Fig. 4, 5 and 6 respectively). This is in agreement with other studies reporting the increase of radiation induced pro-inflammatory cytokines and growth factors^{27, 28} and decrease of anti-inflammatory biomarkers²⁹.

Oral administration of cucurbit extract (Ls, Cp or Lc) to mice under all the three experimental protocols countered the inflammatory effects of radiation, by decreasing pro-inflammatory cytokines NAP-3 (Fig. 1), IFN- γ (Fig. 2) and IL-17 (Fig. 3) and increasing anti-inflammatory cytokines IL-10 (Fig. 4), IL-4 (Fig. 5) and TGF- β (Fig. 6) concurrently as compared to irradiated control (no cucurbit treatment).

Post-irradiation administration of cucurbit extracts was most effective in decreasing the IL-17 or IFN- γ and increasing the IL-10 or IL-4. However, NAP-3 was maximally decreased by Cp and TGF- β was maximally increased by Ls when administered under combined pre + post-irradiation cucurbit treatment protocol. Anti-inflammatory effects of cucurbits have also been demonstrated by other workers; Ls decreased cyclooxygenase and lipoxygenase³⁰, Cp down regulated IFN- γ ³¹ and Lc inhibited IL-6 and ROS generation³². Similarly, anti-inflammatory effects of other prebiotics via induction of IL-10 and down regulation of Th-1 and Th-17 cytokines have also been reported³³.

Anti-inflammatory effect of Ls, Cp and Lc may be attributed to various antioxidants present in cucurbits like, polyphenols, carotenoids, and Vitamins C and E³⁴⁻³⁶ which act as singlet oxygen quenchers and hydrogen ion donors to the free radicals and provide protection against oxidative damages. TGF- β , IL-10 and IL-4 have been known to inhibit ROS/RNS-mediated damage²⁷. Our previous studies on *in-vitro* system^{37, 38} have already demonstrated the antioxidant activities of cucurbit extracts. Flavonoid (epigallocatechin-3-gallate) from other dietary sources have also been reported to decrease inflammatory cytokines like IL-6, IL-17, IFN- γ and TNF- α through decrease in Th-1 and Th-17 population and increase in Tregs population enhancing IL-10 production³⁹. Immunomodulatory effect of cucurbits may also be attributed to various alkaloids, sterols and saponins^{40, 41}. Cucurbitacin, a flagship alkaloid of family cucurbitaceae also rendered anti-oxidant, anti-inflammatory and immunomodulatory effect both *in-vitro* and *in-vivo*¹⁵.

Anti-inflammatory and immuno-modulatory activity of Ls, Cp and Lc may also be attributed to short chain fatty acids (SCFAs) which are produced by metabolism of dietary fiber⁴² and are considered as enactors for prebiotics effects⁴³. SCFAs have been reported to modulate immune responses by inducing both T-cell and B-cell hyporesponsiveness and down-regulating Th-1, Th-2 and Th-17 cytokines^{42, 44}. Administration of B-GOS (5.5g daily for 10 weeks) has been reported to lower circulating levels of inflammatory cytokines such as IL-6, IL-1 β , and TNF- α and to enhance the level of IL-10^{45, 46}. Dietary fiber has been shown to enhance TGF-

beta signalling in the rat intestine through SCFAs⁴⁷. Butyrate (a SCFA) enemas have also been reported to reduce inflammation and oxidative stress in colonic mucosa of patients with ulcerative colitis by increasing colonic IL-10/IL-12 ratio⁴⁸.

Anti-inflammatory and immuno-modulatory effect of Ls, Cp and Lc may also be credited to their positive effects on probiotics growth. Our earlier studies have shown that cucurbits enhanced the activity of Lrh, a probiotics³⁷. Intragastric administration of IL-10 producing *Lactococcus lactis* protected mice from DSS-induced colitis and prevented development of colitis in IL-10-deficient mice⁴⁹. Probiotics have been reported to increase the production of an anti-inflammatory cytokine, IL-10⁵⁰. IL-17 production by lymphocytes has been reported to be decreased by probiotics in the gut^{51, 52}. Our study can be exploited for developing strategies to mitigate radiation induced damage at GI syndrome level.

Effects of Cucurbits on Height of Villi in Jejunum: The changes in the height of villi in jejunum of different mice have been displayed in **Fig. 7** and the microscopic images displaying normal, disrupted and recovering villi have been displayed in **Fig. 9**. Daily administration of a cucurbit extract (Ls, Cp or Lc) orally to mice for 7 days post irradiation (protocol A), significantly increased the villi height from $214 \pm 24\mu\text{m}$ (irradiated control) to 292 ± 32 , 264 ± 29 and $307 \pm 34\mu\text{m}$ respectively. Treatment of mice with cucurbit extract (Ls or Lc) for 7 days before irradiation (Protocol B) significantly increased villi length to 271 ± 30 and $281 \pm 31\mu\text{m}$ respectively as compared to irradiated control ($214 \pm 24\mu\text{m}$) while increase with Cp remained insignificant at $p < 0.05$ ($226 \pm 25\mu\text{m}$). Pretreatment of mice with Ls, Lc or Cp for 7 days followed by irradiation after 24 h and again treatment with Ls, Cp or Lc for 7 days (Protocol C) did not display significant recovery in villi length which remained at 220 ± 24 , 224 ± 25 and $232 \pm 26\mu\text{m}$.

Ls was most effective at increasing the villi height when administered under protocol A, where the increase was 21 and 72 μm more than protocol B and C respectively. Similarly, Cp rendered maximum increase in villi height under protocol A where the increase was 38 and 40 μm more than

protocols B and C respectively. Lc also displayed maximum increase in villi height under protocol A

where the increase was 26 and 75µm more than protocols B and C respectively.




FIG. 7: EFFECT OF CUCURBITS *LAGENARIA SICERARIA* (LS), *LUFFA CYLINDRICA* (LC) AND *CUCURBITA PEPO* (CP) ON THE CHANGES IN HEIGHT OF VILLI IN JEJUNUM OF BALB/C MICE

Each animal in protocol A (Post -R) received a single whole body dose of 6 Gy and cucurbit extract was given each day for 7 days; Protocol B (Pre-R) received cucurbit extract each day for 7 days and 6 Gy dose on 8th day; Protocol C (Pre and Post -R) received a cucurbit extract each day for 7 days, and 6 Gy dose on 8th day and cucurbit extract each day for 7 days. Each bar represented the Mean ± SE of three experiments. Asterisk above a bar represented significance of difference with irradiated control at *p<0.05, **<0.01 and ***<0.001 respectively.

Effects of Cucurbits on Width of Villi in Jejunum: The changes in the width of villi in jejunum of different mice have been displayed in **Fig. 8** and the microscopic images displaying normal, disrupted and recovering villi have been displayed in **Fig. 9**. Daily administration of a

cucurbit extract (Ls, Cp or Lc) orally to mice for 7 days post irradiation (protocol A), significantly increased the villi width from 39 ± 1.1µm (irradiated control) to 45 ± 2.1µm, 43 ± 2.4µm and 46 ± 1.3µm respectively.




FIG. 8: EFFECT OF CUCURBITS *LAGENARIA SICERARIA* (LS), *LUFFA CYLINDRICA* (LC) AND *CUCURBITA PEPO* (CP) ON THE CHANGES IN WIDTH OF VILLI IN JEJUNUM OF BALB/C MICE

Each animal in protocol A (Post -R) received a single whole body dose of 6 Gy and cucurbit extract was given each day for 7 days; Protocol B (Pre-R) received cucurbit extract each day for 7 days and 6 Gy dose on 8th day; Protocol C (Pre and Post -R) received a cucurbit extract each day for 7 days, and 6 Gy dose on 8th day and cucurbit extract each day for 7 days. Each bar represented the Mean ± SE of three experiments. Asterisk above a bar represented significance of difference with irradiated control at *p<0.05, **<0.01 and ***<0.001 respectively.

Treatment of mice with cucurbit extract (Ls, Cp, or Lc) for 7 days before irradiation (Protocol B) also revealed an increase in villi width ($43 \pm 1.9\mu\text{m}$, $38 \pm 1.4\mu\text{m}$ and $43 \pm 1.4\mu\text{m}$ respectively) as compared to irradiated control ($39 \pm 1.1\mu\text{m}$) though the increase was insignificant at $p < 0.05$. Pretreatment of mice with Ls, Lc or Cp for 7 days followed by irradiation after 24 h and again treatment with Ls, Cp or Lc for 7 days (Protocol C) did not display significant increase in villi width with respect to the irradiated group and remained at 39 ± 1.6 , 38 ± 1.4 and $37 \pm 1.5\mu\text{m}$.

Ls was most effective at increasing the villi width when administered under protocol A, where the increase was 2 and 6 μm more than protocol B and C respectively. Similarly, Cp rendered maximum increase in villi width under protocol A where the increase was 5 μm more than both the protocols B and C. Similarly, Lc displayed maximum increase in villi width under protocol A where the increase was 3 and 9 μm more than protocols B and C respectively.




FIG. 9: MICROSCOPIC IMAGE (40X) OF JEJUNUM OF MICE STAINED WITH HEMATOXYLIN AND EOSIN

- a) Un-irradiated control showing healthy villi.
 b) Irradiated control showing disrupted villi.
 c) R + Ls: Irradiation followed *Lgenaria siceraria* once a day for 7 days showing, recovering villi.
 d) R + Cp: Irradiation followed *Cucurbita pepo* once a day for 7 days showing, recovering villi.
 e) R + Lc: Irradiation followed *Luffa cylindrical* once a day for 7 days showing, recovering villi.

The newly formed cells produced by the stem cells, the crypts of Lieberkuhn have been reported to migrate towards the tip of villi reaching the top within about 72 h in mice and thereafter denude in the gut lumen. Irradiation of mice revealed a decrease in villi height (Fig. 7 and 9) and width (Fig. 8 and 9). Radiation damage has been reported to induce apoptosis in crypt cells of the intestine⁵³ and also affected the proliferation of crypt cells adversely^{54, 55}. Daily administration of cucurbit extracts (Ls, Cp or Lc) to mice countered the radiation induced injury to villi by increasing the height (Fig. 7) and width (Fig. 8) in all the three

experimental protocols with respect to irradiated control, however post-irradiation cucurbit treatment rendered maximum effect. Microscopic images of jejunum of mice displaying normal, disrupted and recovering villi have been displayed in Fig. 9. Ls has been reported to counter the radiation induced injury to villi in our previous study¹⁸. The Ls, Cp and Lc have been reported to contain anti-oxidants like flavonoids and other polyphenolic compounds^{30, 36, 56}. Hence, the anti-oxidative property of cucurbit extracts might have helpful in ameliorating ROS induced damage to crypts and hence were able to restore villi structural integrity.

CONCLUSION: The dietary cucurbits reduced the pro-inflammatory cytokines and increased the anti-inflammatory cytokines and the differences in the biochemical constituents of three cucurbits investigated here may be responsible for differential efficacy among themselves. The pre-irradiation administration was studied for prophylactic action against radiation induced inflammatory processes, but the anti-inflammatory molecules in the circulation dwindle with time and only little mitigation of radiation damage occurred. Post-irradiation treatment rendered maximum repair and recovery after radiation damage. Combined modality of pre- and post-irradiation administration of cucurbits yielded still better results in NAP decrease and TGF- β increase. Further studies are necessary to unravel these points. This study revealed the non-toxic anti-inflammatory potential of cucurbits in the therapy of chronic inflammatory disorders.

ACKNOWLEDGEMENT: This work was supported by the Indian Council of Medical Research (ICMR), Government of India vide grant number 5/9/1064/2012-NUT. The authors are sincerely grateful to ICMR for financial support and Amity University Uttar Pradesh (AUUP), Noida (UP) for providing necessary infrastructure facilities.

CONFLICT OF INTEREST: The authors have no conflict of interest.

REFERENCES:

1. Zhao X, Yang H, Jiang G, Ni M, Deng Y, Cai J, Li Z, Shen F and Tao X: Simvastatin attenuates radiation-induced tissue damage in mice. *Journal of Radiation Research* 2014; 55: 257-264.
2. Azzam EI, Jay-Gerin JP and Pain D: Ionizing radiation-induced metabolic oxidative stress and prolonged cell injury. *Cancer Letters* 2012; 327(1-2): 48-60.
3. François A, Milliat F, Guipaud O and Benderitter M: Inflammation and immunity in radiation damage to the gut mucosa. *Biomed Research International* 2013; 123241.
4. Kim JA, Jenrow KA and Brown SL: Mechanisms of radiation-induced normal tissue toxicity and implications for future clinical trials. *Radiation Oncology Journal* 2014; 32(3): 103-115.
5. Berbée M and Hauer-Jensen M: Novel drugs to ameliorate gastrointestinal normal tissue radiation toxicity in clinical practice: what is emerging from the laboratory? *Current Opinion in Support and Palliative Care* 2012; 6(1): 54-59.
6. Koukourakis MI: Radiation damage and radio protectants: new concepts in the era of molecular medicine. *British Journal of Radiology* 2012; 85(1012): 313-330.

7. Lloyd-Price J, Abu-Ali G and Huttenhower C: The healthy human microbiome. *Genome Medicine* 2016; 8: 51.
8. Conlon MA and Bird AR: The impact of diet and lifestyle on gut microbiota and human health. *Nutrients* 2015; 7(1): 17-44.
9. Hardy H, Harris J, Lyon E, Beal J and Foey AD: Probiotics, prebiotics and immunomodulation of gut mucosal defences: homeostasis and immunopathology. *Nutrients* 2013; 5(6): 1869-912.
10. Shih RH, Wang CY and Yang CM: NF-kappa B signaling pathways in neurological inflammation: A mini review. *Frontiers in Molecular Neuroscience* 2015; 8: 77.
11. Habtezion A, Nguyen LP, Hadeiba H and Butcher EC: Leukocyte trafficking to the small intestine and colon. *Gastroenterology* 2016; 150(2): 340-54.
12. Dhiman K, Gupta A, Sharma DK, Gill NS and Goyal A: A review of the medicinally important plants of the family cucurbitaceae. *Asian Journal of Clinical Nutrition* 2012; 4(1): 16-26.
13. Kumar D, Chandrul KK, Singh D, Sharma H, Pawar S and Mishra A: Pharmacognostic evaluation and phytochemical screening of fruits of *Lagenaria siceraria* (molina) Standly. *World Journal of Pharmaceutical Research* 2016; 5(5): 1373-1383.
14. Rajasree RS, Sibi PI, Francis F and William H: Phytochemicals of cucurbitaceae family - A review. *International Journal of Pharmacognosy and Phytochemical Research* 2016; 8(1): 113-123.
15. Chung So, Kim YJ and Park SU: An updated review of cucurbitacins and their biological and pharmacological activities. *EXCLI Journal* 2015; 14: 562-566.
16. Hur SJ, Kang SH, Jung HS, Kim SC, Jeon HS, Kim IH and Lee JD: Review of natural products actions on cytokines in inflammatory bowel disease. *Nutrition Research* 2012; 32(11): 801-16.
17. Sharma D, Rawat I and Goel HC: Anticancer and anti-inflammatory activities of some dietary cucurbits. *Indian Journal of Experimental Biology* 2015; 53(4): 216-21.
18. Sharma D, Goel HC and Chauhan S: Radioprotective potential of *Lagenaria siceraria* extract against radiation-induced gastrointestinal injury. *Applied Physiology, Nutrition, and Metabolism* 2016; 41(12): 1248-1254.
19. Liu Y, Fatheree NY, Mangalat N and Rhoads JM: Human-derived probiotic *Lactobacillus reuteri* strains differentially reduce intestinal inflammation. *American Journal of Physiology: Gastrointestinal and Liver Physiology* 2010; 299: G1087-G1096.
20. Khojasteh SMB, Sheikhzadeh F, Mohammadnejad D and Azami A: Histological, histochemical and ultrastructural study of the intestine of rainbow trout (*Oncorhynchus mykiss*). *World Applied Science Journal* 2009; 6: 1525-1531.
21. Turner MD, Nedjai B, Hurst T and Pennington DJ: Cytokines and chemokines: At the crossroads of cell signaling and inflammatory disease. *Biochim Biophys Acta* 2014; 843(11): 2563-2582.
22. Singh UP, Singh NP, Murphy EA, Price RL, Fayad R, Nagarkatti M and Nagarkatti PS: Chemokine and cytokine levels in inflammatory bowel disease patients. *Cytokine* 2016; 77: 44-9.
23. Shouval DS, Ouahed J, Biswas A, Goettel JA, Horwitz BH, Klein C, Muise AM and Snapper SB: Interleukin 10 receptor signaling: master regulator of intestinal mucosal homeostasis in mice and humans. *Advances in Immunology* 2014; 122: 177-210.
24. Rubtsov YP, Rasmussen JP, Chi EY, Fontenot J, Castelli L, Ye X, Treuting P, Siewe L, Roers A, Henderson WR Jr,

- Muller W and Rudensky A: Regulatory T cell-derived interleukin-10 limits inflammation at environmental interfaces. *Immunity* 2008; 28: 546-558.
25. Johnston CJC, Smyth DJ, Dresser DW and Maizels RM: TGF- β in tolerance, development and regulation of immunity. *Cellular Immunology* 2016; 299: 14-22.
 26. Gajewski T, Schreiber H and Fu YX: Innate and adaptive immune cells in the tumor microenvironment. *Nature immunology* 2013; 14: 1014-1022.
 27. Schaefer D, Kachikwu EL and McBride WH: Cytokines in radiobiological responses: a review. *Radiation Research* 2012; 178(6): 505-23.
 28. Di Maggio FM, Minafra L, Forte GI, Cammarata FP, Lio D, Messa C, Gilardi MC and Bravatà V: Portrait of inflammatory response to ionizing radiation treatment. *Journal of Inflammation (London)* 2015; 18: 12-14.
 29. Teresa Pinto A, Laranjeiro Pinto M, Patricia Cardoso A, Monteiro C, Teixeira Pinto M, Filipe Maia A, Castro P, Figueira R, Monteiro A, Marques M, Mareel M, Dos Santos SG, Seruca R Adolfo Barbosa M, Rocha S and José Oliveira M: Ionizing radiation modulates human macrophages towards a pro-inflammatory phenotype preserving their pro-invasive and pro-angiogenic capacities. *Scientific Reports* 2016; 6: 18765.
 30. Ghule BV, Ghante MH, Urganlawar AB and Yeoli PG: Analgesic and anti-inflammatory activity of *L. siceraria* (Mol.) Stand. fruit juice extract in rats and mice. *Pharmacognosy Magazine* 2006; 2: 232-238.
 31. Winkler C, Wirleitner B, Schroecksnadel K, Schennach H and Fuchs D: Extracts of Pumpkin (*Cucurbita pepo* Linn.) Seeds Suppress Stimulated Peripheral Blood Mononuclear Cells *in vitro*. *American Journal of Immunology* 2005; 1(1): 6-11.
 32. Kao TH, Huang CW and Chen BH: Functional components in *Luffa cylindrica* and their effects on anti-inflammation of macrophage cells. *Food Chemistry* 2012; 135(2): 386-395.
 33. Vieira AT, Teixeira MM and Martins FS: The role of probiotics and prebiotics in inducing gut immunity. *Frontiers in Immunology* 2013; 4: 445.
 34. Deshpande JR, Choudhry AA, Mishra MR, Meghre VS, Wadolkar SG and Dorle AK: Beneficial effects of *Lagenaria siceraria* (Mol.) Standley fruit epicarp in animal models. *Indian Journal of Experimental Biology* 2008; 46: 234-242.
 35. Erasto P and Mbwambo ZH: Antioxidant activity and HPTLC profile of *Lagenaria siceraria* fruits. *Tanzania Journal of Health Research* 2009; 11(2): 79-83.
 36. Irshad MI, Goel HC and Rizvi MMA: Phytochemical screening and high performance TLC analysis of some cucurbits. *Research Journal of Phytochemistry* 2010; 4(4): 242-247.
 37. Sharma D, Rawat I and Goel HC: Antioxidant and prebiotic potential of some dietary cucurbits. *Research Journal of Medicinal Plants* 2012; 6: 500-510.
 38. Rawat I, Sharma D and Goel HC: Antioxidant and anti-inflammatory potential of some dietary cucurbits. *Oxidants and Antioxidants in Medical Science* 2014; 3: 65-72.
 39. Monk JM, Hou TY and Chapkin RS: Recent advances in the field of nutritional immunology. *Expert Review Clinical Immunology* 2011; 7(6): 747-749.
 40. Moses T, Papadopoulou KK and Osbourn A: Metabolic and functional diversity of saponins, biosynthetic intermediates and semi-synthetic derivatives. *Critical Reviews in Biochemistry and Molecular Biology* 2014; 49(6): 439-462.
 41. Bellik Y, Boukraâ L, Alzahrani HA, Bakhotmah BA, Abdallah F, Hammoudi SM and Ouada MI: Molecular mechanism underlying anti-inflammatory and anti-allergic activities of phytochemicals: An update. *Molecules* 2013; 18: 322-353.
 42. Morrison DJ and Preston T: Formation of short chain fatty acids by the gut microbiota and their impact on human metabolism. *Gut Microbes* 2016; 7(3): 189-200.
 43. Den Besten G, Van Eunen K, Groen AK, Venema K, Reijngoud DJ and Bakker BM: The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *Journal of Lipid Research* 2013; 54: 2325-2340.
 44. Collins S and Reid G: Distant site effects of ingested prebiotics. *Nutrients* 2016; 8(9): 523.
 45. Vulevic J, Drakoularakou A, Yaqoob P, Tzortzis G and Gibson GR: Modulation of the fecal microflora profile and immune function by a novel trans-galacto oligosaccharide mixture (B-GOS) in healthy elderly volunteers. *American Journal of Clinical Nutrition* 2008; 88:1438-1446.
 46. Vulevic J, Juric A, Walton GE, Claus SP, Tzortzis G, Toward RE and Gibson GR: Influence of galacto-oligosaccharide mixture (B-GOS) on gut microbiota, immune parameters and metabolomics in elderly persons. *British Journal of Nutrition* 2015; 114: 586-595.
 47. Zhang M, Zhou Q, Dorfman RG, Huang X, Fan T, Zhang H, Zhang J and Yu C: Butyrate inhibits interleukin-17 and generates Tregs to ameliorate colorectal colitis in rats. *BMC* 2016; 16: 84.
 48. Hamer HM, Jonkers DM, Vanhoutvin SA, Troost FJ, Rijkers G, de Bruïne A, Bast A, Venema K and Brummer RJ: Effect of butyrate enemas on inflammation and antioxidant status in the colonic mucosa of patients with ulcerative colitis in remission. *Clinical Nutrition* 2010; 29(6): 738-44.
 49. Steidler L, Hans W, Schotte L, Neiryck S, Obermeier F, Falk W, Fiers W and Remaut E: Treatment of murine colitis by *Lactococcus lactis* secreting interleukin-10. *Science* 2000; 289: 1352.
 50. Goubeyre P, Denery S and Bodinier M: Probiotics, prebiotics, and synbiotics: impact on the gut immune system and allergic reactions. *Journal of Leukocyte Biology* 2011; 89: 685-695.
 51. D' Angelo C, Reale M and Costantini E: Microbiota and probiotics in health and HIV infection. *Nutrients* 2017; 9(6): 615.
 52. Larmonier CB, Shehab KW, Ghishan FK and Kiela PR: T-lymphocyte dynamics in inflammatory bowel diseases: Role of the microbiome. *BioMed Research International* 2015.
 53. Delgado ME, Grabinger T and Brunner T: Cell death at the intestinal epithelial front line. *Federation of European Biochemical Societies* 2016; 283: 2701-19.
 54. Bing SJ, Kim MJ, Ahn G, Im J, Kim DS, Ha D, Cho J, Kim A and Jee Y: Acidic polysaccharide of *Panax ginseng* regulates the mitochondria / caspase-dependent apoptotic pathway in radiation-induced damage to the jejunum in mice. *Acta histochemical* 2014; 116: 514-521.
 55. Zhao X, Yang H, Jiang G, Ni M, Deng Y, Cai J, Li Z, Shen F and Tao X: Simvastatin attenuates radiation-induced tissue damage in mice. *Journal of Radiation Research* 2014; 55: 257-264.
 56. Ojiako OA and Igwe CU: Nutritional and anti-nutritional compositions of *Cleome ruidosperma*, *Lagenaria siceraria*, and *Cucurbita maxima* seeds from Nigeria. *Journal of Medicinal Food* 2007; 10: 735-738.

How to cite this article:

Chauhan S, Sharma D and Goel HC: Therapeutic potential of Cucurbits against radiation induced inflammation and damage in mice gut. Int J Pharm Sci Res 2018; 9(2): 550-62.doi: 10.13040/IJPSR.0975-8232.9(2).550-62.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)