



Received on 08 July, 2017; received in revised form, 06 December, 2017; accepted, 06 January, 2018; published 01 March, 2018

A REVIEW ON TOMATO LYCOPENE

Nishkruti Mehta^{*1,2}, Pragnesh Patani³ and Indrajeet Singhvi²

Arihant School of Pharmacy and Bioresearch Institute¹, Adalaj, Gandhinagar - 382421, Gujarat, India.

Pacific Academy of Higher Education and Research University², Udaipur - 313003, Rajasthan, India.

A- One Pharmacy College³, Naroda, Ahmedabad - 382330, Gujarat, India.

Keywords:

Lycopene, Tomato,
Antioxidant, Chronic disease

Correspondence to Author:

Nishkruti Mehta

Arihant School of Pharmacy and
Bioresearch Institute, Adalaj,
Gandhinagar - 382421, Gujarat, India.

E-mail: nishkrutimehta@yahoo.com

ABSTRACT: Lycopene, a non-provitamin a carotenoid, found in tomatoes, processed tomato products and other fruits, due to which red to pink coloration is observed. Among all the dietary carotenoids, lycopene is one of the most potent naturally and abundantly occurring antioxidant. Lycopene has unique structural features and chemical features which contribute to specific biological properties and pharmacological activities. The antioxidant potential of lycopene may be responsible for its associated health benefits. Dietary consumption of tomatoes and tomato products rich in lycopene are reported to be associated with a decreased risk of chronic diseases, like cancer and various cardiovascular diseases. Further investigations will be required to understand and evaluate mechanism whereby lycopene and its metabolites are proven to possess biological activities in human. This review summarizes the current understanding of lycopene with respect to its role in human health and prevention of various diseases.

INTRODUCTION: Chemistry of Lycopene:

Lycopene is specifically an acyclic isomer of β -carotene, which is one of the members of carotenoid family. It is a lipid soluble antioxidant which is naturally synthesized by various plants, microorganisms and of certain algae and fungi but not synthesized by animals and humans¹. It acts as an accessory light-gathering pigment and protects the plants and microorganisms against the toxic effects of oxygen and light. Chemically, lycopene is a highly unsaturated hydrocarbon having 11 conjugated and 2 unconjugated double bonds. As a polyene, it undergoes *cis-trans* isomerization induced by light, thermal energy and chemical reactions (Fig. 1)^{2,3}.

Lycopene obtained from plant sources exists predominantly in an all-*trans* configuration, which is the most stable form, thermodynamically^{2,3}. Lycopene is present as an isomeric mixture, with 50% as *cis* isomers, in human plasma⁴. It is one of the most potent antioxidants,⁵⁻⁸ which has a singlet-oxygen-quenching ability twice as high as that of β -carotene and it has 10 times higher ability than that of α -tocopherol⁸.

Description of Lycopene:

Molecular Formula: C₄₀H₅₆

Molecular Weight: 536.85Da

Melting Point: 172-175 °C

Density: 0.889 gm./cm³

Stability: Sensitive to light, oxygen, high temperature, acids, catalysts and metal ions.

Solubility: Synthetic lycopene occurs in form of red to dark violet crystalline powder. It is insoluble in water and nearly insoluble in methanol and

	<p style="text-align: center;">DOI: 10.13040/IJPSR.0975-8232.9(3).916-23</p>
	<p style="text-align: center;">Article can be accessed online on: www.ijpsr.com</p>
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.9(3).916-23</p>	

ethanol whereas freely soluble in chloroform and tetra hydrofuran. It is sparingly soluble in solvents like ether, hexane, and vegetable oils².




FIG. 1: STRUCTURES OF TRANS AND CIS ISOMERS OF LYCOPENE

Mechanism of Action: The biological activities of carotenoids like β -carotene are related in general to their ability to form vitamin A within the body⁹. Lycopene doesn't have β -ionone ring structure, so it does not form Vitamin A¹⁰. Its biological effects in humans have therefore been attributed to mechanisms other than Vitamin A.

Two major hypotheses have been proposed to explain the anticarcinogenic and antiatherogenic activities of lycopene: nonoxidative and oxidative mechanisms. Among the nonoxidative mechanisms, the anticarcinogenic effects of lycopene have been reported to be due to regulation of gap-junction communication in mouse embryo fibroblast cells^{11, 12}. Lycopene is hypothesized to suppress carcinogen-induced phosphorylation of regulatory proteins such as p53 and Rb anti-oncogenes and stop cell division at the G0-G1 cell cycle phase¹³.

Astorg *et al.*,¹⁴ demonstrated that lycopene-induced alteration of the liver metabolizing enzyme like cytochrome P450E1, was initial *in-vitro* evidence also indicated that lycopene was found to reduce cellular proliferation induced by insulin-like growth factors, which are potent mitogens, in various cancer cell lines¹⁵. Regulating intrathymic T-cell differentiation (immunomodulation) was predicted to be the underlying mechanism for suppression of mammary tumor growth by lycopene treatments in SHN retired mice^{16, 17}. It has also been demonstrated that lycopene acts as a

hypo-cholesterolemic agent by inhibiting HMG-CoA (3-hydroxy-3-methylglutaryl- coenzyme A) reductase enzyme¹⁸. It has been hypothesized that lycopene possesses anti-carcinogenesis and anti-atherogenesis by protecting critical cellular biomolecules, like lipids, proteins, lipoproteins and DNA^{19, 20, 21}.

In healthy volunteers, lycopene or tomato free diets resulted in loss of lycopene and showed increase in lipid oxidation²², whereas dietary supplementation or dietary consumption with lycopene for 1 week, reported in increased serum lycopene levels and reduced endogenous levels of lipid oxidation, protein oxidation, oxidation of lipoproteins and DNA^{19, 20}. Patients having prostate cancer were reported to have low levels of lycopene and high levels of oxidation of serum lipids and proteins²³.

Pharmacological Activities: It has been reported from various studies that a tomato rich diet and products containing tomatoes rich in lycopene have protective effects against number of chronic diseases by removing or reducing oxidative stress. Dietary lycopene protected lipids, DNA and proteins against oxidation. Such oxidized products are considered to possess key role in cancer and various chronic diseases and also have been reported to increase in the chronic disease condition significantly. In addition, dietary intake or consumption of lycopene reduces the risk of various chronic diseases and the serum and tissue levels are inversely related to the risk of these diseases. Various studies have been reported which show an inverse correlation with tomatoes and lycopene-rich diets and the incidence of several cancers and CHD. The beneficial effects are attributed due to the anti-oxidant properties but several other mechanisms like alteration or modulation of intercellular gap junction, communication, hormonal and immune system, metabolic pathways may also be involved^{22, 24, 25, 26}.

Role of Lycopene in Cancer: The affirmation in support of lycopene in prevention of various chronic diseases comes from epidemiological studies²⁷⁻³¹, studies using human cancer cell lines³¹⁻³³ pre-clinical studies^{34, 35} and also human clinical trial³⁶. From all the different types of cancer, the key role of lycopene for prevention of prostate cancer has been widely studied.

Prostate cancer is one of the most common malignancy and leading causes of death occurring in male, worldwide. Although, age and genetic factors are also the leading and important risk factors, environmental exposures, like diet are increasingly being associated with the development of disease. A follow up met analysis of 72 various studies reported that intake of lycopene as well as level of serum lycopene were inversely related to various cancers like breast cancer, prostate cancer, cervical cancer, ovarian cancer, liver cancers and other organ sites. Since then, various studies reported that with increased lycopene intake and serum levels of lycopene the risk of cancers were reported to reduce significantly³⁶⁻³⁸.

Rao et al.,³⁹ demonstrated the level of oxidative stress and antioxidants in the patients of prostate cancer. The results reported the notable differences in the level of prostate specific antigen (PSA), serum carotenoids levels and biomarkers of oxidation. Although there were no significant alterations in the levels of lutein, β -carotene, Vitamin A and E, cryptoxanthin, between the subjects with cancer and their controls, lycopene levels were noted reduced significantly in cancer patients. The PSA levels were significantly increased in the patients with cancer, along with elevated levels of proteins and lipid oxidation reporting the increased levels of oxidative stress in the patients with cancer.

Furthermore, lycopene is utilized to reduce the lipid oxidation suggesting that reduced lycopene levels in cancer cases were caused due to the disease rather than the cause of the disease³⁹. Lycopene was reported to reduce the PSA levels and also the development of prostate cancer in the patients newly diagnosed with cancer, who were receiving 15 mg of lycopene for 3 weeks daily, prior to radical prostatectomy^{36,40}.

Studies indicated that daily consumption of food rich in lycopene has been reported to be associated with 30 to 40 % reduction in the risk of prostate cancer²⁸. According to other study it was demonstrated that consuming tomato sauce, providing 30 mg lycopene for 3 weeks daily prior to prostatectomy in male diagnosed with prostate cancer showed significant elevation in serum and prostate levels of lycopene⁴¹. Oxidative damage to

DNA was reduced and serum PSA declined significantly by 20 % with lycopene treatment.

In pilot studies with the patients having prostate cancer, it was illustrated that daily ingestion of lycopene obtained from tomato sauce or extract of tomato inhibited tumor growth and invasiveness and reduced the growth of prostate cancer probably by up regulation of connexin 43 (Cx43) (tumor suppressor protein⁴²).

Similarly, Heber and Lu⁴³ demonstrated that a gene, connexin 43, whose expression was up regulated by lycopene and allowed direct intercellular gap junctional communication (GJC) was identified from preclinical and clinical studies. It was also found that there is deficiency of GJC in many tumors of humans and its restoration or up regulation is associated with decreased proliferation. In the same manner it is confirmed from various preclinical studies that there is the inverse association between lycopene obtained from diet and growth of both spontaneous and transplanted tumors⁴⁴. It has also been reported to inhibit the proliferation of various cancer cells like lungs, breast, and endometrium and also suppress insulin like growth factor I stimulated growth. Moreover, it inhibits growth and development of KB-I human oral tumor cells⁴⁵ and C-6 glioma cells transplanted into rats⁴⁶.

Nkondjock et al.,⁴⁷ demonstrated that tomato rich diet and consumption of tomato based products with increased lycopene content may help to decrease the risk of cancer of pancreas. From various studies it has been investigated that dietary lycopene (10ppm) showed significant reduction in the lipid and protein oxidation and illustrated that it had an apparent protective effect against azoxymethane induced colonic preneoplastic lesions in rats²⁶.

Overall epidemiological analysis, *in-vitro* tissue culture studies, preclinical studies and some clinical studies involving human intervention demonstrated that increased consumption of lycopene would indicate increased circulatory and tissue lycopene levels. From *in-vivo* studies it was reported, that lycopene can play a key role as an anti-oxidant potentially and protect cells against oxidative damage and thereby reduces or prevents

the risk of development of various cancers. Further investigations are required for detailed understanding of the underlying mechanisms.

Role of Lycopene in Cardiovascular Disease:

The normal function of the cardiovascular system involving heart and blood vessels is remarkably affected by cardiovascular disease (CVD), which is the major and leading cause of deaths in western world and an important contributor of morbidity and mortality in developing countries.

As per reports of the World Health Organization (WHO) ⁴⁸, it was found that CVD is the world's largest killer, claiming 17.1 million lives a year. Tobacco use, unhealthy diet, physical inactivity and high intake of alcohol are the known risk factors leading to CVD. Plasma low density lipoprotein (LDL) is the major risk factor of CVD. Elevation in oxidation of LDL has been proposed to be associated with elevating risk of atherosclerosis and coronary heart disease. Cumulative evidences in literature support the role of lycopene for the prevention of cardiovascular diseases (CVD) ⁴⁹⁻⁵².

Arab and Steck ⁵² reported that it has a protective effect against intimal wall thickness and myocardial infarction. It was proposed that the protective effect may be due to some other mechanism beside its antioxidant effect. Similarly, Rissanen *et al.*, ⁵³ concluded that low plasma lycopene concentration is associated with early atherosclerosis and increased intima media thickness of common carotid artery wall (CCA-IMT). Serum cholesterol levels have been used as a biomarker for the risk of CHD traditionally. Oxidation of the circulating low density lipoprotein (LDL), which is carrying cholesterol into the blood stream, to oxidized LDL (LDLox) is also thought to play a key role in the pathogenesis of arteriosclerosis which is the main disorder leading to heart attack and ischemic strokes ⁵⁴⁻⁵⁶.

Owing to its lipophilic nature, lycopene concentrates in LDL and VLDL fractions and not in HDL fractions ⁵⁷ It has been reported to significantly reduce the levels of oxidized LDL and lipid peroxidation in the subjects who consume tomato sauce, tomato juice and lycopene oleo-resin capsules and also protects LDL from oxidation *in-vitro*.⁵⁸ The *in-vivo* and *in-vitro* studies, demonstrated that lycopene can suppress cholesterol synthesis by

inhibiting 3-hydroxy-3-methyl glutaryl coenzyme A (HMG Co A) ^{52, 59}.

Ahuja *et al.*, ⁶⁰ reported that the diet high in olive oil and rich in lycopene decreased the risk of CHD by improving the serum lipid profile as compared to high carbohydrate and low fat diet. Although from the epidemiological studies, convincing evidence in support of the lycopene's protective role in CVD has been provided. Well controlled human intervention studies are required to be conducted for validation of these observations in future.

Role of Lycopene in Other Diseases: Researchers have started to investigate role of lycopene in various human diseases, because of its recognition as a potent antioxidant and its preventive role in oxidative stress mediated chronic diseases.

Osteoporosis: Oxidative stress may be one of the contributing factors involved in the pathogenesis of skeletal system like development of osteoporosis, which is the most common metabolic bone disease. Lycopene has a stimulatory effect on cell proliferation and the differentiation marker alkaline phosphatase of osteoblasts as well as inhibitory effects on osteoclasts formation and resorption. There have been results of a possible decrease in bone turnover and oxidative stress markers and an increase in antioxidant status in postmenopausal women taking tomato juice or lycopene capsules. Thus lycopene plays a key role in maintaining bone health and provides dietary alternative to drug therapy especially for women with the risk of this disease ⁶¹⁻⁶⁵.

Male Infertility: It is the most common male reproductive disorder is now being associated with oxidative damage of the sperm which leads to the loss of its quality and functionality. Significant levels of ROS are detectable in the semen of up to 25 % of infertile men whereas fertile men do not produce detectable levels of ROS in their semen ^{66, 67}. Researchers are beginning to investigate the role of lycopene in protecting sperm from oxidative damage leading to infertility. Studies show that men with antibody mediated infertility were found to have lower serum lycopene levels than their fertile controls ⁶⁸. According to other study, a significant increase in serum lycopene concentration

and improvement in sperm motility, sperm motility index, sperm morphology and functional sperm concentration was reported in infertile men when administered with 8 mg lycopene for 12 months. Furthermore, it was found lycopene treatment resulted in 36 % successful pregnancies.

Hypertension: Hypertension, a 'silent killer,' is a disorder which is asymptomatic until a more advanced and a fatal stage is reached. Lycopene's antioxidant property has attracted scientific research into its protective role in hypertension. As per recent study it was demonstrated that lycopene supplementation of 15 mg/ day for 8 weeks remarkably decreased systolic blood pressures from baseline values to 144 mm Hg to 134 mm Hg in mildly hypertensive subjects.

In another study a significant reduction in plasma lycopene was observed in the hypertensive patients compared to normal subjects. As the antioxidants are important in the management of hypertension, a 'dietary approach for controlling hypertension (DASH)', diet containing substantially higher levels of lycopene along with other carotenoids, polyphenols, flavanols, flavanones and flavan-3-ols is recommended⁶⁹⁻⁷².

Neurodegenerative Diseases: Rao and Balachandran⁷³ suggested that lycopene plays important role in neurodegenerative diseases like Alzheimer's disease. It was reported that lycopene was able to cross the blood brain barrier and be present in central nervous system (CNS) in low concentration. Significant reduction in the levels of lycopene was reported in patients suffering from Parkinson's disease and vascular dementia⁷⁴. Similarly, Saganuma *et al.*,⁷⁵ reported that tomato ingestion might serve as a preventive therapy against neurodegenerative diseases such as Parkinson's disease caused by 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine (MPTP) and other environmental toxins. Moreover, it was also suggested to provide protection against amyotrophic lateral sclerosis (ALS) disorder in humans⁷⁶.

Respiratory Infections: Intake of tomatoes was also inversely and significantly associated with respiratory infections⁷⁷. A study demonstrated that lycopene has a protective role for prevention of emphysema in a mouse model. At a conference held to deliberate on the role of processed tomatoes

in human health, data was provided for the lycopene's protective role for emphysema's prevention in Japanese population⁷⁸.

Cataract: It was reported that persons with a high intake of carotene showed reduced incidence of risk of cataract⁷⁹ and the relationship between nuclear cataract and intakes of α - carotene, β -carotene, lutein, lycopene and cryptoxanthin stratifying by gender and by regular multivitamin use⁸⁰. It has been confirmed that lycopene prevents cataract genesis *in-vivo* and *in-vitro* due to its antioxidant potentials⁸¹. In previous studies, it was found that lycopene prevented sugar induced diabetic cataract⁸².

Erythema: Lycopene plays an important role in the protection against photooxidative processes. It acts as singlet molecular oxygen and peroxy radicals scavengers. It can also interact synergistically with other antioxidants. Administration of tomato paste daily for 10 weeks, protected against UV light induced erythema on the dorsal skin⁸³. However, it is completely depleted from skin upon exposure to solar radiation⁸⁴ and undergoes oxidative or enzymatic cleavage to form apocarotenoids⁸⁵.

Diarrhea: In a prospective study, Fawazi *et al.*,⁸⁶ found that an intake of tomatoes for 2-3 days compared with zero days has been associated with significant reduction in mortality (48%) and with a decreased risk of death associated with diarrhea.

HIV: Several studies indicate that lycopene plays important role in the human organism's natural defense mechanism which gives protection from the harmful oxidizing agents. Lower serum lycopene levels were also reported in human immuno deficiency virus (HIV) positive women and children^{87, 88}. Further investigations can explore lycopene's role in various human diseases like diabetes, rheumatoid arthritis, periodontal diseases and inflammatory disorders³⁸. Antioxidant potential of lycopene are opening up new applications in pharmaceutical; nutraceutical and cosmeceutical products and can inhibit the progression or development of many human diseases at an initial stage and may improve the quality of life⁸⁹.

Diabetes: Treatment of lycopene (1, 2 and 4 mg/kg; p.o.) in streptozotocin-induced diabetic rats

has significantly attenuated cognitive deficit, increased acetyl cholinesterase activity, oxidative nitrosative stress and inflammation⁹⁰.

Memory and Cognitive Behavior: The treatment of lycopene using 3-nitropropionic acid-induced rats has significantly improved the memory and restored glutathione system functioning⁹¹. Akbaraly *et al.*,⁹² also suggested that low plasma lycopene levels could contribute to cognitive impairment.

CONCLUSION: Lycopene is a naturally occurring substance found in many fruits and vegetables including tomatoes. At least 85% of lycopene comes from consumption of tomatoes and tomato based products like soups, sauces, juice, ketchups, salad etc. Carotenoids in general have undergone a number of research studies as to their possible benefits against diseases, among other health issues. Owing to the numerous pharmacological actions of tomato lycopene as discussed in above section, and associated health benefits, tomato consumption should be promoted in daily diet.

ACKNOWLEDGEMENT: Nil

CONFLICTS OF INTEREST: Nil

REFERENCES:

1. Paiva SAR and Russell RM: β -Carotene and other carotenoids as antioxidants. *J Am Coll Nutr*, 1999; 18(5): 426-433.
2. Nguyen ML and Schwartz SJ: Lycopene: chemical and biological properties. *Food Technol* 1999; 53: 38-45.
3. Zechmeister L, LeRosen AL, Went FW and Pauling L: Prolycopene, a naturally occurring stereoisomer of lycopene. *Proc Natl Acad Sci USA* 1941; 21: 468-74.
4. Clinton SK, Emehiser C, Schwartz SJ, Bostwick DG, Williams AW, Moore BJ, *et al.*, cis-trans lycopene isomers, carotenoids and retinol in the human prostate. *Cancer Epidemiol Biomarkers Prev* 1996; 5: 823-33.
5. Miller NJ, Sampson J, Candeias LP, Bramley PM and Rice-Evans CA: Antioxidant activities of carotenes and xanthophylls. *FEBS Lett* 1996; 384: 240-6.
6. Mortensen A and Skibsted LH: Relative stability of carotenoid radical cations and homologue tocopheroxyl radicals. A real time kinetic study of antioxidant hierarchy. *FEBS Lett* 1997; 417: 261-6.
7. Woodall AA, Lee SWM, Weesie RJ, Jackson MJ and Britton G: Oxidation of carotenoids by free radicals: relationship between structure and reactivity. *Biochim Biophys Acta* 1997; 1336: 33-42.
8. DiMascio P, Kaiser S and Sies H: Lycopene as the most effective biological carotenoid singlet oxygen quencher. *Arch Biochem Biophys* 1989; 274: 532-8.
9. Clinton SK: Lycopene: chemistry, biology and implications for human health and disease. *Nutr Rev* 1998; 56: 35-51.
10. Stahl W, Schwarz W, Sundquist AR and Sies H: cis-trans isomers of lycopene and beta-carotene in human serum and tissues. *Arch Biochem Biophys* 1992; 294: 173-7.
11. Zhang LX, Cooney RV and Bertram JS: Carotenoids enhance gap junctional communication and inhibit lipid peroxidation in C3H/10T1/2 cells: relationship to their cancer chemopreventive action. *Carcinogenesis* 1991; 12: 2109-14.
12. Zhang LX, Cooney RV and Bertram JS: Carotenoids up-regulate connexin 43 gene expression independent of their provitamin A or antioxidant properties. *Cancer Res* 1992; 52: 5707-12.
13. Matsushima NR, Shidoji Y, Nishiwaki S, Yamada T, Moriwaki H and Muto Y: Suppression by carotenoids of microcystin-induced morphological changes in mouse hepatocytes. *Lipids* 1995; 30: 1029-34.
14. Astorg P, Gradelet S, Berges R and Suschetet M: Dietary lycopene decreases the initiation of liver preneoplastic foci by diethyl nitrosamine in the rat. *Nutr Cancer* 1997; 29(1): 60-8.
15. Levy J, Bosin E, Feldmen B, Giat Y, Miinster A, Danilenko M, *et al.*, Lycopene is a more potent inhibitor of human cancer cell proliferation than either α carotene or β -carotene. *Nutr Cancer* 1995; 24: 257-66.
16. Nagasawa H, Mitamura T, Sakamoto S and Yamamoto K: Effects of lycopene on spontaneous mammary tumour development in SHN virgin mice. *Anticancer Res* 1995; 15: 1173-8.
17. Kobayashi T, Iijima K, Mitamura T, Toriizuka K, Cyong JC and Nagasawa H: Effects of lycopene, a carotenoid, on intrathymic T cell differentiation and peripheral CD4/CD8 ratio in a high mammary tumor strain of SHN retired mice. *Anticancer Drugs* 1996; 7: 195-8.
18. Fuhramn B, Elis A and Aviram M: Hypo-cholesterolemic effect of lycopene and α - carotene is related to suppression of cholesterol synthesis and augmentation of LDL receptor activity in macrophage. *Biochem Biophys Res Commun* 1997; 233: 658-62.
19. Agarwal S and Rao AV: Tomato lycopene and low density lipoprotein oxidation: a human dietary intervention study. *Lipids* 1998; 33: 981-4.
20. Rao AV and Agarwal S: Bioavailability and antioxidant properties of lycopene from tomato products. *Nutr Cancer*, 1998a; 31: 199-203.
21. Pool-Zobel BL, Bub A, Muller H, Wollowski I and Rechkemmer G: Consumption of vegetables reduces genetic damage in humans: first result of a human intervention trial with carotenoid-rich foods. *Carcinogenesis* 1997; 18: 1847-50.
22. Rao AV and Agarwal S: Effect of diet and smoking on serum lycopene and lipid peroxidation. *Nutr Res* 1998; 18: 713-21.
23. Rao AV, Fleshner N and Agarwal S: Serum and tissue lycopene and biomarkers of oxidation in prostate cancer patients: a case control study. *Nutr Cancer* 1999; 33: 159-64.
24. Agarwal S and Rao AV: Tomato lycopene and its role in human health and chronic diseases. *Can Med Assoc J*, 2000; 163(6): 739-744.
25. Wang M, Dhingra K, Hittelman WN, Liehr JG, de Andrade M and Donghui L: Lipid peroxidation-induced putative malondialdehyde- DNA adducts in human breast tissues, *Cancer Epiemiol Biomarkers Prev*, 1996; 5: 705-710.

26. Rao AV and Agarwal S: Role of lycopene as antioxidant carotenoid in the prevention of chronic diseases: a review. *Nutr Res* 1999; 19: 305-23.
27. Giovannucci E: Tomatoes, tomato-based products, lycopene and cancer: review of the epidemiologic literature. *Journal of the National Cancer Institute* 1999; 91(4): 317-331
28. Giovannucci E, Rimm E, Liu Y, Stamfer M and Willet WA: Prospective study of tomato products, lycopene and prostate cancer risk. *Journal of the National Cancer Institute* 2002; 94: 391-398.
29. Giovannucci E, Ascherio A, Rimm EB, Stampfer MJ, Colditz GA and Willett WC: Intake of carotenoids and retinol in relation to risk of prostate cancer. *J Natl Cancer Inst*, 1995; 87: 1767-1776.
30. LaVecchia C: Mediterranean epidemiological evidence on tomatoes and the prevention of digestive tract cancers. *Proc Soc Exp Bio Med*, 1997; 218: 125-12.
31. Hall TJ, Schaeublin M, Fuller K and Chambers TJ: The role of oxygen intermediates in osteoclastic bone resorption. *Biochem Biophys Res Commun*, 1995; 207: 280-287.
32. Kotake-Nara E, Kushiro M, Zhang H, Sugawara T, Miyashita K and Nagao A: Carotenoids affect proliferation of human prostate cancer cells. *J Nutr*, 2001; 131: 3303-3306.
33. Kim L, Rao AV and Rao LG: Effect of lycopene on Prostate LNCaP cancer cells in culture. *J Med Food*, 2002; 5(4): 181-187.
34. Heath E, Seren S, Sahin K and Kucuk O: The role of tomato lycopene in the treatment of prostate cancer, In: Rao AV, editor, *Tomatoes, lycopene and human health*, Caledonian Science Press, Scotland, 2006: 127-140.
35. Jain CK, Agarwal S and Rao AV: The effect of dietary lycopene on bioavailability, tissue distribution, in-vivo antioxidant properties and colonic preneoplasia in rats. *Nutr Res* 1999; 19: 1383-91.
36. Kucuk O, Sarkar FH and Sakr W: Phase II randomized clinic trial of lycopene supplementation before radical prostatectomy. *Cancer Epidemiol Biomarkers Prev*, 2001; 10: 861-868
37. Rao AV and Rao LG: Lycopene and human health. *Curr Top Nutr Res*, 2004; 2: 127-36.
38. Rao AV, Ray MR and Rao LG: Lycopene. *Adv Food Nutr Res*, 2006; 51: 99-164.
39. Rao AV, Fleshner N and Agarwal S: Serum and tissue lycopene and biomarkers of oxidation in prostate cancer patients: a case-control study. *Nutr Cancer*, 1999; 33: 159-164.
40. Kucuk O and Wood Jr DP: Response of hormone refractory prostate cancer to lycopene. *J Urol*, 2002; 167: 651.
41. Bowen P, Chen L, Stacewicz-Sapuntzakis M, Duncan C, Sharifi R, Ghosh L, Kim HS, Christov-Tzelkov K and van Breemen R: Tomato sauce supplementation and prostate cancer: lycopene accumulation and modulation of biomarkers of carcinogenesis. *Exp Biol Med* 2002; 227: 886-893
42. Kucuk O, Sarkar FH, Djuric Z, Sakr W, Pollak MN, Khachik F, Banerjee M, Bertram JS and Wood DP Jr: Effects of lycopene supplementation in patients with localized prostate cancer. *Exp Biol Med*, 2002; 227: 881-885.
43. Heber D and Lu QY: Overview of mechanisms of action of lycopene. *Exp Biol Med*, 2002; 227: 920-923.
44. Sharoni Y, Giron E, Rise M and Levy J: Effects of lycopene enriched tomato oleoresin on 7, 12-dimethylbenz [a] anthracene-induced rat mammary tumors. *Cancer Detect Prev*, 1997; 21: 118-123.
45. Livny O, Kaplan I, Reifen R, Polak-Charcon S, Madar Z and Schwartz B: Lycopene inhibits proliferation and enhances gap-junction communication of KB-1 human oral tumor cells. *J Nutr*, 2002; 132: 3754-3759.
46. Wang CJ, Chou MY and Lin JK: Inhibition of growth and development of the transplantable C-6 glioma cells inoculated in rats by retinoids and carotenoids. *Canc Lett*, 1989; 48: 135-142.
47. Nkondjock A, Ghadirian P, Johnson KC and Krewski D: And the Canadian cancer registries epidemiology research group, Dietary intake of lycopene is associated with reduced pancreatic cancer risk. *J Nutr*, 2005; 135: 592-597.
48. WHO, Cardiovascular Diseases: World Heart Day; World Health Organization (World Heart Federation), Geneva, Switzerland, 2009.
49. Fuhramn B, Elis A and Aviram M: Hypo-cholesterolemic effect of lycopene and β -carotene is related to suppression of cholesterol synthesis and augmentation of LDL receptor activity in macrophage. *Biochem Biophys Res Commun*, 1997; 233: 658-662.
50. Rissanen T, Voutilainen S, Nyyssonen K, Salonen R and Salonen JT: Low plasma lycopene concentration is associated with increased intima-media thickness of the carotid artery wall. *Arterioscler Thromb Vasc Biol*, 2000; 20: 2677-2687.
51. Rao AV: Lycopene, tomatoes and the prevention of coronary heart disease, *Exp Biol Med*, 2000; 227: 908-913.
52. Arab L and Steck S: Lycopene and cardiovascular disease. *Am. J Clin Nutr*, 2000; 71(S-1): 1691S-1695S.
53. Rissanen T, Voutilainen S, Nyyssonen K and Salonen JT: Lycopene, atherosclerosis and coronary heart disease. *Exp Biol Med*, 2002; 227: 900-907.
54. Parthasarathy S, Steinberg D and Witztum JL: The role of oxidized low density lipoproteins in pathogenesis of atherosclerosis. *Ann Rev Med*, 1992; 43: 219-225.
55. Witztum JL: The oxidation hypothesis of atherosclerosis. *Lancet*, 1994; 344: 793-796.
56. Heller FR, Descamps O and Hondekjijn JC: LDL oxidation: therapeutic perspectives. *Atherosclerosis*, 1998; 137: S25-S31.
57. Stahl W and Sies H: Lycopene: a biologically important carotenoid for human? *Arch Biochem Biophys*, 1996; 336: 1-9.
58. Safari MR: Effects of lycopene on the susceptibility of low density lipoproteins to oxidative modification. *Iran J Phar Res*, 2007; 6(3): 173-177.
59. Blum A, Monir M, Wirsansky I and Ben Arzi S: The beneficial effects of tomatoes. *Eur J Intern Med*, 2005; 16: 402-404.
60. Ahuja KDK, Pittaway JK and Ball MJ: Effects of olive oil and tomato lycopene combination on serum lycopene, lipid profile and lipid oxidation. *Nutr* 2006; 22: 259-265.
61. Kim L, Rao AV and Rao LG: Lycopene II-Effect on osteoblasts: the carotenoid lycopene stimulates cell proliferation and alkaline phosphatase activity of SaOS-2 cells. *J Med Food*, 2003; 6(2): 79-86.
62. Park CK, Ishimi Y, Ohmura M, Yamaguchi M and Ikegami S: Vitamin A and carotenoids stimulate differentiation of mouse osteoblastic cells. *J Nutr Sci Vitaminol*, 1997; 43: 281-96.
63. Ishimi Y, Ohmura M, Wang X, Yamaguchi M and Ikegami S: Inhibition by carotenoids and retinoic acid of osteoclast-like cell formation induced by bone-resorbing agents *in-vitro*. *J Clin Biochem Nutr*, 1999; 27: 113-122.

64. Rao LG, Krishnadev N, Banasikowska K and Rao AV: Lycopene I- Effect on osteoclasts: Lycopene inhibits basal and parathyroid hormone stimulated osteoclast formation and mineral resorption mediated by reactive oxygen species in rat bone marrow cultures. *J Med Food*, 2003; 6(2): 69-78.
65. Rao LG, Mackinnon ES, Josse RG, Murray TM, Strauss A and Rao AV: Lycopene consumption decreases oxidative stress and bone resorption markers in postmenopausal women. *Osteoporosis Int*, 2007; 18(1): 109-115.
66. Iwasaki A and Gagnon C: Formation of reactive oxygen species in spermatozoa of infertile patients. *Fertil Steril*, 1992; 57: 409-416.
67. Zini A, de Lamirande E and Gagnon C: Reactive oxygen species in semen of infertile patients: levels of superoxide dismutase and catalase-like activities in seminal plasma and spermatozoa. *Int J Androl*, 1993; 16: 183-188.
68. Palan P and Naz R: Changes in various antioxidant levels in human seminal plasma related to immunofertility. *Arch Androl*, 1996; 36: 139-43.
69. Paran E and Engelhard Y: Effect of Lyc-O-Mato, standardized tomato extract on blood pressure, serum lipoproteins plasma homocysteine and oxidative stress markers in grade 1 hypertensive patients. *Am J Hypertens*, 2001; 14: 141A Abstract P-333.
70. Paran E: Reducing hypertension with tomato lycopene, In: Rao AV, editor, *Tomatoes, lycopene and human health*, Caledonian Science Press, Scotland, 2006; 169-182.
71. Moriel P, Sevanian A, Ajzen S, Zanella MT, Plavnik FL, Rubbo H and Abdal DSP: Nitric oxide, cholesterol oxides and endothelium dependent vasodilation in plasma of patients with essential hypertension. *Braz J Med Biol Res*, 2002; 35: 1301-1309.
72. Most MM: Estimated phytochemical content of the dietary approaches to stop hypertension (DASH) diet is higher than in the control study diet. *Am Diet Assoc*, 2004; 104: 1725-1727.
73. Rao AV and Balachandran B: Role of oxidative stress and antioxidants in neurodegenerative disease. *Nutr Neurosci*, 2003; 5(5): 291-309.
74. Foy CJ, Passmore AP, Vahidassr MD, Young IS and Lawson JT: Plasma chain-breaking antioxidants in Alzheimer's disease, vascular dementia and Parkinson's disease. *QJM*, 1999; 92: 39-45.
75. Suganuma H, Hirano T, Arimoto Y and Inakuma T: Effect of tomato intake on striatal monoamine level in a mouse model of experimental parkinson's disease. *J Nutr Sci Vitiminol*, 2002; 48: 251-254.
76. Longnecker MP, Kamel F, Umbach DM, Munsat TL, Shefner JM, Lansdell LW and Sandler DP: Dietary intake of calcium, magnesium and antioxidants in relation to risk of amyotrophic lateral sclerosis. *Neuroepidemiology*, 2000; 19: 210-216.
77. Fawzi W, Herrera MG and Nestel P: Tomato intake in relation to mortality and morbidity among Sudanese children. *J Nutr*. 2000; 130(10): 2537-2542.
78. Rao AV and Rao LG: Carotenoids and human health. *Pharm Res*, 2007; 55: 207-216.
79. Cumming RG, Mitchell P and Smith W: Diet and cataract: The Blue Mountains Eye studies, *Ophthalmology*, 2000; 107: 450-456.
80. Mares-Perlman JA, Brady WE, Klein BE, Klein R, Haus GJ and Palta M: Diet and nuclear lens opacities. *Am J Epidemiol*, 1995; 41: 322-334.
81. Gupta SK, Trivedi D, Srivastava S, Joshi S, Halder N and Verma SD: Lycopene attenuates oxidative stress induced experimental cataract development: an in-vitro and in-vivo study. *Nutr*, 2003; 19: 794-799.
82. Mohanty I, Joshi S, Trivedi D, Srivastava S and Gupta SK: Lycopene prevents sugar-induced morphological changes and modulates antioxidant status of human lens epithelial cells. *Br J Nutr*, 2002; 88: 347-354.
83. Stahl W, Heinrich U, Wiseman S, Eichler O, Sies H and Tronnier H: Dietary tomato paste protects against ultraviolet light induced erythema in humans. *J Nutr*, 2001; 131: 1449-1451.
84. Ribago-Mercado JD, Garmyn M, Gilchrest BA and Russell RM: Skin lycopene is destroyed preferentially over β -carotene during UV irradiation in humans. *J Nutr*, 1995; 125: 1854-1859.
85. Tang G, Wang XD, Russel RM and Krinsky NI: Characterization of β -apo-13-carotenone and β -apo-14'-carotenal as enzymatic products of the excentric cleavage of β -carotene. *Biochem*, 1991; 30: 9829-9834.
86. Fawzi W, Herrera MG and Nestel P: Tomato intake in relation to mortality and morbidity among Sudanese children. *J Nutr*, 2000; 130(10): 2537-2542
87. Coodley GO, Coodley MK and Nelson HD: Micronutrients in HIV infected women. *J Womens Health*, 1995; 4: 303-311.
88. Periquet BA, Jammes NM, Lambert WE, Tricoire J, Moussa MM, Garcia J, Ghisolfi J and Thouvenot J: Micronutrients levels in HIV-1-infected children. *AIDS*, 1995; 9: 887-893.
89. Stahl W: Tomato lycopene in photoprotection and skin care, In: Rao AV, editor, *Tomatoes, lycopene and human health*, Caledonian Science Press, Scotland, 2006: 199-211.
90. Kuhad A, Sethi R and Chopra K: Lycopene attenuates diabetes associated cognitive decline in rats. *Life Sci*, 2008; 83: 128-134.
91. Kumar P and Kumar A: Effect of lycopene and epigallocatechin-3- gallate against 3-nitropropionic acid induced cognitive dysfunction and glutathione depletion in rat: A novel nitric oxide mechanism. *Food Chem Toxicol*, 2009; 47: 2522-2530.
92. Akbaraly NT, Faure H, Gourlet V, Favier A and Berr C: Plasma carotenoid levels and cognitive performance in an elderly population: Results of the EVA Study. *J Gerontol Ser A*, 2007; 62: 308-316.

How to cite this article:

Mehta N, Patani P and Singhvi I: A review on tomato lycopene. *Int J Pharm Sci & Res* 2018; 9(3): 916-23. doi: 10.13040/IJPSR.0975-8232.9(3).916-23.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)