



Received on 21 July, 2017; received in revised form, 07 October, 2017; accepted, 12 October, 2017; published 01 May, 2018

## SCREENING OF BIOACTIVE COMPOUNDS FROM LEAF OF *CENCHRUS CILIARIS* L. FROM THAR REGION OF RAJASTHAN, INDIA

S. Arora\* and G. Kumar

Department of Botany, Jai Narain Vyas University, Jodhpur - 342001, Rajasthan, India.

### Keywords:

*Cenchrus ciliaris* L.,  
Gas Chromatography - Mass  
Spectrometry, Retention Time (RT),  
Methyl commate B, Vitamin E,  
Hexadecanoic acid 2-hydroxy-1-  
(hydroxymethyl) ethyl ester

### Correspondence to Author:

**Dr. Sunita Arora**

Professor,  
Department of Botany,  
Jai Narain Vyas University,  
Jodhpur - 342001, Rajasthan, India.


**E-mail:** jnvusunitorora@gmail.com

**ABSTRACT:** *Cenchrus ciliaris* L. (Poaceae) commonly known as “Dhaman grass”, is a common famine food used during drought and scarcity of food in Rajasthan. The present investigation was carried out to determine the phytochemical composition of leaf of *C. ciliaris* using gas chromatography - mass spectrometry technique. The mass spectrum of the compounds found in the extract was matched with the national institute of standards and technology (NIST) and Willey 8 library. This study could segregate several natural bioactive compounds from the leaf using solvents of different polarity *i.e.* methanol, ethyl acetate and hexane. Sixty one biologically active compounds were extracted amongst them Tetracontane is present in maximum amount (30.09%) with RT = 25.302 min in the hexane leaf extract. Hence the present study helps to predict the formula, structure and activity of phytoconstituents from the leaf of *C. ciliaris* that can be used for drug design and justifying the use of this plant to treat many ailments as herbal medicine.

**INTRODUCTION:** Medicinal plants having high therapeutic value are becoming popular in the area of medicine for their less expense and less side effects as compared to modern allopathic drugs. The traditional medicines in the last few decades emerged to have immense acknowledgements in terms of their potential therapeutic value and it is estimated that 80% of community depends upon traditional medicine for their primary healthcare<sup>1</sup>. India is one of the 12 mega biodiversity countries of the World, which represents 11% of World's flora in about 2.4% of global land mass. Approximately 28% of the total Indian flora and 33% of angiosperms occurring in India are endemic. Higher human population density in India puts undue pressure on medicinal plants<sup>2</sup>.

Natural compounds extracted from plants have been suggested as alternative sources for antibiotics. Natural products are widely viewed as templates for structure optimization to make perfectly effective drugs. Natural products have historically been incredible as a source of therapeutic agents. Natural products have the characteristics of high chemical diversity, biochemical specificity and other molecular properties that make them favourable as lead structures for drug discovery<sup>3</sup>.

Extraction is the main step for the recovery and isolation of bioactive phytochemicals from plant materials. Analysis and extraction of plant matrices are important processes for the development, modernization and quality control of herbal formulations<sup>4</sup>. Hence for the discovery of lead compounds to be used as therapeutic drugs, the active principal in medicinal plants needs to be identified<sup>5</sup>. GC-MS method can serve as an interesting tool for testing the amount of some active principle of herbs.

<b>QUICK RESPONSE CODE</b>	<b>DOI:</b> 10.13040/IJPSR.0975-8232.9(5).1878-85
	<b>Article can be accessed online on:</b> <a href="http://www.ijpsr.com">www.ijpsr.com</a>
<b>DOI link:</b> <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.9(5).1878-85">http://dx.doi.org/10.13040/IJPSR.0975-8232.9(5).1878-85</a>	

It combines two analytical techniques to a single method of analyzing mixtures of chemical compounds. Gas chromatography separates the components of the mixture and mass spectroscopy analyzes each of the components separately<sup>6</sup>. *Cenchrus ciliaris*, known as Dhaman grass is extremely variable, tufted (sometimes shortly rhizomatous) with branching culms and perennial with types ranging in habit from ascendant to erect.

Leaf blades are linear, long, green, scabrous and hairy at the base. Panicle erect, grey, hairy with false spike. Each bur-like fascicle comprises of a single spikelet or cluster of 2 - 4 spikelets that in turn remain surrounded by involucre of bristles. Bristles are hairy, making the fascicle an adhesive quality<sup>7</sup>.

The desert ecosystem covers about 2% of the total landmass (Spreading over states of Rajasthan, Gujarat, Punjab and Haryana) in India, and is characterized by low precipitation and largely barren arid lands with only sparse or seasonal vegetal cover<sup>8</sup>.

This grass is gaining attention in various fields of research as it is more efficient C<sub>4</sub> plant at gathering CO<sub>2</sub> and utilizing nitrogen from the atmosphere and recycling of N in the soil. It is more competitive under the conditions of high temperature, solar radiation and low moisture<sup>9</sup>.

## MATERIAL AND METHODS:

**Collection of Plant Material:** Leaves of *C. ciliaris* were collected from Jodhpur district (Rajasthan) India, during August-October, 2016. Further identification and authentication of the specimens was done from Botanical Survey of India, Jodhpur (Rajasthan). The leaves were thoroughly washed with tap water followed by distilled water, shade dried for 20 days and grounded to fine powder. After sieving they were transferred to airtight polyethylene zipper bags for further use.

**Preparation of Plant Extracts:** The powdered plant leaves (5 g) were successively extracted with methanol, ethyl acetate and hexane. The extraction was done by hot continuous Soxhlet extraction<sup>10</sup>. The extracts were stored at - 4 °C till further use.

**GC-MS Analysis:** Gas chromatography - mass spectrometry (GC-MS) analysis of the extracts was

performed using a GC-MS (Model; QP 2010 ultra series, Shimadzu, Tokyo, Japan) equipped with thermal desorption system TD 20. Injection mode was split with a ratio of 50.0 and flow control mode was linear.

Velocity was maintained at 40.5 cm/sec and pressure was maintained at 81.9 kPa. For GC-MS detection Helium gas (99.99%) was used as a carrier gas at a constant flow rate. Total flow was 64.7 mL/min and column flow was 1.21 mL/min. Injector and mass transfer line temperature were set at 200 °C and 240 °C respectively.


The oven temperature was programmed at 80.0 °C and injection temperature maintained at 260.0 °C. Total running time of GC-MS was 46.28 minutes. The relative % amount of each component was calculated by comparing its average peak area to the total area, software adopted to handle mass spectra and chromatograms was Turbo mass. The relative percentage of the each extract constituents was expressed as percentage with peak area.

**Identification of Components:** Interpretation on mass spectrum of GC-MS was done using the database from National Institute of Standard and Technology (NIST), USA and WILEY8 library. Library contains more than 62,000 patterns. The mass spectrum of the unknown component was compared with the spectrum of the known components stored in this library. Molecular formula, molecular structure and weight of the compounds of the test samples were ascertained.

**RESULTS:** The analysis and extraction of plant material play an important role in the development, modernization and quality control of herbal formulations.

Total 61 bio-active constituents were identified including both major and minor constituents shown in **Table 1**. GC-MS chromatogram of the methanol, ethyl acetate and hexane extracts of leaf of *C. ciliaris* revealed 66, 78 and 58 peaks are shown in **Fig. 1** indicating the presence of 63, 66 and 47 bioactive compounds respectively.

Molecular structure of important and common compounds are presented in **Fig. 2**.



**FIG. 1: GC-MS CHROMATOGRAM OF *CENCHRUS CILIARIS* LEAF IN (A) METHANOL (B) ETHYL ACETATE AND (C) HEXANE**

Pentadecanoic acid is present in maximum amount (19.78%) followed by 9, 12, 15-Octadecatrienoic acid, (z,z,z)- (10.67%), Stigmasta-5, 22-Dien-3-OL (8.75%), 2- Hexadecen- 1- OL, 3, 7, 11, 15-Tetramethyl-, [R-[R\* R\*, (E)]] (4.08%), Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester (3.96%), Stigmast- 5- en- 3-ol, (3.beta.)- (3.49%), Ergost- 5- en- 3- ol, (3.beta., 24r)- (2.99%) and Vitamin E (2.82%) in methanolic leaf extract of *C. ciliaris*. Tetacontane is present in maximum amount (17.91%) followed by Hexadecanoic acid (9.15%), Stigmasta-5, 22-Dien-3-OL (6.83%),

Vitamin E (5.96%), Octadecanal (5.75%), 9, 12, 15-Octadecatrienoic acid, (z,z,z)- (5.15%) and Ergost-5-en-3-ol, (3.beta.) (3.11%) in ethyl acetate leaf extract of *C. ciliaris*. Tetracontane is present in maximum amount (30.09%), followed by Penta-decanoic acid (11.36%) Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester (10.37%), Stigmasta-5, 22-Dien-3-OL (7.83%), Octadecanoic acid, 2,3-dihydroxypropyl ester (4.13%), Vitamin E (3.75%), 9, 12, 15-Octadecatrienoic acid, (z,z,z)- (3.25%) and Ergost- 5- en- 3- ol, (3.beta., 24r)- (2.74%) in hexane leaf extract.


**TABLE 1: BIOACTIVITY OF PHYTOCOMPOUNDS IDENTIFIED IN *CENCHRUS CILIARIS* LEAF**

S. no.	Compound	Mol. Weight	Mol. formula	Solvents			Bioactivities
				Met	Et. Act	Hex	
1	2,3-Dihydro-3,5-Dihydroxy-6-Methyl-4H-Pyran	144	C <sub>6</sub> H <sub>8</sub> O <sub>4</sub>	+	-	-	Antimicrobial, AntiInflammatory
2	2-Methoxy-4-Vinylphenol	150	C <sub>9</sub> H <sub>10</sub> O <sub>2</sub>	+	-	-	Antibacterial
3	1-Tetradecene	196	C <sub>14</sub> H <sub>28</sub>	+	-	-	Antituberculosis
4	Guanosine	283	C <sub>10</sub> H <sub>13</sub> N <sub>5</sub> O <sub>5</sub>	+	-	-	Cytotoxic against T cells lines, Anti viral against Vero cells infected with HSV-1
5	Heptadecane	240	C <sub>17</sub> H <sub>36</sub>	+	-	-	Antioxidant
6	Dodecanoic acid	200	C <sub>12</sub> H <sub>24</sub> O <sub>2</sub>	+	-	-	Antimicrobial
7	Megastigmatrienone	190	C <sub>13</sub> H <sub>18</sub> O	+	-	-	Aroma
8	Nonadecane	268	C <sub>19</sub> H <sub>40</sub>	-	+	-	Antimicrobial, Cytotoxic


9	n- Tridecan-1-ol	200	C <sub>13</sub> H <sub>28</sub> O	+	+	-	Natural mosquito control agent
10	5-Octadecene, (E)-	252	C <sub>18</sub> H <sub>36</sub>	+	-	-	Stronger sexual characters
11	2-Methyltetracosane	352	C <sub>25</sub> H <sub>52</sub>	+	+	+	Free radical scavenging
12	2-Methylhexacosane	380	C <sub>27</sub> H <sub>56</sub>	-	+	+	Antimicrobial, Hypocholesterolemic
13	7,9-Di-tert-butyl-1-oxaspiro (4,5)deca-6,9-diene-2,8-dione	276	C <sub>17</sub> H <sub>24</sub> O <sub>3</sub>	-	-	+	Antimicrobial
14	2,6,10-Trimethyl, 14-Ethylene-14- Pentadecne	296	C <sub>20</sub> H <sub>38</sub>	-	+	+	Antiproliferative
15	2-Pentadecanone, 6,10,14- Trimethyl-	268	C <sub>18</sub> H <sub>36</sub> O	+	+	+	Allelopathic, Antibacterial
16	2-Hexadecen-1-OL, 3,7,11,15- Tetramethyl-, [r-[r* r*, (e)]]	296	C <sub>20</sub> H <sub>40</sub> O	+	+	+	Antimicrobial, Sedatives and Anesthetics
17	1,2-Benzenedicarboxylic Acid	390	C <sub>8</sub> H <sub>6</sub> O <sub>4</sub>	-	-	+	Antioxidant, Antimicrobial, Antifouling
18	1,2-Benzene dicarboxylic acid, Diethyl ester	222	C <sub>12</sub> H <sub>14</sub> O <sub>4</sub>	-	+	-	Cosmetics, Insecticides, Plasticizer
19	1,2-Benzenedicarboxylic Acid, bis(2-Methylpropyl) Ester	278	C <sub>16</sub> H <sub>22</sub> O <sub>4</sub>	-	+	+	Antimicrobial, Antifouling
20	Hexadecanoic acid	256	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	-	+	-	Antitumor
21	Hexadecanoic acid, methyl ester	270	C <sub>17</sub> H <sub>34</sub> O	+	-	-	Antioxidant, Insecticide,hemolytic, Hypo - cholesterolemic
22	Hexadecanoic acid , ethyl ester	284	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	+	+	+	Antioxidant, Hypocholesterolemic Antiandrogenic, Hemolytic, 5-Alpha reductase inhibitor
23	13-Docosanoic acid	338	C <sub>22</sub> H <sub>42</sub> O <sub>2</sub>	+	-	-	Surfactant, Precursor to Biodiesel fuel
24	cis-13-Octadecenoic acid	282	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	+	-	-	Therapeutic uses in medicine, Surgery
25	Pentadecanoic acid	242	C <sub>15</sub> H <sub>30</sub> O <sub>2</sub> C <sub>14</sub> H <sub>28</sub> O <sub>2</sub>	+	-	+	Lubricants, Adhesive agents Antioxidant,
26	Tetradecanoic acid	228		+	+	+	Anticancer, Hypocholesterolemic
27	Tetratetracontane	618	C <sub>44</sub> H <sub>90</sub>	-	+	-	Hypoglycaemic, Antioxidant
28	Tetracontane	562	C <sub>40</sub> H <sub>82</sub>	-	+	+	Anti Inflammatory
29	Heptadecanoic acid	270	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	+	-	+	Antimicrobial
30	9,12-Octadecadienoic acid (z,z,-)	280	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	+	+	+	AntiInflammatory, Antibacterial, Antiarthritic, Hepatoprotectiv, Anti-histaminic, Anticoronary
31	9,12-Octadeca dienoic acid (z,z,-), methyl ester	294	C <sub>19</sub> H <sub>34</sub> O <sub>2</sub>	+	-	-	Hepatoprotective, Anti-histaminic, Antieczemic, Hypocholesterolemic
32	9,12,15-Octadecatrienoic acid, (z,z,z)-	278	C <sub>18</sub> H <sub>30</sub> O <sub>2</sub>	+	+	+	Preventive against cardiovascular diseases
33	9,12,15-Octadecatrienoic acid, methyl ester, (z,z,z)-	292	C <sub>19</sub> H <sub>32</sub> O <sub>2</sub>	+	-	-	Anti-inflammatory, Hypocholesterolemic, Cancer preventive, Hepatoprotective
34	Octadecanoic acid	284	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	+	+	+	Antifungal, Antitumor, Antibacterial
35	(E)-9-Octadecenoic acid, ethyl ester	310	C <sub>20</sub> H <sub>38</sub> O <sub>2</sub>	+	-	-	Flavoring ingredient
36	Phytol, acetate	338	C <sub>22</sub> H <sub>42</sub> O <sub>2</sub>	+	+	+	Flavor and fragrances
37	Squalene	410	C <sub>30</sub> H <sub>50</sub>	+	+	+	Antioxidant, Anticancer Pesticide, Sunscreen, Perfumery, Chemo preventive
38	.gamma.-Tocopherol Vitamin E	416	C <sub>28</sub> H <sub>48</sub> O <sub>2</sub> C <sub>29</sub> H <sub>50</sub> O <sub>2</sub>	+	-	+	Antioxidant, Cardio protective, Anticancer, Anti-inflammatory Antiaging, Analgesic, Antidiabetic, Antidermatitic, Antileukemia, Anticancer, Vasodilator,
39		430		+	+	+	Hepatoprotective, Hypocholesterolemic, Antibronchitic, Anticoronary
40	Ergost-5-en-3-ol, (3.beta.,24r)-	400	C <sub>28</sub> H <sub>48</sub> O	+	-	+	Liver disease, Jaundice, Arthrosclerosis
41	Ergost-5-en-3-ol, (3.beta.)-	400	C <sub>28</sub> H <sub>48</sub> O	+	+	-	Antioxidant, Hypocholesterolemic
42	Stigmasta-5,22-dien-3-ol	412	C <sub>29</sub> H <sub>48</sub> O	+	+	+	Synthetic Progesterone
43	Stigmast-5-en-3-ol, (3.beta.)-	414	C <sub>29</sub> H <sub>50</sub> O	+	+	+	Anti Inflammatory, Antipyretic, Anti ulcer, Antiarthritic
44	Methyl Commate B	470	C <sub>31</sub> H <sub>50</sub> O <sub>3</sub>	+	+	+	Antimicrobial, anti-inflammatory
45	Naphthalene	128	C <sub>10</sub> H <sub>28</sub>	-	+	-	Antiseptic, Carcinogenic
46	Tetradecane	198	C <sub>14</sub> H <sub>30</sub>	-	+	+	Antifungal, Antibacterial, Nematicidal
47	Dibutyl phthalate	278	C <sub>16</sub> H <sub>22</sub> O <sub>4</sub>	+	-	-	Antifungal, Antimicrobial, Anti malarial, Plasticizer, Ectoparasiticide
48	Octadecanal	268	C <sub>18</sub> H <sub>38</sub> O	+	-	-	Sex Pheromone

49	Eucalyptol	154	$C_{10}H_{18}O$	-	-	+	Antioxidant, antibacterial, Anti-inflammatory, antileukaemia and antirrhinosinusitis
50	Dodecane	170	$C_{12}H_{26}$	-	-	+	Solvent, distillation chaser and scintillator component
51	Pentadecane	212	$C_{15}H_{32}$	-	+	+	Suger-phosphatase inhibitor, Chymosin inhibitor, Antibacterial
52	Pentadecane, 2,6,10,14-tetramethyl	268	$C_{19}H_{40}$	-	+	+	Pathogenesis of Rheumatoid arthritis and lupus
53	Hexadecane	226	$C_{16}H_{34}$	-	+	+	Antifungal, Antibacterial, Antioxidant
54	Heptadecane	240	$C_{17}H_{36}$	-	+	+	Antioxidant
55	Tetracosane	338	$C_{24}H_{50}$	-	+	-	Cytotoxicity against colon, breast and gastric cancer cells
56	Pentacosane	352	$C_{25}H_{52}$	-	+	-	Antibacterial activity
57	Eicosane	282	$C_{20}H_{42}$	-	+	+	Antifungal, Antitumor, Antibacterial, Larvicidal, Cytotoxic, Antimicrobial
58	Hexadecane, 2,6,10,14-tetramethyl	282	$C_{20}H_{42}$	-	+	+	Bio markers in Petroleum studies
59	tridecane	184	$C_{13}H_{28}$	-	+	-	Fragrance agent
60	Phenol, (1,1-dimethylethyl)-4-methoxy-	180	$C_{11}H_{16}O_2$	+	-	-	Antioxidant, Antitumor, Antimutagen, Radical scavenging
61	Hexadecanoic acid 2-hydroxy-1-(hydroxymethyl)ethyl ester	330	$C_{19}H_{38}O_4$	+	-	+	Antioxidant


Met: Methanol, Et. act: Ethyl acetate, Hex: Hexane, +: present, -: absent




A




B




C




D




E




F




G



H



I



J






FIG. 2: MOLECULAR STRUCTURE OF (A) 2-METHYLTETRACOSANE (B) 2-PENTADECANONE, 6, 10, 14-TETRAMETHYL (C) 2-HEXADECEN-1-OL, 3, 7, 11, 15- TETRAMETHYL-, [R- [R\*, R\*-(E)]]- (D) HEXADECANOIC ACID, ETHYL ESTER (E) TETRADECANOIC ACID (F) 9, 12- OCTADECADIENOIC ACID (Z, Z)- (G) 9, 12, 15-OCTADECATRIENOIC ACID (Z, Z, Z) - (H) OCTADECANOIC ACID (I) PHYTOL, ACETATE (J) SQUALENE (K) VITAMIN E (L) STIGMASTA-5,22-DIEN-3-OL (M) STIGMAST-5-EN-3-OL, (3.BETA.)- (N) METHYL COMMATE B AND (O) ERGOST-5-EN-3-BETA-OL

**DISCUSSION:** Phenolic compounds show antioxidant, anti-mutagenic and scavenging activity on free radicals and prevent cancer and cardiovascular heart diseases<sup>11</sup>. 2-Hexadecen-1-ol, 3, 7, 11, 15- tetramethyl-, [R- [R\*, R\*-(E)]]- and Phytol, acetate both are phenolic compounds in nature. 2-Methyltetraacosane, also known as Isopentacosane belongs to the family of acyclic alkanes, shows a free radical scavenging activity. 9, 12-Octadecadienoic acid (Z, Z)- Linoleic acid shows anti-inflammatory, hypocholesterolemic, cancer preventive, hepatoprotective, nematocidal, insectifuge, antihistaminic, antieczemic, antiacne, 5- Alpha reductase inhibitor, antiandrogenic, antiarthritic, anticoronary and insectifuge properties<sup>12</sup>. 9, 12, 15-Octadecatrienoic acid (z, z, z)- an  $\alpha$ -linolenic acid (ALA) is one of the two essential fatty acids in humans.

ALA, a nonproprietary, naturally occurring omega-3 fatty acid found in foodstuffs shows anti-inflammatory property. ALA reported as a potent nutraceutical that protects the brain from stroke, characterized by its pleiotropic effects in

neuroprotection, vasodilation of brain arteries and neuroplasticity<sup>13</sup>. Octadecanoic acid and stearic acid exhibit antifungal, antitumor and antibacterial properties<sup>14</sup>. Tetradecanoic acid, a saturated fatty acid occurring in most animal and vegetable fats, particularly butterfat, coconut palm and nutmeg oil, is used to synthesize flavour and as an ingredient in soaps and cosmetics. Tetradecanoic acid has an antioxidant, lubricant, hyper-cholesterolemic and cancer-preventive activity<sup>15</sup>. Monohydroxy commic acid B, commonly known as Methyl commate B is found in the resin of *Commiphora glandulosa* Schinz<sup>16</sup>. Methyl commate B is a triterpene glycoside in nature. Triterpene glycosides are natural highly polar compounds with low volatility and were discovered in higher plants. They are well-known for their cytotoxic, antimicrobial, anticoagulant, hemolytic, antiviral, antiparasitic and antitumor properties<sup>17</sup>. 2-Pentadecanone, 6, 10, 14- trimethyl is a diterpene in nature. Diterpenes show potent and selective activity towards platelet-activating factor increasing in conditions of shock, burns, ulceration and inflammation.

As per clinical trial report, its importance in bladder hyperreflexia and diabetic neuropathy and as an anticancerous agent against ovarian, breast and lung cancer as well proved<sup>18</sup>. Squalene is a triterpene that is an intermediate in the cholesterol biosynthesis pathway. It was so named because of its occurrence in shark liver oil; it is the main component of skin surface polyunsaturated lipids and shows some advantages for the skin as an emollient and antioxidant for hydration and antitumor activities. Squalene considered as an interesting natural molecule, with broad applications in food industry, cosmetics and in prevention and treatment of human diseases<sup>19</sup>.

Vitamin E is a group of eight different compounds, but only two of the forms,  $\alpha$ -tocopherol and  $\gamma$ -tocopherol, are commonly found in the human body. Vitamin E is the most potent lipid-soluble antioxidant<sup>20</sup>. Phytosterols produce health benefits in animals/humans such as reduction of cholesterol level with decreased risk of coronary heart diseases, anti-inflammatory activities, induction of apoptosis in cancer cells and in prevention and treatment of diseases<sup>21</sup>.

Ergost-5-en-3 beta-ol, also called "campesterol" competes with cholesterol and reduces the absorption of its in human intestine<sup>22</sup>. Stigmasterol, naturally occurring steroid alcohol shows antiarthritic, anti-inflammatory<sup>23</sup> actions as well as lipid lowering and antiatherogenic effects<sup>24</sup>.

Stigmasterol inhibits OVA induced asthma in guinea pigs<sup>25</sup>. Phytosterols such as  $\beta$ -sitosterol structurally resembles to cholesterol and able to inhibit the absorption of cholesterol, cancer-cell growth, angiogenesis, invasion and metastasis.  $\beta$ -sitosterol is well known natural sterol in composition of known herbal drugs used for the treatment of benign prostatic hyperplasia and prostate cancer. Besides the compound elevated enzymatic and non-enzymatic antioxidant in cells making it effective anti-diabetic, neuroprotective and chemoprotective agent as well<sup>26</sup>.

**CONCLUSION:** This study was aimed at identifying the best solvent system to obtain maximum number of phytochemicals with strong bioactivities from this  $C_4$  plant of stressed conditions. GC-MS itself is a best tool and it has

been found out that ethyl-acetate is a best solvent for extraction of compounds from *C.iliaris*. We observed that *C.iliaris* is a potent grass yielding compounds of high therapeutic value that after clinical and other trials can be used by drug-developing companies. A further study involving purification of these phytochemicals individually and determining their mechanistic actions will be the future scope of this study which will help their safer application in therapeutics.

**ACKNOWLEDGEMENT:** We are grateful to the Department of Botany, Jai Narain Vyas, University, Jodhpur and AIRF (Advanced Instrumentation Research Facility), Jawaharlal Nehru University, New Delhi, for assisting laboratory and instrumentation facilities.

**CONFLICT OF INTEREST:** Nil

## REFERENCES:

1. Kalaisezhiyen P and Sasikumar V: GC-MS evaluation of Chemical constituents from methanolic leaf extract of *Kedrostis foetidissima* (Jacq.) Cogn. Asian Pharm Clin Res 2012; 5(4): 77-81.
2. Chitale VS, Behera MD and Roy PS: Future of endemic flora of biodiversity hotspot in India. PLoS One 2014; 9(12): 1-15.
3. Siddiqui AA, Iram F, Siddiqui S and Sahu K: Role of natural products in drug Discovery process. Int J Drug Dev Res 2014; 6(2): 172-204.
4. Ghasemzadeh A, Jaafar HZE and Rahmat A: Optimization protocol for the extraction of 6-gingerol and 6-shogaol from *Zingiber officinale* var. rubrum theilade and improving antioxidant and anticancer activity using response surface methodology. BMC Complement Altern Med 2015; 15: 258.
5. Vuorela P, Leinonen M, Saikku P, Tammela P, Rouha JP, Wennberg T and Vuorela H: Natural products in the process of finding new drug candidates. Curr Med Chem 2004; 11(11): 1375-1389.
6. Gupta AK and Bhat JL: GC-MS analysis of methanol extract of *Acacia nilotica* (L.) leaves. Int J Pharm Chem 2016; 6(02): 50-53.
7. Singariya P, Mourya KK and Kumar P: Gas Chromatography-Mass Spectrometric analyses of acetone extract of *Cenchrusiliaris* (Dhama grass). Ind J Sci Nat 2015; 6(4): 652-661.
8. Arisdason W and Lakshminarasimhan P: Status of plant diversity in India: an overview. ENVIS Centre on floral diversity, BSI, Kolkata, India 2016.
9. Agrawal P: Ecophysiological and biochemical studies related to drought adaptation in grasses of Indian desert. Ph.D. Thesis, J. N. V University 2007.
10. Harborne JB: Phytochemical Methods. Chapman and Hall, London, Edition 2, 1984; 5-6.
11. Ozcan T, Akpınar-Bayazit A, Yılmaz - Ersan L and Delikanlı B: Phenolics in human health. Int J Chem Engin Appl 2014; 5(5): 393-396.

12. Rajeswari G, Murugan M and Mohan VR: GC-MS analysis of bioactive components of *Hugonia mystax* L. (Linaceae). Res J Pharm, Bio Chem Sci 2012; 3(4): 301-308.
13. Blondeau N, Lipsky RH, Bourourou M, Duncan MW, Gorelick PB, and Marini AM: Alpha-linolenic acid: an omega-3 fatty acid with neuroprotective properties-ready for use in the stroke clinic? BioMed Res Int 2015; 2015: 519830.
14. Vijisarl ED and Subramanian A: GC – MS analysis of ethanol extract of *Cyperus rotundus* Leaves. Int J Curr Biotechnol 2014; 2(1): 19-23.
15. Anburaj G, Marimuthu M, Rajasudha V and Manikandan R: Phytochemical screening and GC-MS analysis of ethanolic extract of *Tecoma stans* (family: Bignoniaceae) yellow bell flowers. J Pharmacog Phytochem 2016; 5(4): 172-175.
16. Thomas AF and Willhalm B: triterpene of Commiphora; mass spectra and organic analysis, mass spectroscopic studies and the structure of commic acid A and B. Tetrahedron letters 1964; 5(43): 3177-3183.
17. Bahrami Y and Franco CMM: Acetylated triterpene glycosides and their biological activity from holothuroidea reported in the past six decades. Mar Drugs 2016; 14(8): 147.
18. Lanzotti V: Natural products. Springer-Verlag, Heidelberg, Berlin 2013; 3173-3191.
19. Popa O, Babeanu NE, Popa I, Nița S, and Dinu-Parvu CE: Methods for obtaining and determination of squalene from natural sources. Bio Med Res Int 2015; 2015: 367202.
20. Gregory III JF: Accounting for differences in the bioactivity and bioavailability of vitamers. Food Nutr Res 2012; 56(1):5809.
21. Ogbe RJ, Ochalefu DO, Mafulul SG and Olaniru OB: A review on dietary phytosterols: Their occurrence, metabolism and health benefits. Asian J Plant Sci Res 2015; 5(4): 10-21.
22. Choudhary SP and Tran LS: Phytosterols: perspectives in human nutrition and clinical therapy. Curr Med Chem 2011; 18(29): 4557–4567.
23. Chen WP, Yu C, Hu PF, Bao JP, Tang JL and Wu LD: Stigmasterol blocks cartilage degradation in rabbit model of osteoarthritis. Acta Biochimica Polonica. 2012; 59(4): 537-41.
24. Othman RA and Moghadasian MH: Beyond cholesterol-lowering effects of plant sterols: clinical and experimental evidence of anti-inflammatory properties. Nut Rev 2011; 69(7): 371-382.
25. Antwi AO, Obiri DD and Osafo N: Stigmasterol modulates allergic airway inflammation in guinea pig model of ovalbumin-induced asthma. Mediat Inflamm 2017; 2017: 2953930.
26. Saeidnia S, Manayi A, Gohari AR and Abdollahi M: The story of beta- sitosterol- a review. EJMP2014;4(5):590-60.

**How to cite this article:**

Arora S and Kumar G: Screening of bioactive compounds from leaf of *Cenchrus ciliaris* L. from Thar region of Rajasthan, India. Int J Pharm Sci Res 2018; 9(5): 1878-85. doi: 10.13040/IJPSR.0975-8232. 9(5).1878-85.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)