



Received on 18 June, 2017; received in revised form, 28 August, 2017; accepted, 17 December, 2017; published 01 May, 2018

ANTICANCER POTENTIAL OF LEAFLESS MISTLETOE (*VISCUM ANGULATUM*) FROM WESTERN GHATS OF INDIA

Saurabh Sharma¹, Radhey Shyam Sharma¹, Milind Madhav Sardesai² and Vandana Mishra^{*1}

Bioresources and Environmental Biotechnology Laboratory¹, Department of Environmental Studies, University of Delhi, Delhi - 110007, Delhi, India.

Department of Botany², Savitribai Phule Pune University, Pune - 411007, Maharashtra, India.

Keywords:

Mistletoe, Anticancer activity, Bioprospecting, Parasitic plant, Antioxidant potential, Phenolics and flavanoids

Correspondence to Author:

Dr. Vandana Mishra

Assistant Professor (Stage III),
Bioresources and Environmental
Biotechnology Laboratory,
Department of Environmental
Studies, University of Delhi,
Delhi - 110007, Delhi, India.


E-mail: mistletoe_h@hotmail.com

ABSTRACT: Phenolics have been shown to attenuate the initiation and progression of cancer. Owing to the parasitic nature, mistletoes are a rich source of phenolics and flavonoids. European mistletoe (*Viscum album*) formulations provide the first line of treatment for cancer by stimulating the immune system and improve the quality of life. In fact, host-associated variation has been shown to affect the therapeutic potential of the *V. album* preparations. Leafless Mistletoe (*Viscum angulatum*), a hemiparasitic plant of the Western Ghats widely used in folk medicine, contains diverse phytochemicals having anti-inflammatory properties. However antioxidant and anticancer potential of *Viscum angulatum* is still not investigated. In the present study, we analyzed the contents of total phenolics (TPC), flavonoids (TFC), antioxidant potential and cytotoxic potential of extracts of leafless mistletoe plants parasitic on two different host trees, *Olea dioica* and *Flacourtia indica*. The whole plant of leafless Mistletoe was extracted in organic solvents (methanol and ethanol) and water (aqueous) using rotary evaporator. In the methanolic extract, TPC of plants parasitic on *O. dioica* was higher than that of *F. indica*, however, in ethanolic and aqueous extracts, TPC of plants parasitic on *O. dioica* was lower than that of *F. indica*. Interestingly, the pattern of antioxidant activity truly corresponds with that of TPC. In both methanolic and ethanolic extracts, the TFC of the plants parasitic on *O. dioica* was lower than that of *F. indica*. The plants parasitic on *O. dioica*, showing high antioxidant activity also showed high cytotoxicity (LC₅₀ - 79.33 µg/ml) on MDA-MB-231 breast cancer cells. The study provides experimental evidence on the anticancer potential of the *V. angulatum* plant. It also suggests the involvement of host tree in determining the therapeutic potential of *V. angulatum* which needs to be established further.

INTRODUCTION: Plants used in traditional folk medicine have been the source of the novel therapeutic compounds^{1, 2}. Most of the traditionally used medicinal plants are an excellent source of natural antioxidants having the potential for prevention and treatment of several inflammatory and degenerative diseases³.

Phenolics have been identified as the key components responsible for the antioxidant potential of these plants. The antioxidants can reduce the oxidative stress in cells and impart protection against oxidative damage from the free radicals through various mechanisms^{4, 5}. In fact, approximately 50 % of the anticancer drugs that got approved in recent times (between 1981 - 2010) were either natural compounds or their semi-synthetic or synthetic derivatives⁶.

Recently, there has been a growing interest in herbal medicines due to their role in disease prevention. European mistletoe (*Viscum album*), a hemiparasitic plant that grows on diverse host trees,

QUICK RESPONSE CODE 	DOI: 10.13040/IJPSR.0975-8232.9(5).1902-07
	Article can be accessed online on: www.ijpsr.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.9(5).1902-07	

has shown significant anticancer effects^{7, 8}. In European countries, mistletoe is extensively used in cancer therapeutics⁹. Mistletoe preparations made of plants parasitizing different host trees have been shown to vary in therapeutic potential. Mistletoe is also used in the form of a herbal tea known for its potential for immuno-stimulation and cancer prevention¹⁰. Biologically active components of mistletoe extract such as mistletoe lectins (ML) and secondary metabolites have been identified as active principle responsible for anti-inflammatory and anti-cancerous effect^{11, 12}.

Although there are many species of *Viscum* worldwide only European mistletoe, i.e., *V. album* has been studied for its anticancer potential. The Western Ghats in India, a hotspot of biodiversity is a home to 16 - 18 species of *Viscum* which may serve as an alternative source of phytomedicines for cancer therapeutics⁸. Various studies have demonstrated that phytochemical profile of mistletoe might vary with the host plant as the hemiparasite depends on a host tree for its water and mineral requirement^{13, 14}.

Therefore, the present work aimed to determine the pharmaceutical potential of a leafless mistletoe (*V. angulatum*) inhabiting Western Ghats region of India. The total phenolics, flavonoids, and antioxidant potential were analyzed in the different extracts of mistletoe parasitic on two different host trees, i.e., *Olea dioica* and *Flacourtia indica*. The present study establishes the cytotoxic potential of *V. angulatum* against breast cancer and also suggests the role of host variation in determining the therapeutic potential of leafless mistletoe.

MATERIALS AND METHODS:

Materials: Folin-Ciocalteu reagent (FCR), 2,2-diphenyl-1-picrylhydrazyl (DPPH), L-ascorbic acid, potassium acetate (CH₃COOK), aluminium chloride (AlCl₃) were purchased from Merck (Germany). Gallic acids, sodium carbonate (Na₂CO₃) were purchased from Sigma Chemicals (Sigma-Aldrich, USA). Cell proliferation reagents were obtained from Invitrogen (USA) and HyClone (USA). All other chemicals used in experiments were of analytical grade.

Preparation of Crude Extract: *V. angulatum* parasitic on *O. dioica* and *F. indica* trees were collected from Western Ghats region of India and

stored at -20 °C till further use. The plant samples (50 g) were crushed in the presence of liquid N₂ and homogenized in 200 ml of the respective extraction solvent, i.e., ethanol, methanol and aqueous. Homogenized samples were incubated overnight in shaking condition at 4 °C. The mixture was centrifuged at 5000 rpm for one hour, and the supernatant was filtered. The filtrate was concentrated using a rotary evaporator, and the extracted phytochemicals were further dissolved in respective solvents.

Determination of Total Phenolic Contents: Total phenolics of different crude extracts were determined using Folin-Ciocalteu reagent (Roy et al., 2010) with slight modifications. Briefly, 8 µl of each extract solution (10 mg/ml) were separately mixed with 200 µl of Folin-Ciocalteu reagent and incubated for 5 min at RT. After that 1 ml of 15 % sodium carbonate solution was added and the mixture was allowed to stand for 30 min. The absorbance was measured at OD₇₆₀ nm using a UV-visible spectrophotometer. The concentration of the total phenolics was expressed as gallic acid equivalents. Three replicates were kept for each sample and results are represented as mean ± standard deviation¹⁵.

Determination of Flavonoids: The flavonoids content was determined using the aluminium chloride colorimetric method. Briefly, 100 µl of the methanolic, ethanolic and aqueous extract (10 mg/ml) were separately mixed with 0.1 ml of 10 % aluminium chloride. Subsequently, the reaction mixtures were diluted with 0.1 ml of 1 M potassium acetate and 2.8 ml of distilled water. The absorbance of the reaction mixture was measured at 415 nm after 30 min incubation. Blank samples for each extract were prepared from 100 µl of samples and diluted with 3 ml of respective solvent. The quercetin dissolved in the respective solvent was used as a standard. The total flavonoid content of each extract was expressed in terms of a milligram of quercetin equivalent per gram of dry weight. All the experiments were performed in replicates of three, and results are presented as mean ± standard deviation¹⁶.

Quantitative DPPH Radical Scavenging Assay: The DPPH radical scavenging activity was determined using the method proposed by Brand - Williams et al., (1995).

The reaction was performed in 96 well plate. A volume of 30 μ l sample and 200 μ l of 4 mM DPPH solution were added to each well. The 30 μ l of respective extraction solvent and 200 μ l of DPPH were set as blank. The negative control was made up of 30 μ l sample and 200 μ l the respective solvent. The decrease in the absorbance of the resulting solution was examined at OD₅₁₇ nm after 30 minutes of incubation. Ascorbic acid (10 mg/mL) was used as the positive control. The percent scavenging effect of the different extracts against DPPH radicals was calculated using the following equation:

$$\text{DPPH scavenging effect (\%)} = [(A_o - A_s)/A_o] * 100$$

Where, A_o is absorbance of blank, and A_s is absorbance of the respective samples¹⁷.

Cell Cytotoxicity Assay: To demonstrate the anti-cancer potential of *V. angulatum*, the cytotoxic effect of the methanolic extract was assessed in MDA-MB-231 breast cancer cells using MTT assay in a 96-well plate. The cells were challenged with different concentration of methanolic extract of *V. angulatum*. Cell lines were maintained as a subconfluent monolayer in Dulbecco's modified Eagles's medium with 100 units/ml penicillin and 10% fetal bovine. Cells were incubated in a humidified chamber at 37 °C with 5 % CO₂. The culture was allowed to grow to confluence and the viability of the cells was checked using trypan blue (0.4 % trypan blue in PBS) exclusion method. The cells were incubated with MTT reagent (20 ml of 5 mg/ml per well) at 37 °C for 4 h. The formazan product was solubilized by the addition of 100 ml of 0.1 N HCl in isopropanol. Cell viability was estimated by recording the absorbance at 595 nm in an ELISA plate reader¹⁸.

Statistical Analysis: The results were statistically analyzed by independent sample t-test using SPSS ver. 16.0 for Windows (SPSS Inc., Chicago, Illinois, USA)¹⁹. The '*p*' value < 0.05 was considered to be a significant value. The results are presented as the mean \pm standard deviations.

RESULTS AND DISCUSSION: Mistletoe or *Viscum* sp. have been documented to treat a wide variety of diseases such as cancer, diabetes, skin diseases, high blood pressure, etc.^{20, 21, 22}. Various phyto-constituents of *V. album* and *V. articulatum*

such as viscotoxins, phenylpropanes, flavonoids, phenols, amines, polysaccharides, ribosome inactivating proteins and lectins have been identified as the active components^{23, 24, 25, 26}. Mechanism of anticancer potential of *V. album* and *V. articulatum* has also been studied^{8, 25}. *V. Album* based mistletoe preparations varying in host plants are available in the market and are in use for the treatment of different cancers²⁷. *V. angulatum* is a mistletoe sp. (**Fig. S1**) prevalent in Western Ghat region, one of the hot spots of biodiversity. A few studies have revealed phytochemical profiling²⁸ and diuretic activity of this plant²⁹.



FIG. 1: LEAFLESS MISTLETOE (*VISCUM ANGULATUM*) PARASITIC ON *OLEA DIOICA* SHOWING A CHARACTERISTIC DROOPING HABIT

In fact a unique antioxidant protein, Van Prx has also been purified from *V. angulatum*³⁰. Despite its use in folk medicine, the pharmaceutical potential of *V. angulatum* is still unexplored. Therefore, in the present study, extract of *V. angulatum* parasitic on two different host trees, *O. dioica* and *F. indica* trees were compared in terms of their phenolic content, flavonoid content, antioxidant potential, and cytotoxicity against cancer cells. Phenolic phytochemicals are most abundant secondary metabolites and widely distributed in the plant kingdom³¹. Plants phenolics and flavonoids are mostly known for their broad range of biological properties specifically the antioxidant activities³². In this study, the total phenolics (TPC) and flavonoids (TFC) of *V. angulatum* parasitizing on *O. dioica* and *F. indica* trees were extracted in methanol, ethanol and distilled water. The TPC and TFC of the methanol and ethanol extract of *V. angulatum* were significantly higher than that of aqueous extract. TPC of methanolic extract of the plants parasitic on *O. dioica* was higher than that of plant parasitizing *F. indica* (**Fig. 2A**).

On the contrary, the TPC of ethanolic extract of *V. angulatum* (41.80 µg/ml of dry extract) parasitic on *F. indica* tree was significantly higher ($p < 0.02$) than that of plant parasitic on *O. dioica*. Further, the TFC of different extracts of *V. angulatum* were

measured. Both methanolic and ethanolic extract of *V. angulatum* parasitic on *O. dioica* showed a decreased level of TFC as compared to plants parasitic on *F. indica* (Fig. 2B).




FIG. 2: TOTAL PHENOLIC CONTENT (TPC) AND FLAVONOID CONTENT (TFC): The phenolic and flavonoid content was estimated by Folin-Ciocalteu method and aluminium chloride colorimetric method. The ethanolic extract of *V. angulatum* parasitic on *F. indica* tree showed the significantly high ($p < 0.02$) TPC content (A), however methanolic extract of *V. angulatum* parasitic on *F. indica* showed significantly elevated ($p < 0.012$) TFC content (B). (Note: ‘*’ between the bars represent the significant difference ($p < 0.05$) between the means).

The antioxidant activity of *Viscum angulatum* plant extract was estimated based on the scavenging capacity of the free DPPH radical. The methanolic extract of *Viscum angulatum* possessed higher antioxidant activity (57.88 + 0.66 %) in comparison with the ethanolic and aqueous extract. In accordance with the TPC, the antioxidant activity of the methanolic extract was higher in leafless mistletoe parasitizing *Olea dioica*, however, the ethanolic and aqueous extract of plant parasitic on *F. indica* showed high antioxidant activity (Fig. 3).

activity of the methanolic extract of the leafless mistletoe suggest an important role of phenolics other than flavonoids.




FIG. 3: ANTIOXIDANT POTENTIAL OF *V. ANGULATUM*: Antioxidant potential of different Mistletoe extract was evaluated by DPPH assay. Methanolic extract of *V. angulatum* parasitic on *F. indica* showed strong antioxidant potential as compared with that growing on *O. dioica*

Including flavonoids, different phenolic compounds have been shown to contribute to the antioxidant activity of plants. However, the contrasting pattern observed in the flavonoid content, and antioxidant

It may be noted that three phenolic glycosides have been reported from *V. angulatum*²⁸. The effect of mistletoe extract was tested on breast carcinoma (MDA-MB-231 cells) using MTT assay. Since methanol extract of *V. angulatum* showed significantly higher antioxidant activity in comparison to the ethanolic and aqueous extract, the cytotoxicity of methanolic extract was evaluated. The methanolic extract of *V. angulatum* showed cytotoxicity within 24h of treatment indicating its high anticancer potential. In fact, the extract of *V. angulatum* parasitic on *O. dioica* which showed high antioxidant potential also showed high cytotoxicity than that of *V. angulatum* parasitic on *F. indica*. As evident in *in-vitro* studies, the methanol extract isolated from mistletoe hosted on *O. dioica* tree decreased the proliferation (LC_{50} - 79.33 µg/ml) of the MDA-MB-231 cancer cells in a dose-dependent manner (0.1, 1.0, 10, 50, 100 µg/ml). On the contrary, *V. angulatum* parasitic on *F. indica* exhibited no significant activity (LC_{50} - 500.82µg/ml) against the cancer cells (Fig. 4). Above observation suggest that antioxidant activity of *V. angulatum* might be responsible for its anticancer potential against breast cancer cells.




FIG. 4: CYTOTOXICITY ASSAY: Cytotoxic activity of methanolic extract of *V. angulatum* was examined on MDA-MB-231 cells. The cytotoxicity of *V. angulatum* extract (0.1-100 µg/ml) parasitic on different tree species was analyzed on MDA-MB-231 cells after an exposure of 24h. The methanolic extract of *V. angulatum* parasitic on *O. dioica* showed high cytotoxicity (LC_{50} – 79.33µg/ml) to MDA-MB-231 cancerous cells in comparison with that parasitic on *F. indica* (LC_{50} – 500.82 µg/ml). The values are expressed as the means ± S.D. (n = 3)

CONCLUSION: In conclusion, we report the host-associated variations in the phenolic, and flavonoid contents in the methanolic, ethanolic and aqueous extract of *V. angulatum* parasitic on host *O. dioica* and *F. indica* trees, which show differential antioxidant potential, and cytotoxicity against cancer cells. Despite the low phenolic and flavonoid content, the methanolic extract of plants parasitic on *O. dioica* exhibited strong scavenging activity against DPPH free radicals and cytotoxicity (LC_{50} - 79.33 µg/ml) against MDA-MB-231 cells. In contrast, the ethanolic extract of plants parasitic on *F. indica* showed a high phenolic content but low antioxidant activity and cytotoxicity. These results suggest the importance of antioxidant activity of *V. angulatum* in determining its anticancer potential. The study provides scientific evidence on the anticancer potential of the *V. angulatum* used in traditional medicine. It also suggests the significance of host tree in determining the therapeutic potential of *V. angulatum*. Further *in-vitro* and *in-vivo* studies along with isolation of active constituents are needed to get further insight into the role of the host plant in improving the efficacy of phytomedicines.

AUTHORS' CONTRIBUTIONS: SS and VM designed and supervised the experiments. SS performed the experiments. SS, VM, and RSS analyzed the data. SS wrote first draft and VM, RSS and MMS edited and finalized the manuscript. MMS helped during collection of plant material.

ACKNOWLEDGEMENT: VM and RSS thank University of Delhi for Faculty Research Grant and SS thanks, University Grant Commission, Government of India, for the Non-NET fellowship.

CONFLICTS OF INTEREST: The authors declare no conflict of interest.

REFERENCES:

- Nile SH and Park SW: Edible berries: Bioactive components and their effect on human health. *Nutrition*. 2014; 30: 134-44.
- Zaid H, Silbermann M, Amash A, Gincel D, Abdel-Sattar E and Sarikahya NB: Medicinal plants and natural active compounds for cancer chemoprevention/chemotherapy. *Evid Based Complement Alternat Med*. 2017; 7952417.
- Li S, Chen G, Zhang C, Wu M, Wu S and Liu Q: Research progress of natural antioxidants in foods for the treatment of diseases. *Food Science and Human Wellness*, 2014; 3(3-4): 110-16.
- Liu N, Lin Z, Guan L, Gaughan G and Lin G: Antioxidant enzymes regulate reactive oxygen species during pod elongation in *Pisum sativum* and *Brassica chinensis*. *PLoS ONE*, 2014; 9: 1-9. doi:10.1371/journal.pone.0087588
- Krishnaiah D, Sarbaty R and Nithyanandam R: A review of the antioxidant potential of medicinal plant species. *Food Bioprod Process*. 2011; 89: 217-233. doi:10.1016/j.fbp.2010.04.008
- Newman DJ and Cragg GM: Natural products as sources of new drugs over the 30 years from 1981 - 2010. *J Nat Prod*. 2012; 75(3): 311-335.
- Mishra V, Sharma RS, Yadav S, Babua CR and Singh TP: Purification and characterization of four isoforms of Himalayan mistletoe ribosome-inactivating protein from *Viscum album* having unique sugar affinity. *Arch Biochem Biophys*. 2004; 423: 288-301.
- Mishra R, Das MK, Singh S, Sharma RS, Sharma S and Mishra V: Articulatin-D induces apoptosis via activation of caspase-8 in acute T-cell leukemia cell line. *Mol Cell Biochem*. 2017; 426(1-2): 87-99.
- Beuth J, Ko HL, Gabius HJ, Burchard H, Oette K and Pulverer G: Behavior of lymphocyte subsets and expression of activation markers in response to immunotherapy with galactoside-specific lectin from mistletoe in breast cancer patients. *Clin Invest*. 1992; 70(8): 658-661.
- Patel S and Panda S: Emerging roles of mistletoes in malignancy management. *Biotech*. 2014; 4(1): 13-20. doi: 10.1007/s13205-013-0124-6.
- Hajto T, Hostanska K, Frei K, Rordorf C and Gabius HJ: Increased secretion of tumor necrosis factor- α , interleukin-1, and interleukin-6 by human mononuclear cells exposed to β -galactoside-specific lectin from clinically applied mistletoe extract. *Cancer Res*. 1990; 50: 3322-3326.
- Peumans WJ, Verhaert P, Fuller U and Van Damme EJ. Isolation and partial characterization of a small chitin-binding lectin from mistletoe (*Viscum album*). *FEBS Lett*. 1996; 396: 261-265.
- Büssing A and Schietzel M: Apoptosis-inducing properties of *Viscum album* Linn. extracts from different host trees, correlate with their content of toxic mistletoe lectins. *Anticancer Res*. 1999; 19: 23-28.
- Pietrzak W, Nowak R, Gawlik-Dziki U, Lemieszek MK, Rzeski W: LC-ESI-MS/MS identification of biologically

- active phenolic compounds in mistletoe berry extracts from different host trees. *Molecules*. 2017; 22(4): 624.
15. Song FL, Gan RY, Zhang Y, Xiao Q, Kuang L and Li HB: Total phenolic contents and antioxidant capacities of selected Chinese medicinal plants. *Int J Mol Sci*. 2010; 11: 2362-2372. doi:10.3390/ijms11062362
 16. Baba SA and Malik SA: Determination of total phenolic and flavonoid content, antimicrobial and antioxidant activity of a root extract of *Arisaema jacquemontii* Blume. *J Taibah Univ Sci*. 2015; 9(4): 449-454.
 17. Jha DK, Panda L, Ramaiah S and Anbarasu A: Evaluation and comparison of radical scavenging properties of solvent extracts from *Justicia adhatoda* leaf using DPPH assay. *Appl Biochem Biotechnol*. 2014; 174(7): 2413-25. doi: 10.1007/s12010-014-1164-z
 18. Marvibaigi M, Amini N, Supriyanto E, Abdul Majid FA, Jaganathan SK, Jamil S, Almaki JH and Nasiri R: Antioxidant activity and ROS-dependent apoptotic effect of *Scurrula ferruginea* (Jack) Danser methanol extract in human breast cancer cell MDA-MB-231. *PLoS ONE*. 2016; 11(7): e0158942.
 19. Singh S, Mishra R, Sharma RS and Mishra V: Phenol remediation by peroxidase from an invasive mesquite: Turning an environmental wound into wisdom. *J Hazard Mater*. 2017; 334: 201-211.
 20. Nazaruk J and Orlikowski P: Phytochemical profile and therapeutic potential of *Viscum album* Linn. *Nat Prod Res*. 2016; 30(4): 373-85. doi: 10.1080/14786419.2015.1022776
 21. Beuth J: Clinical relevance of immunoactive mistletoe lectin-I. *Anticancer Drugs*. 1997; 8: S53-S55.
 22. Onay-Uçar E, Karagöz A and Arda N: Antioxidant activity of *Viscum album* sp. album. *Fitoterapia*. 2006; 77(7-8): 556-60.
 23. Mishra V, Bilgrami S, Sharma RS, Kaur P, Yadav S and Krauspenhaar R: Crystal structure of himalayan mistletoe ribosome-inactivating protein reveals the presence of a natural inhibitor and a new functionally active sugar-binding site. *J Biol Chem*. 2005; 280: 20712-20721.
 24. Das MK, Sharma RS and Mishra V: A cytotoxic type-2 ribosome inactivating protein (from leafless mistletoe) lacking sugar binding activity. *Int J Biol Macromol*. 2011; 49(5): 1096-103.
 25. Das MK, Sharma RS and Mishra V: Induction of apoptosis by ribosome inactivating proteins: importance of N-glycosidase activity. *Appl Biochem Biotechnol*. 2012; 166(6): 1552-61.
 26. Mishra V, Ethayathulla AS, Sharma RS, Yadav S, Krauspenhaar R, Betzel C, Babu CR and Singh TP: Structure of a novel ribosome-inactivating protein from a hemi-parasitic plant inhabiting the northwestern Himalayas. *Acta Crystallogr D Biol Crystallogr*. 2004; 60(Pt 12 Pt 2): 2295-304.
 27. Luczkiewicz M, Cisowski W, Kaiser P, Ochocka R and Piotrowski A: Comparative analysis of phenolic acids in mistletoe plants from various hosts. *Acta Poloniae Pharmaceut-Drug Res*. 2001; 58(5): 373-379.
 28. Lin J, Chiou Y and Lin Y: Phenolic glycosides from *Viscum angulatum*. *J. Nat. Prod*. 2002; 65: 638-640.
 29. Jadhav RB, Bhatnagar SP and Surana SJ: Diuretic activity of squamate mistletoe, *Viscum angulatum*. *Pharmaceutical Biology*, 2010; 48(4): 417-421.
 30. Das MK, Sharma RS and Mishra V: A novel cationic peroxidase (VanPrx) from a hemi-parasitic plant (*Viscum angulatum*) of Western Ghats (India): Purification, characterization and kinetic properties. *J Mol Catal B Enzym* 2011; 71: 63-70.
 31. Kesarwani A, Chiang P and Chen S: Distribution of phenolic compounds and antioxidative activities of rice kernel and their relationships with agronomic practice. *Scientific World Journal*. 2014; 620171.
 32. Arts ICW and Hollman PCH: Polyphenols and disease risk in epidemiologic studies. *The American Am J Clin Nutr*. 2005; 81(1): 317S-325S.

How to cite this article:

Sharma S, Sharma RS, Sardesai MM and Mishra V: Anticancer potential of leafless mistletoe (*Viscum angulatum*) from Western Ghats of India. *Int J Pharm Sci Res* 2018; 9(5): 1902-07. doi: 10.13040/IJPSR.0975-8232.9(5).1902-07.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)