



Received on 28 July, 2017; received in revised form, 13 January, 2018; accepted, 27 January, 2018; published 01 May, 2018

## SYNTHESIS, CHARACTERISATION AND ANTIMICROBIAL ACTIVITY OF SOME NEW CHALCONES

M. Elavarasan, M. Thamizh Thendral and S. Syed Shafi \*

Department of Chemistry, Thiruvalluvar University, Serkkadu, Vellore - 632115, Tamil Nadu, India.

### Keywords:

Claisen-Schmidt condensation, Chalcone, Anti-bacterial, Anti-fungal

### Correspondence to Author:

**S. Syed Shafi**


Department of Chemistry,  
Thiruvalluvar University, Serkkadu,  
Vellore - 632115, Tamil Nadu, India.

**E-mail:** suban\_shafi@yahoo.com

**ABSTRACT:** In this work, an attempt was made to synthesize new chalcones by Claisen - Schmidt condensation of substituted benzaldehydes with 4-aminoacetanilide in the presence of base catalyst. The synthesized chalcone derivatives was characterized by FT-IR, <sup>1</sup>H-NMR, and <sup>13</sup>C-NMR and also tested for their antimicrobial activity and compared with the standard drugs.

**INTRODUCTION:** Chalcones are an important class of natural products and are considered as the precursors of flavonoids and isoflavonoids. Chemically chalcones are 1, 3-diaryl-2-propen-1-ones in which two aromatic rings are joined by a three carbon bridge having a carbonyl moiety and  $\alpha$ ,  $\beta$  -unsaturation. Chalcones come under an aromatic ketone that forms the central core for a variety of biological compounds. Claisen - Schmidt condensation between acetophenone and benzaldehyde gives chalcones, but in this reaction instead of acetophenone, we are using acetanilide with substitution benzaldehyde gives chalcones. This reaction is catalyzed by acids and bases under homogeneous or heterogeneous condition. Chalcones derivatives have received a great deal of attention due to their relatively simple structures and wide variety of pharmacological activities reported for these compounds include anti-inflammatory<sup>1,2</sup> antifungal<sup>3,4</sup>,

Antibacterial<sup>5</sup>, anti-malarial<sup>6</sup> and antitumor activities<sup>7</sup>. For these reasons, the synthesis of chalcones and their functionalized derivatives is a primary objective. Herein we report the activity of chalcone derivatives by synthesizing a series of molecules (2a-d) and evaluating their antibacterial activity against eight microorganism strains of Gram positive and negative antifungal profile against *Staphylococcus aureus* and *Penicillium* sps. In this study, only the feature associated with "B" ring of the chalcone moiety was changed by keeping the basic skeleton intact. Three compounds found most active *in-vitro* against *Staphylococcus aureus* and *Penicillium* sps. compared to standard drug.



**FIG. 1: SKELETON OF CHALCONE MOIETY**

**MATERIALS AND METHOD:** Melting points were determined in open capillary tubes and are uncorrected. All the chemicals and solvents used were of laboratory Grade and solvents were purified by suitable methods FTIR Spectra were

<p><b>QUICK RESPONSE CODE</b></p>	<p><b>DOI:</b> 10.13040/IJPSR.0975-8232.9(5).1969-73</p>
<p>Article can be accessed online on: <a href="http://www.ijpsr.com">www.ijpsr.com</a></p>	
<p>DOI link: <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.9(5).1969-73">http://dx.doi.org/10.13040/IJPSR.0975-8232.9(5).1969-73</a></p>	

recorded using KBr disc,  $^1\text{H}$ NMR and  $^{13}\text{C}$ NMR spectra were recorded on a Bruker 400MHz NMR spectra meter using TMS as an internal standard (chemical shift in  $\delta$ ppm) in  $\text{CDCl}_3$ . The Homogeneity of the products was checked by the TLC using silica gel 'G' and the eluent system was a mixture of hexane, ethyl acetate in 5:5 proportion.

**General Procedure for the Preparation of Chalcones:** A solution of 4-aminoacetanilide (0.002 mol) in absolute ethanol (20 ml) is kept in constant stirring with various aromatic aldehydes in the presence of 10% NaOH (20 ml) for 4 hr. After the reaction completes, the reaction mixture was poured in cooled water and kept in freezer at overnight, the product were purified by recrystallization from suitable solvents.


**N- (4-Aminophenyl) cinnamamide, (2a):** Orange-yellow powder, Yield 60%, m.p. 159 °C; TLC (Hexane: Ethylacetate, 5:5). IR: (KBr,  $\text{cm}^{-1}$ ) 3294 (N-H), 3185 (Aromatic C-H stretch), 2875 (C-H stretch), 1749 (NH-C=O), 1603 (CH=CH of carbonyl conjugated double bond), 1519 (C=C of aromatic ring).  $^1\text{H}$ NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ /ppm: 8.71 (s, 1H, N-H), 8.32-8.34 (t, 2H, -CO-CH=CH), 7.99-7.97 (d, 1H, =CH-Ar), 7.49-7.47 (d, 2H, Ar-H), 7.84-7.81 (d, 2H, Ar-H).  $^{13}\text{C}$ NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ /ppm:  $\delta$  168.57 (-NH-C=O),  $\delta$  135.94 (=C-Ar),  $\delta$  120.86; 121.69; 128.90; (Aromatic C), 135.29 (Carbonyl-C=).

**(E)- N- (4-Aminophenyl)- 3- (4-methoxyphenyl) acrylamide (2b):** Orange-yellow powder, Yield 80%, m.p. 145 °C; TLC (Hexane: Ethylacetate, (5:5). IR: (KBr,  $\text{cm}^{-1}$ ) 3238 (N-H), 3058 (Ar CH

stretch), 2846 (C-H stretch), 1658 (NH-C=O), 1540 (C=C of aromatic ring), 1310 (C-N).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ /ppm: 8.44 (ss, 1H, N- H), 7.51-7.52 (d, 1H, -CO-CH=), 7.25 (d, 1H, =CH-Ar), 6.95 (d, 2H, Ar-H), 7.16-7.17 (d, 2H, Ar-H), 3.86 (s, 2H, (NH<sub>2</sub>)), 2.16(s, O-CH<sub>3</sub>)  $^{13}\text{C}$ NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ /ppm:  $\delta$  162.38 (-NH-C=O),  $\delta$  135.94 (=C-Ar),  $\delta$  130.58; 129.30, 121.89; 121.58; 120.90 (Aromatic C), 114.30 (Carbonyl-C=),  $\delta$  55.40 (-O-CH<sub>3</sub>).

**(E)-N- (4-Aminophenyl)-3-(benzo[d][1, 3]dioxol-5-yl) acrylamide (2c):** Yellow powder, Yield 62%, m.p. 130 °C; TLC (Hexane: Ethylacetate, 2:8). IR: (KBr,  $\text{cm}^{-1}$ ) 3449 (N-H), 3221 (Ar C-H stretch), 3048 and 2901 (C-H stretch), 1657 (NH-C=O), 1593 (CH=CH of -Carbonyl-CH=CH-), 1540 (C=C of aromatic ring).  $^1\text{H}$ NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ /ppm: 8.33 (ss, 1H, N-H), 6.86-6.88 (d, 1H, -CO-CH=), 7.08-7.51 (m, 7H, Ar-H), 6.03(s, 2H, O-CH<sub>2</sub>-O).  $^{13}\text{C}$ NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ /ppm:  $\delta$  168.41 (-NH-C=O),  $\delta$  148.58 (=C-Ar),  $\delta$  106.92; 108.37; 120.81; 121.67; 121.93; 125.83; 131.31; 135.95 (Aromatic C), 120.01 (Carbonyl-C=).

**(E)- N- (4- Aminophenyl)- 3- (4- chlorophenyl) acrylamide (2d):** Light yellow powder, Yield 67%, m.p. 148 °C; TLC (Hexane: Ethylacetate, 5:5). IR: (KBr,  $\text{cm}^{-1}$ ) 3294 (N-H), 3186 (Ar C-H stretch), 3121 and 3061 (C-H stretch), 1661 (NH-C=O), 1568 (CH=CH of -Carbonyl-CH=CH-), 1542 (C=C of aromatic ring).  $^1\text{H}$ NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ /ppm: 10.01(s, 1H, OC-NH), 8.65 (d, 1H,-CO-CH=), 7.92-7.93(d, 2H, Ar-H), 7.57-7.63 (m, 4H, Ar-H), 2.05(s, 2H, NH<sub>2</sub>).



SCHEME - 1: SYNTHESIS OF CHALCONE DERIVATIVES (2a-d)

a) BENZALDEHYDE, NaOH/EtOH b) ANISALDEHYDE c) PIPERANAL d) CHLOROBENZALDEHYDE

Compound	R	M. P. (°C)	Yield (%)	Mol. formula	Mol. weight	Recrystal
2a	H	158-159	60	C <sub>15</sub> H <sub>14</sub> N <sub>2</sub> O	238.12	1
2b	O-CH <sub>3</sub>	144-145	80	C <sub>16</sub> H <sub>16</sub> N <sub>2</sub> O <sub>2</sub>	268.09	1
2c	O-CH <sub>2</sub> -O	130-132	62	C <sub>16</sub> H <sub>14</sub> N <sub>2</sub> O <sub>3</sub>	282.53	1
2d	4-Cl	148-149	67	C <sub>15</sub> H <sub>13</sub> N <sub>2</sub> OCl	274.64	1




FIG. 2: COMPOUND - 2A IR




FIG. 3: 2a <sup>1</sup>H NMR




FIG. 4: 2a <sup>13</sup>C NMR




FIG. 5: 2b IR




FIG. 6: 2b <sup>1</sup>H NMR




FIG. 7: 2b <sup>13</sup>C NMR




FIG. 8: 2c IR




FIG. 9: 2c <sup>1</sup>H NMR




FIG. 10: 2c <sup>13</sup>C NMR





FIG. 11: 2d IR

FIG. 12: 2d <sup>1</sup>H NMR

**RESULTS AND DISCUSSION:** In this reaction synthesis of chalcones carried out by the reaction of 4-aminoacetanilide and different aromatic aldehyde with 10% NaOH in absolute ethanol and the products were purified by recrystallization from suitable solvents. The synthesized product has been fully characterized by IR, <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectroscopy data. The IR spectrum of the synthesized chalcones was recorded and it gives an absorption band near 1700 - 1650 cm<sup>-1</sup> representing the presence of -C=O group. The absorption band at 1650 - 1580 cm<sup>-1</sup> confirms the aromatic -C=C- group. The <sup>1</sup>H NMR and <sup>13</sup>C NMR of synthesized chalcones give <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ/ppm: 6.70-6.38 (d, 1H, -CO-CH=), 7.99-7.97 (d, 1H, =CH-Ar) and <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>) δ/ppm: δ 166.13 (-NH-C=O), δ 138.48 (=C-Ar), 128.13 (Carbonyl-C=) confirms formation of chalcones.

### Pharmacological Results:

**Antibacterial Activity:** Nutrient Agar (NA) plates were seeded with 8 h broth culture of different bacteria. Sterile paper disc (6 mm in diameter) impregnated with 50 μl of different concentrations of samples were allowed to dry before being placed onto the seeded top layer of the agar plates. Each of the discs was gently placed at equidistance on top of the agar layer to give better contact with agar.

The plates were then incubated at 37 °C for 24 h. gentamycin (10 μg) was used as positive controls and DMSO / chloroform as negative control. The antibacterial activity was evaluated by measuring the diameter of inhibition zone. Solvent and growth controls were kept, the zones of inhibition and minimum inhibitory concentrations (MIC) noted. Results of these studies are compared with the standard.

**Antifungal Activity:** All the synthesized compounds were also screened for their in vitro antifungal activity against *Mucor*, *Aspergillus niger* and *Penicillium* strains. The zone of inhibition was measured in millimeters. Antifungal activities of all compounds were screened by the turbidometry method<sup>8,9</sup>. Activity of extract was compared with standard antibiotics fluconazole fungi. DMSO was used as solvent. All compounds are active against *Mucor*, *A. niger* and *Penicillium*.

### Results of Antibacterial Activity:

**Plates:** Inhibition zone of compounds K1, K2, K3 on *Staphylococcus aureus* and *Klebsiella pneumoniae* at 50 μg/mL.

Standard = Ciprofloxacin A – Compound K1, B – Compound K2, C – Compound K3

FIG. 13: INHIBITION ZONE OF COMPOUNDS K1, K2, K3 ON *STAPHYLOCOCCUS AUREUS* AND *KLEBSIELLA PNEUMONIAE*

S. no.	Microorganisms	Control (Solvent)	A	B	C	Ciprofloxacin
1	<i>Staphylococcus aureus</i>	-	6 mm	5 mm	10 mm	12 mm
2	<i>Klebsiella pneumonia</i>	-	4 mm	8 mm	7 mm	20 mm

### Results of Antifungal Activity:

**Plates:** Inhibition zone of compounds K1, K2, K3 on *Penicillium* sps and *Mucor* sps at 50µg/mL.

Standard = Amphotericin-B

A – Compound K1; B – Compound K2  
C – Compound K3



FIG. 14: INHIBITION ZONE OF COMPOUNDS K1, K2, K3 ON *PENICILLIUM* SPS AND *MUCOR* SPS

S. no.	Microorganisms	Control (Solvent)	D	E	F	Amphotericin-B
1	<i>Penicillium</i> sps	DMSO	7 mm	18 mm	10 mm	12 mm
2	<i>Mucor</i> sps	DMSO	7 mm	11 mm	8 mm	15 mm

**ACKNOWLEDGEMENT:** The author's wishes to express their thanks to, Department of Chemistry, Thiruvalluvar University for providing laboratory facilities and also Pondicherry University for spectral studies.

**CONFLICTS OF INTEREST:** Nil

### REFERENCES:

1. Yang HM, Shin HR, Cho SH, Bang SC, Song GY and Ju JH: Structural requirement of chalcones for the inhibitory activity of interleukin-5. *BioorgMedChem*2007;15:104-11.
2. Panchal AD, Prashant D and Patel KM: Synthesis and Biological Evaluation of Chalcone Derivatives *IJPSDR* 2011; 3(4): 331-337.
3. Sortino M, Delgado P, Juarez S, Quiroga J, Abonia R and Insuasty B: Synthesis and antifungal activity of (Z)-5-arylidenetherodanines. *Bioorg Med Chem* 2007; 15: 484-94.
4. Dominguez JN, Leon C, Rodrigues J, Gut J and Rosenthal PJ: Synthesis and Evaluation of New Antimalarial Phenylurenyl Chalcone Derivatives. *J. Med. Chem.* 2005; 48: 3654-8.
5. Valla A, Valla B, Cartier D, Guillou RL, Labia R and Florent L: New syntheses and potential antimalarial activities of new 'retinoid-like chalcones. *Eur. J. Med. Chem.* 2006; 41: 142-6.
6. Seo WD, Ryu YB, Curtis-Long MJ, Lee CW, Ryu HW and Jang KC: Evaluation of anti-pigmentary effect of synthetic sulfonylamino Chalcone. *Eur. J. Med. Chem.* 2010; 45: 2010-7.
7. Heindel ND and Reid JR: 4-Amino-3-mercapto-4H-1, 2, 4-triazoles and propargyl aldehydes: A new route to 3-R-8-aryl-1, 2, 4-triazolo[3, 4- b]-1, 3, 4-thiadiazepines. *J. Heterocycl. Chem.* 1980; 17(5): 1087.
8. Barry AL: The antimicrobial susceptibility test: principles and practices. Lea and Febiger Philadelphia, 1976; In A. L. Barry (ed.): 163-218.
9. Cooper KE: Analytical microbiology II. Academic Press Inc. New York, In F. Kavanagh (ed.) 1972; 13-30.

#### How to cite this article:

Elavarasan M, Thendral MT and Shafi SS: Synthesis, characterisation and antimicrobial activity of some new chalcones. *Int J Pharm Sci Res* 2018; 9(5): 1969-73. doi: 10.13040/IJPSR.0975-8232. 9(5).1969-73.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)