



Received on 30 July, 2017; received in revised form, 30 October, 2017; accepted, 27 January, 2018; published 01 May, 2018

## CHARACTERIZATION AND STABILITY OF EXTRACELLULAR ALKALINE PROTEASES FROM HALOPHILIC AND ALKALIPHILIC BACTERIA ISOLATED FROM SALINE HABITAT OF LONAR SODA LAKE, INDIA

Vijayshree M. Hemke<sup>1</sup> and Vishal R. Dhundale<sup>\*2</sup>

Jijamata Mahavidyalay<sup>1</sup>, Buldhana - 443203, Maharashtra, India.

New Arts Commerce and Science College<sup>2</sup>, Ahmednagar - 414001, Maharashtra, India.

### Keywords:

Bacillus, Extremozymes,  
Protease, Metalloenzyme

### Correspondence to Author:

**Vishal R. Dhundale**

Department of Microbiology,  
New Arts Commerce and  
Science College, Ahmednagar -  
414001, Maharashtra, India.


**E-mail:** vrdvishal@gmail.com

**ABSTRACT:** Culture dependent phenotypic characterization analyses were applied to study the protease producing bacteria from Indian Soda (Lonar crater). The uniqueness of the Lonar lake water is its salinity and alkalinity. A total of one hundred and fourteen bacterial cultures were isolated from the water and sediment samples collected from the hyper alkaline saline environment of Lonar lake. The thirty two bacterial strains were found protease producing bacterial culture, out of them strain EW2 was selected for 16S rDNA sequencing, production and partial characterizations of protease on the basis of their maximum proteolytic activity. The phylogenetic position indicated the strain was related to phylum Firmicutes. Optimum enzyme activity was found to be at 80 °C and pH 9.5. The protease was highly stable over broad temperature from 40 to 90 °C, pH 7.0 - 12.0, showing excellent thermostability and alkaline tolerant nature. The enzyme activity has enhanced by all metal ion which was tested in this studies indicating it was a metalloenzyme but inhibited by organic solvent except acetone. This is valuable information for enzyme production and optimization of protease from *Bacillus pumilus* EW2 has bright future towards the improvement and production of novel enzymes for leather, detergent and pharmaceuticals industries and biotechnological potential.

**INTRODUCTION:** The extracellular enzymes are secreted by wild group of organism comprise fungi bacteria, yeast actinomycetes<sup>1</sup>. Alkaliphilic micro-organisms, in particular *Bacillus* species, have attracted much interest because of their ability to produce extracellular metabolites that are active and stable at high pH. The unusual properties of these metabolites offer a potential opportunity for their utilization in processes demanding such extreme conditions<sup>2</sup>.

The reason for this is their high pH exhibiting range of pH 6 - 12 and temperature stability. Alkaline proteases belong to the group of serine proteases, or metalloenzyme<sup>3</sup>. Those from *Bacillus* species have wide use and importance in several industrial economies such as in food, detergent and in the synthesis of biologically active peptides<sup>4</sup>. Extremozymes have a great biotechnological potential, including agricultural, chemical and pharmaceutical applications<sup>5,6</sup>.

Due to the harsh condition in industrial production process, the present enzymes are not sufficient to meet most of the potential, so in last decades isolation and characterization of extremophiles have attracted attention<sup>7</sup>. Halalkaliphiles are salt-loving organisms survive in alkaline environment that inhabit hyper-saline alkaline environments.

<p style="text-align: center; font-weight: bold;">QUICK RESPONSE CODE</p> <div style="text-align: center;">  </div>	<p style="text-align: center;">DOI: 10.13040/IJPSR.0975-8232.9(5).1974-79</p> <hr/> <p style="text-align: center;">Article can be accessed online on: <a href="http://www.ijpsr.com">www.ijpsr.com</a></p>
<p>DOI link: <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.9(5).1974-79">http://dx.doi.org/10.13040/IJPSR.0975-8232.9(5).1974-79</a></p>	

They are characterized by large amounts of NaCl saturation due to the evaporative concentration. The progression and formation of alkalinity and salts also concentrate, giving rise to an alkaline saline environment<sup>8</sup>. Such type of condition generated in the soda lakes. Soda lakes are the most stable alkaline environments in the world, with pH values generally higher than 10 - 12<sup>9</sup>.

The bacteria survive in the soda lakes are mostly alkaliphilic, it is to be considered that extracellular enzymes synthesized by alkaliphilic bacteria would be diligent under alkaline stipulations. These extremozymes are active in absence or presence of trace amount of some metal ions and such enzymes are of biotechnological potential<sup>6</sup>. The Lonar lake is the great soda lake which is located in Lonar, District Buldhana, Maharashtra, India (latitude 19° 8, longitude 76° 36).

The alkaline Lonar crater is a unique basaltic rock meteorite impact crater, ranking third in the world and is filled with saline water having an average pH of 9.5 - 10. The Lonar lake was extensively studied for their bacterial diversity based on traditional and modern techniques with potential applications<sup>10 - 13</sup>. In view of the above facts, the present study focused on the screening and characterization of extracellular protease production from *Bacillus pumilus*.

## MATERIALS AND METHODS:

### Enrichment and Isolation of Microorganisms:

Lonar lake water and sediment sample were collected in sterile bottles and polythene bags respectively, from defined sampling site. Enrichment of water samples and sediment samples were carried out in Horikoshi I, Horikoshi II and nutrient agar at pH 10, nutrient agar at pH 10.0 with 30 g l<sup>-1</sup> sodium chloride. All flasks were incubated at 37 °C on a rotary shaker (100 rpm) for 48 h. After enrichment, well isolated and morphologically distinct colonies from these plates were transferred on the respective medium slants.

**Screening of Bacterial Alkaliphiles:** Individual bacterial colonies were screened for proteolytic activities on Skim milk agar medium. The inoculated plates were incubated at 37 °C for 48 hrs and observed for zones of clearance, represents proteolytic activities<sup>14, 15</sup>.

**Identification of the Proteolytic Isolates:** These isolates were identified in accordance with the methods recommended in Bergey's manual of determinative bacteriology and diagnostic microbiology. The identified strains were maintained on nutrient agar slants having pH 10 at 4 °C.

### 16S rDNA Sequences and Phylogenetic Analysis:

DNA was extracted from bacilli culture using standard phenol chloroform protocol<sup>16</sup>. The partial sequence of the 16S rRNA gene was amplified by using polymerase chain reaction and universal primer Eubacteria specific primers, 16F 27 (5' CCAGAATTGATCMTGGCTCAG- 3') and 16R 1525 (5' TTCTGCAGTCTAGAAGGAGGTGWT CCAGCC- 3'). The amplified 16S rRNA gene PCR products from these isolates were directly sequenced after purification by precipitation with polyethylene glycol and NaCl procedure<sup>17</sup>.

**Preparation of Crude Enzyme Extracts:** The 100 ml Casein nutrient broth (casein 1 gm, peptone 0.5 gm, yeast extract 0.15 gm, beef extract 0.15 gm, sodium chloride 0.5 gm, pH 10) was prepared. The sterile broth was inoculated with culture and incubated for 48 hrs at 37 °C. After 48 hrs incubation, centrifugation of the broth at 5000 rpm for 15 min was carried out<sup>18</sup>.

**Enzyme Assay:** The standard graph of tyrosine was prepared by adding different concentration of standard tyrosine (1 mg/ml). Estimation of proteases was carried out with 2.5 ml of casein in a test tube; 1 ml of enzyme source was added and incubated for 10 min at room temperature. After incubation 2 ml of TCA was added to stop the reaction and centrifuged the reaction mixture at 5000-8000 rpm for 15 min<sup>19</sup>.

**Partial Characterization of Protease:** Partial characterization of protease from *Bacillus* was carried out by effect of pH, temperature, substrate, enzyme, metal ions and organic solvents on alkaline protease activity were then measured as per assay procedure.

### Effect of Different Metal Ions on Protease Activity:

The effect of different metal ions on alkaline protease activity was determined. The enzyme assay was performed in the reaction mixture as described above in the presence of various metal ions at a final concentration of 1 mM.

The activity of the enzyme without any additives was taken as 100 %. The influence of various metal ions (1 mM each) on protease activity was studied by pre-incubating the enzyme with the compounds for 15 min at 37 °C. Then, the remaining activity was measured under the enzyme assay conditions.

#### Enzyme Stability in Presence of Solvents:

Stability of the protease enzyme was determined in the presence of various organic solvents (10 mM each). 0.25 ml of organic solvent was added to 1 ml of the protease solution in a 1.5 ml micro centrifuge tube with a screw cap. The mixture was shaken at 150 rpm for 100 min at 37 °C. Then, the remaining activity was measured under the enzyme assay conditions.

### RESULTS AND DISCUSSION:

**TABLE 1: MORPHOLOGICAL AND BIOCHEMICAL CHARACTERS OF BACILLUS SP. EW - 2**

S. no.	Test	Observation
1	Isolation Media	E
2	Color	White
3	Shape of Colony	Circular
4	Size of Colony	1 mm
5	Elevation	Convex
6	Edge	Entire
7	Gram Reaction	Positive
8	Shape	Short rod
9	Spore Reaction	+
10	Motility	+
11	Catalase	+
12	Oxidase	+
13	Glucose	+
14	Arabinose	+
15	Mannitol	-
16	Xylose	+
17	Sucrose	-
18	Maltose	-
19	Fructose	+

#### Screening of Bacterial Isolates, Identifications and Characterization of Bacterial Cultures:

There is very less studies were performed on the about the microbial diversity of hyper-saline alkaline Lonar crater especially from sediment and water characterized by alkaline and saline conditions. Bacillus EW-2, isolated from those microsites, shares with other alkaliphilic Bacillus species the capacity to grow at high pH values<sup>20, 21</sup>. During a previous study carried out from 2008 to 2017, total one hundred and fourteen alkaliphilic bacterial cultures were isolated from Lonar lake<sup>13, 22</sup>. In the present study, out of one hundred and fourteen, thirty two bacterial cultures were found with

proteolytic activity. All the bacterial strains were found alkaliphilic. According to Bergey's manual of systematic bacteriology, the isolates were identified as Bacillus sp. All the bacterial cultures were found gram positive and spore forming bacilli. Out of these thirty two, one bacterial isolate was selected for the further production and characterizations on the basis of their maximum proteolytic activity of Bacillus sp. EW-2. In the present studies the 16S rDNA analysis of isolates EW2 indicated that the bacterium affiliated to Firmicutes, which included family of bacillace, belong to *Bacillus pumilus* EW2.

#### Effect of pH on Activity of Enzyme Protease:

The effect of pH on protease activity of *Bacillus pumilus* EW-2 was determined by incubating the enzyme in different pH buffers ranging from 7 - 12 for 10 minutes at 37 °C. The enzyme was stable over the pH range of 7 - 12. The protease was active in a wide pH range with optimum activity at pH 9.5 and was stable over a broad pH range 7 - 12 with retention of more than 58 - 73 % of initial activity. The optimum activities of this enzyme were at pH 9.5 with 35 units/mL. The enzyme activity decreased dramatically at pH 10 (70%) and 12 (58%). The optimum pH range of alkaline proteases is between pH 9 and 11<sup>23</sup>. Alkaline protease from Bacillus EW-2 novel feature in the extremozyme which can produce the protease at alkaline pH which was beneficial for biotechnological potential and industrial application.

The similar studies were performed by Sookkheo *et al.*,<sup>24</sup> reported from thermophilic *Bacillus stearothermophilus* TLS33, optimum pH values of 8.5. To determine the optimum pH for enzyme activity, reactions were performed at various pH but optimum activity was found at pH 9.0 for *Elizabethkingia meningoseptica* KB042 isolated from chicken feathers by Nagal *et al.*,<sup>25</sup>.

#### Effect of Temperature on Activity of Enzyme Protease:

Influence of temperature on Bacillus sp. EW-2 protease activity was observed by incubating the enzyme at different temperature ranging from 35 - 100 °C and residual activity were determined under enzyme assay condition. The optimum activity of enzyme was taken as 100 %. The temperature profile of protease activity of Bacillus sp. EW-2 were showed maximal enzymatic activity

of (52 Units/ml) at 80 °C, which indicated that the enzyme was thermostable at high temperature. The protease retained more than 50% of the highest activity between 40 - 70 °C. Subsequently, the enzyme activity progressively decreased at 100 °C. In our study, although the bacterium was able to

grow well at temperature up to 50 °C, but it preferred thermophilic range of temperature for optimum activity at 80 °C (52 U/ml). The protease from *Bacillus sp* P2 was isolated and produce the thermostable protease and optimum activity was found at 90 °C<sup>26</sup>.




FIG. 1: EFFECT OF pH ON PROTEASE




FIG. 2: EFFECT OF TEMPERATURE ON PROTEASE




FIG. 3: EFFECT OF SUBSTRATE CONCENTRATION




FIG. 4: EFFECT OF ENZYME CONCENTRATION ON PROTEASE




FIG. 5: EFFECT OF ORGANIC SOLVENTS ON PROTEASE




FIG. 6: EFFECT OF METAL IONS ON PROTEASE

**Effect of Substrate Concentration on Activity of Enzyme Protease:** The influence of different concentrations of substrate was assayed ranging from 1 - 12 ml under constant assay conditions. The maximum enzymatic activity (80 Units/ml) at 4 ml of substrate concentration was considered as 100%. However, the activity was dramatically decreased as the substrate concentration increases more than 8 ml.

**Effect of Enzyme Concentration on Activity of Enzyme Protease:** The effects of different enzyme concentrations ranging from 1 - 12 ml was carried out under assay conditions. The enzyme shows maximum enzymatic activity (120 Units/ml) at 8 ml of enzyme concentration and it was considered as 100%. The activity of protease decreases as the enzyme concentration increases more than 9 ml.



**Influence of Different Metal Ions on Activity of Enzyme Protease:** The influence of different metal ions on activity of *Bacillus* sp. EW-2 protease was carried out under the assay conditions. Metal ions have different effects on activity of protease. The enzyme activity without any additives was taken as 100%. The enzyme activity was enhanced by KCl, NaCl, FeCl<sub>3</sub>, BaCl<sub>2</sub>, CaCl<sub>2</sub> and MgCl<sub>2</sub>. The optimum protease activity (316.67 %) was enhanced in presence of FeCl<sub>3</sub>.

**Influence of Various Organic Solvents on Activity of Enzyme Protease:** The effect of organic solvents on the activity of *Bacillus* sp. EW-2 protease was determined. Organic solvents have different effects on activity of protease. The enzyme activity without any additives was taken as 100%. The enzyme activity was enhanced only by Acetone. However, the protease activity was inhibited by ethanol, methanol, chloroform, benzene, hexane, ethyl acetate and acetic acid.

**CONCLUSION:** The *Bacillus pumilus* EW-2 strain isolated and extracellular synthesis of thermostable alkaline metalloprotease. Effects of metal ions and enhance the activity indicated that the alkaline protease belongs to the family of serine proteases. The optimum temperature at 80 °C and pH 9.5 for protease activity thereby give an opportunity to its biotechnological potential and to take super advantage in leather, detergent and pharmaceuticals industries as an eco-friendly.

**ACKNOWLEDGEMENT:** Nil

**CONFLICT OF INTEREST:** Author has no conflict of interest.

## REFERENCES:

- Hameş-Kocabaş EE and Uzel A: Alkaline protease production by an actinomycete MA1-1 isolated from marine sediments. *Annals Microbiol.* 2007; 57(1): 71-75.
- Dhundale VR, Nagarkar RD, More VR and Hemke VM: Partial characterization and optimization of alkaline amylase production from *Bacillus lehensis* isolated from alkaline saline Lonar lake. *Int. J. Advn. Biotech. Res.* 2014; 53: 1531-1536.
- Gençkal H and Tari C: Alkaline protease production from alkalophilic *Bacillus* sp. isolated from natural habitats. *Enz. Microbial Technol.* 2006; 39(4): 703-710.
- Kumar D and Bhalla TC: Microbial proteases in peptide synthesis: approaches and applications. *Appl. Microbial. Biotechnol.* 2005; 68(6): 726-736.
- Van den Burg B: Extremophiles as a source for novel enzymes. *Curr. Opin. Microbiol.* 2003; 6: 213-218.
- Ates O, Oner TE, Arıkan B, Denizci AA and Kazani D: Isolation and identification of alkaline protease producer halotolerant. *Bacillus licheniformis* strain BA17. *Annals of Microbiol.* 2007; 57 (3): 369-375.
- Patel R, Dodia M and Singh SP: Extracellular alkaline protease from a newly isolated haloalkaliphilic *Bacillus* sp. Production and optimization. *Proc. Biochem.* 2005; 40: 3569-3575.
- Grant WD and Tindall BJ: The isolation of alkaliphilic bacteria. In: Gould GW, Corry ICL (Eds.) *Microbial Growth and Survival in Extremes of Environment.* Academic Press, London, 1980; 27-36
- Jones BE, Grant WD, Duckworth AW and Owenson GG: Microbial diversity of Soda lakes. *Extremophiles.* 1998; 2: 191-200.
- Chaphalakar S and Dey S: Some aspects of production of extracellular protease from *Streptomyces diastaticus*. *J. of Micro. Biotechnol.* 1994; 9: 85- 100.
- Joshi AA, Kanekar PP, Kelkar AS, Shouche YS, Vani AA, Borgave SB and Sarnaik SS: Cultivable bacterial diversity of alkaline Lonar lake, India. *Microbial Ecology.* 2008; 55(2): 163-172.
- Surakashi VP, Antony CP, Sharma S, Patole MS and Shouche YS: Temporal bacterial diversity and detection of putative methanotrophs in surface mats of Lonar crater lake. *J. of Basic Microbiol.* 2010; 50: 465-474.
- Tambekar DH and Dhundale VR: Studies on the physiological and cultural diversity of bacilli characterized from Lonar lake (MS) India. *Biosci Discov.* 2012; 3(1): 34-39
- Dhundale VR, More VR, Nagarkar RD, Hemke VM and Tambekar DH: Isolation and characterization of a novel amylase from *Bacillus pseudofirmus* DW4 (1). *Indian J Life Science.* 2014; 4(1): 69-76.
- Hemke VM, Joshi SS, Fule NB, Dhundale VR and Tambekar DH: Phenetic diversity of alkaliphilic protease producing bacteria from alkaliphilic environment. *Indian J Sci Res.* 2015; 10(1): 47-52.
- Sambrook J, Fritsch EF and Maniatis T: *Molecular cloning: a Laboratory Manual*, Cold Spring Harbor, New York. Edition 2, 1989.
- Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA and Struhl, K: *Short protocols in molecular biology.* Wiley, New York, Edition 5, Vol. II, 1992.
- Gessesse A and Gashe BA: Production of alkaline protease by an alkaliphilic bacteria isolated from an alkaline Soda lake. *Biotechnology Letters.* 1997; 19(5): 479
- Tambekar DH, Tambekar SD, Jadhav AS and Babhulkar BV: Isolation and partial characterization of protease from *Bacillus halodurans* (AJ302709) from alkaline Lonar lake. *International Journal of Pharmaceutical Sciences and Research*, 2016; 7(11): 4546.
- Nielsen P, Fritze D and Priest FG: Phenetic diversity of alkaliphilic *Bacillus* strains: proposal for nine new species. *Microbiology.* 1995; 141: 1745-1761.
- Olivera N, Sequeiros C, Siñeriz F and Breccia JD: Characterization of alkaline proteases from a novel alkali-tolerant bacterium *Bacillus patagoniensis*. *World J. Microbiol Biotechnol.* 2006; 22(7): 737-743.
- Dhundale VR and Hemke VM: Phylogenetic analysis of Bacilli from haloalkaline Lonar Soda crater. *Int J Pharm Bio Sci* 2015; 6(4): (B) 27 -290.
- Kumar CG and Takagi H: Microbial alkaline protease; from a bio industrial view point. *Biotechnol Adv.* 1999; 17: 561-594.
- Sookkheo B, Sinchaikul S, Phutrakul S and Chen ST: Purification and characterization of the highly

- thermostable proteases from *Bacillus stearothermophilus* TLS33. Prot. Exp. Pur. 2000; 20: 142-151.
25. Nagal S, Kango N and Jain PC: Production of alkaline protease from *Elizabethkingia meningoseptica* KB042 using chicken feathers. Annals of microbiology. 2010; 60(4): 629-635.
26. Kaur S, Vohra RM, Kapoor M, Beg QK and Hoondal GS: Enhanced production and characterization of a highly thermostable alkaline protease from *Bacillus* sp. P-2. World J. Microbiol Biotechnol.2001; 17(2): 125-129.

**How to cite this article:**

Hemke VM and Dhundale VR: Characterization and stability of extracellular alkaline proteases from halophilic and alkaliphilic bacteria isolated from saline habitat of Lonar soda lake, India. Int J Pharm Sci Res 2018; 9(5): 1974-79. doi: 10.13040/IJPSR.0975-8232.9(5).1974-79.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)