



Received on 28 November, 2011; received in revised form 12 January, 2012; accepted 25 March, 2012

COMPLEMENTARY THERAPEUTIC POTENTIAL: A FOCUS ON POLYHERBAL PRODUCTS IN HYPERGLYCEMIA

S. Sujatha*, Jemima Shalin J. and Baby Joseph

International Centre for Bioresources Management, Malankara Catholic College, Mariagiri, Kaliyakkavilai - 629153, Kerala, India

ABSTRACT

Keywords:

Polyherbal,
Antidiabetics,
Hyperglycemia,
Antioxidant

Correspondence to Author:

S. Sujatha

International Centre for Bioresources
Management, Malankara Catholic College,
Mariagiri, Kaliyakkavilai - 629153, Kerala,
India

Nature has always been stands as a golden mark to amplify the outstanding phenomenon of symbiosis. Ayurvedic remedies for diabetes are usually mixed formulations containing blood sugar lowering herbs in combinations with immunomodulator, detoxicants with the rationale behind such formulations being provided by modern research. Polyherbal formulations have plant-based pharmacological agents which may exert synergistic, potentiative, agonistic antagonistic actions by virtue of its diverse active principles within themselves. These pharmacological principles work together in a dynamic way to produce maximum therapeutic efficiency with minimum side effects. Medicinal plants existing even before human being made their appearance on the earth. The development procedures of herbal drugs for world-wide use has to been different form that of synthetic drugs. The raw materials for Ayurvedic medicines were mostly obtained from plant sources in the form of crude drugs such as dried herbal powders or their extracts or mixture of products or mixture of product. Apart from these systems there has been a rich heritage of Ethno-botanical usage of herbs by various colorful tribal communities in the country. Hence, the present review provides the supportive evidence consideration of a therapeutic approach combining the beneficial effects of Polyherbal preparation in association with adaptive physical activity for effective management of diabetic complications.

INTRODUCTION: Diabetes affects about 5% of the global population (Chakraborty and Rajagopalan, 2002) and management of diabetes without any side effects is still a challenge to the medical system (Kameswara *et al.*, 2003). Diabetes is one of the most common chronic diseases in the world. The number of people with diabetes is increasing dramatically due to population growth, aging, urbanization, and increasing prevalence of obesity and physical inactivity that is finally associated with major health and socio-economic problems. The extent and severity of these problems are reflected in extra mortality and at-risk

people. For example, according to the last International Diabetes Federation (IDF) report, about 25,66,000 people (6% total population) are suffering from diabetes in Iran and its prevalence is increasing like other developing countries expecting to reach 5114900 in 2025 (Sicree *et al.*, 2007). Current estimates demonstrate that the world prevalence of diabetes will increase to 7.7% (439 million) adults by 2030. Between 2010 and 2030, there will be a 69% increase in number of diabetic patients in developing countries and 20% in developed countries (Shaw *et al.*, 2010). So, with regard to the issue of socioeconomic

burden of diabetes, discovery of more effective and less side effect therapies are necessary. In the recent years good data have been obtained from traditional medicines indicating usefulness of many herbal medicines (Hasani-Ranjbar *et al.*, 2008; 2009).

Hyperglycemia, Diabetes and Oxidative Stress:

Hyperglycemia is a connector between diabetes with diabetic complications (Brownlee, 2001; Rolo and Palmeira, 2006). In the review of Rolo four of the most important molecular mechanisms have been involved in hyperglycemia-induced tissue damage. Hyperglycemia-induced overproduction of superoxide is the causal link between high glucose and the pathways responsible for hyperglycemic damage. In fact, diabetes is typically associated with increased generation of free radicals and/or impaired antioxidant defense qualifications, representing a central contribution for reactive oxygen species in the onset, progression, and pathological consequences of diabetes.

Besides oxidative stress, some evidence has demonstrated a link between various disturbances in mitochondrial functioning and type-2 diabetes. Mutations in mitochondrial DNA and decreases in mitochondrial DNA copy number have been connected to the pathogenesis of type-2 diabetes (Rolo and Palmeira, 2006). Oxidative stress is increased in diabetes and is more definite in women and this leads to cardiovascular disease (Marra *et al.*, 2002). Modifications of life style through increased physical activity and reduced intake of calories can help lower the number of future cases of diabetes (Lakka *et al.*, 2002). Early recognition, treatment, and prevention of the metabolic syndrome present a major challenge for health care professionals Confronting an epidemic of overweight and sedentary lifestyle (Lakka *et al.*, 2002).

People with diabetes do not have enough antioxidant defenses (Martin-Gallan *et al.*, 2003), but, in contrast, too much of the free radical-induced damage can occur. Oxidative stress takes place while oxygen free radicals are produced in a very large amount through the diminution of oxygen. Diverse surveys were intended to establish the levels of stress associated biomarkers in type-1 and 2 (Varvarovska *et al.*, 2003).

Medicinal plants and its anti-diabetic potential:

Interntional Scenario: For a very long time, plants have been an important role part of treatment of many diseases. The use of plants to treat diabetes is a centuries-old practice. More than 400 traditional plant treatments for diabetes have been recorded but only a small number of these have received scientific and medical evaluation to assess their efficacy. Hypoglycemic action from some treatments has been confirmed in animal models, and various hypoglycemic compounds have been identified. Traditional treatments may provide valuable clues for the development of new oral hypoglycemic agents and simple dietary adjuncts.

Urtica dioica (Stinging Nettle) has been used for centuries for food and medical purposes. The species name *dioica* means two houses because the plant usually contains either male or female flowers. It is abundant in North America, Northern Europe and most of Asia, usually found in the rural area. It contains flavonoids (0.7-1.8%), silicic acid (1-4%), Potassium-ions (0.6%), Nitrates (1.5-3%), volatile oil, Histamine, Serotonin, acetylcholine, formic acid and leukotrienes (LTB₄, LTC₄, LTD₄). The blood sugar lowering effect of *Urtica dioica* has been mentioned in old script such as those written by Avicenna. There have been other reports indicating the benefits of using the infusion or the extract of the leaves or other parts of this plant for the use in diabetes (Ramos *et al.*, 1992; Swanston-Flatt *et al.*, 1989). Moreover, it is used internally and externally as supportive therapy for prostatic hyperplasia (Krzieski *et al.*, 1993; Kayser *et al.*, 1995), inflammation (Obertreis *et al.*, 1996), rheumatoid arthritis, hypertension and allergic rhinitis (Mittman, 1990).

Medicinal plants and its anti-diabetic potential:

Indian Scenario: India represented by rich traditional practices, the present research was focused on culture, traditions and natural biodiversity, offers a unique the antibiotic activity of one new species leads to an opportunity for drug discovery researchers (Jachak and Saklani, 2007). Since ancient times, plants have been an exemplary source of medicine. Ayurveda and other Indian literature mention the use of plants in treatment of various human ailments. India has about 45 000 plant species and among them, several thousands have been claimed to possess medicinal

properties. Many researchers conducted in last few decades on plants mentioned in ancient literature or used traditionally for diabetes have shown anti-diabetic property. Previously, Grover et al., 2002 reviewed 45 such plants and their products (active, natural principles and crude extracts) that have been mentioned/ used in the Indian traditional system of medicine and have shown experimental or clinical anti-diabetic activity. Indian plants which are most effective and the most commonly studied in relation to diabetes and their complications are: *Allium cepa*, *Allium sativum*, *Aloe vera*, *Cajanus cajan*, *Coccinia indica*, *Caesalpinia bonducella*, *Ficus bengalensis*, *Gymnema sylvestre*, *Momordica charantia*, *Ocimum sanctum*, *Pterocarpus marsupium*, *Swertia chirayita*, *Syzgium cumini*, *Tinospora cordifolia* and *Trigonella foenum graecum*. Among these we have evaluated *M. charantia*, *Eugenia jambolana*, *Mucuna pruriens*, *T. cordifolia*, *T. foenum graecum*, *O. sanctum*, *P. marsupium*, *Murraya koeingii* and *Brassica juncea*. All plants have shown varying degree of hypoglycemic and anti-hyperglycemic activity (Table 1).

TABLE 1: INDIAN MEDICINAL PLANTS WITH HYPOGLYCEMIC ACTIVITY

Name of the plants	Family
<i>Acacia arabica</i> (Lam.) Muhl. ex Willd.	Mimosaceae
<i>Aegle marmelos</i> (L.) Correa ex Roxb.	Rutaceae
<i>Allium cepa</i> L.	Liliaceae
<i>Allium sativum</i> L.	Alliaceae
<i>Aloe vera</i> (L.) Burm.f.	Aloaceae
<i>Areca catechu</i> L.	Arecaceae
<i>Artemisia pallens</i> Wall. ex DC.	Compositae
<i>Annona squamosa</i> L.	Annonaceae
<i>Andrographis paniculata</i> Nees	Acanthaceae
<i>Aerva lanata</i> (L.) Juss. ex Schult.	Amaranthaceae
<i>Azadirachta indica</i> A. Juss.	Meliaceae
<i>Biophytum sensitivum</i> (L.) DC.	Oxalidaceae
<i>Bombax ceiba</i> L.	Bombacaceae
<i>Beta vulgaris</i> L.	Chenopodiaceae
<i>Brassica juncea</i> (L.) Czern.	Brassicaceae
<i>Barleria lupulina</i> Lindl.	Acanthaceae
<i>Boerhavia diffusa</i> L.	Nyctaginaceae
<i>Cassia auriculata</i> L.	Leguminosae
<i>Caesalpinia bonducella</i> (L.) Roxb.	Cesalpinaceae
<i>Capparis decidua</i> (Forsk.) Edgew.	Capparidaceae
<i>Cajanus cajan</i> (L.) Millsp.	Fabaceae
<i>Citrullus colocynthis</i> (L.) Schrad.	Cucurbitaceae
<i>Coccinia indica</i> Wight & Arn.	Cucurbitaceae
<i>Casearia esculenta</i> Roxb.	Flacourtiaceae
<i>Catharanthus roseus</i> (L.) G. Don.	Apocynaceae
<i>Camellia sinensis</i> Kuntze	Theaceae
<i>Eugenia uniflora</i> L.	Myrtaceae
<i>Eucalyptus globulus</i> Labill.	Myrtaceae

<i>Enicostemma littorale</i> Blume	Gentianeae
<i>Biophytum sensitivum</i>	Laksmana
<i>Eugenia jambolana</i> Lam.	(Myrtaceae)
<i>Ficus bengalensis</i> L.	Moraceae
<i>Gymnema montanum</i> Hook.f.	Asclepiadaceae
<i>Gymnema sylvestre</i> R. Br.	Asclepiadaceae
<i>Glycyrrhiza glabra</i> L.	Fabaceae
<i>Hibiscus rosa sinensis</i> L.	Malvaceae
<i>Helicteres isora</i> L.	Sterculiaceae
<i>Ipomoea batatas</i> (L.) Lam.	Convolvulaceae
<i>Lantana camara</i> L.	Verbenaceae
<i>Mangifera indica</i> L.	Anacardiaceae
<i>Memecylon umbellatum</i> Burm. f.	Melastomataceae
<i>Momordica cymbalaria</i> Fenzl ex Naudin	Cucurbitaceae
<i>Musa sapientum</i> L.	Musaceae
<i>Momordica charantia</i> L.	Cucurbitaceae
<i>Morus alba</i> L.	Moraceae
<i>Murraya koeingii</i> (L.) Spreng.	Rutaceae
<i>Nelumbo nucifera</i> Gaertn.	Nymphaeaceae
<i>Ocimum sanctum</i> L.	Lamiaceae
<i>Picrorrhiza kurroa</i> Royle ex Benth.	Scrophulariaceae
<i>Phyllanthus amarus</i> Schumach. & Thonn.	Euphorbiaceae
<i>Pterocarpus marsupium</i> Roxb.	Fabaceae
<i>Punica granatum</i> L.	Punicaceae
<i>Pterocarpus santalinus</i> L. f.	Leguminosae
<i>Salacia reticulata</i> Wight.	Celastaceae
<i>Salacia oblonga</i> Wall.	Celastaceae
<i>Swertia chirayita</i> (Roxb. ex Fleming) H. Karst.	Gentianaceae
<i>Scoparia dulcis</i> L.	Scrophulariaceae
<i>Syzgium alternifolium</i> Walp.	Myrtaceae
<i>Sida cordifolia</i> L.	Malvaceae
<i>Trigonella foenum graecum</i> L.	Fabaceae
<i>Terminalia catappa</i> L.	Combretaceae
<i>Terminalia pallida</i> Brandis	Combretaceae
<i>Tinospora cordifolia</i> (Willd.) Hook.f. & Thomson	Menispermaceae
<i>Vinca rosea</i> (Catharanthus roseus):	Roseaceae
<i>Zingiber officinale</i> Roscoe	Zingiberaceae
<i>Zizyphus sativa</i> Gaertn.	Rhamnaceae

Enumeration of Antidiabetic plants: Several medicinal plants have found potential use as hypoglycemic in the Indian system of medicines, including ayurveda. Many Indian plants have been investigated for their beneficial use in different types of diabetes and reports occur in numerous scientific journals. This article aims to provide a comprehensive review on various plant species from Indian biosphere and their constituents, which have been shown to display potent hypoglycemic activity. The use of herbs as hypoglycemic is a major avenue in Indian perspectives particularly for treating diabetes, which require to be explored more effectively as there are so many

literatures available on these aspects. This paper describes the chemistry, activity and usage of the constituents isolated from these plants from India for the treatment of diabetes recorded by Pulok *et al.* (2006). Furthermore, Nagendra *et al.*, (2010) enumerated that the *C. orehioides* is the black gold with numerous health benefits which is considered to have aphrodisiac, immunostimulant, hepatoprotective, antioxidant, anticancer and antidiabetic activity.

Traditional Herbal Medicines Used for the Treatment of Diabetes: In recent years, there has been renewed interest in the treatment against different diseases using herbal drugs as they are generally non-toxic and World Health Organization has also recommended the evaluation of the effectiveness of plants in condition where we lack safe modern drugs (Mohseni-Salehi *et al.*, 2010). Plant derivatives with hypoglycaemic properties have been used in folk medicine and traditional healing systems around the world (Yeh *et al.*, 2003) from very ancient time. Despite the introduction of hypoglycaemic agents from natural and synthetic sources, diabetes and its secondary complications continue to be a major medical problem to people (Ravi *et al.*, 2005). In addition, Medicinal plants used to treat hypoglycemic and hyperglycemic conditions are of considerable interest to ethnobotanical community as they are recognized to contain valuable medicinal properties in different parts of the plant depicted by very recently (Lan *et al.*, 2011).

In traditional medicine diabetes mellitus is treated with diet, physical exercise and medicinal plants, even though, more than 1200 plants are used around the world in the control of diabetes mellitus and approximately 30% of the traditionally used antidiabetic plants were pharmacologically and chemically investigated (Alarcon-Aguilar *et al.*, 2002). Again Ayyanar *et al.*, 2008 studied the potential hypoglycaemic agents have also been detected for more than 100 plants used in antidiabetic therapy. Traditional treatments may provide the valuable clues for the development of new oral hypoglycemic agents and simple dietary adjuncts Mukherjee *et al.*, 2006. More than 100 medicinal plants are mentioned in the Indian system of medicines including folk medicines for the management of diabetes, which are effective either separately or in combinations (Kar *et al.*, 2003).

As per the ethnobotanical literature on traditional Phytotherapy of Indian medicinal plants, the species like *Asparagus racemosus*, *Butea monosperma*, *Catharanthus roseus*, *Coccinia indica*, *Gymnema sylvestre*, *Syzygium cumini* and *Momordica charantia* are consistently used by the tribal communities for the treatment of diabetes (Kumarappan *et al.*, 2007) as well as in modern medicine.

The oxidative stress due to free radicals (H_2O_2 , oxy radicals O_2^- etc.) lead to many diseases such as atherosclerosis (DeWitt Dawn. and Hirsch, 2009) and *Diabetes mellitus* (Rahimi *et al.*, 2005). It is essential to counteract this oxidative stress and there by enhance the immunity of body system. Allopathic drugs are available for counteracting the oxidative stress and hence improve immunity, but the side effects and prohibitive cost of these allopathic drugs makes it necessary to search for an alternative (Fazeli *et al.*, 2008). The Ayurvedic system of medicines not only provides that alternative, but also scores over the side effects and cost factor of allopathic medicine (Donga *et al.*, 2011).

Guard Sansar (GS) is a Polyherbal formulation, developed by Pradhan Herbal Company, Bangalore. GS consists of extracts of seven medicinal plants, which are *Embllica officinalis*, *Tinospora cardifolia*, *Terminalia arjuna*, *Piper longum*, *Comiphora mukul*, *Nardostachys jatamansi* and *Boerhavia diffusa* as well as corals and asphaltum. These plant extracts are classified, in ayurveda, as Rasayanas which improve defense mechanism of body and possess antioxidant activity. Earlier studies indicate that *Embllica officinalis* (Sarkhail *et al.*, 2010), *Tinospora cardifolia*, *Terminalia arjuna* and *Piper longum* possess immunomodulatory and antioxidant activity (Petlevski *et al.*, 2001). As there is a need to generate scientific evidence/preclinical data to support the clinical use of the ayurvedic medicine (s), the present study was undertaken to investigate and to validate the immunomodulatory and antioxidant activity of Guard Sansar.

Current trends in D.M: Current available therapies for diabetes include insulin and various oral antidiabetic agents such as sulfonylurea, biguanides, α -glycosidase inhibitors and glinides, which are used immunotherapy or in combination to achieve better glycerin regulation (Varvarovska *et al.*, 2003). The use of oral drugs is

limited due to adverse side effects including hematological, cardiovascular and gastrointestinal reactions, hypoglycemic coma and disturbance of liver and kidney functions in addition, they are not suitable for use during pregnancy (Alarcon *et al.*, 2000). In the treatment of diabetic mellitus non-pharmacologic measures [leg diet, exercise, relaxation therapy, meditation and music therapy remains a critical component of therapy (Shaw *et al.*, 2010). To avoid the serious side effects of complementary the therapies, technically another survey of alternative treatments used by patients with diabetes (Mobaseri *et al.*, 2010). A part from currently available therapeutic options many herbal medicines have been recommended for the Treatment of diabetes medicinal plants have the advantage of having no side effects (Mohseni-Salehi *et al.*, 2010).

Effectuated Biological product in D.M: Plants have always been utilized as sources of drugs and many of the currently available drugs have been directly or indirectly obtained from plant sources (Jachak and Saklani, 2007). Herbal medicines derived from plant extract are being used to treat a wide variety of clinical disease (Kar *et al.*, 2003). More attention has paid to the protective effects of natural antioxidants against drug induced toxicity studies especially whenever free radical generation is involved (Mobaseri *et al.*, 2010). Flavonoids have been found to play a very important role in protection against oxidative stress (Said *et al.*, 2008).

In accordance with the recommendations of the world health organization (WHO) committee on diabetes mellitus it is important to investigate the hypoglycemic actions from plants which were originally used in traditional medicine (Jachak and Saklani, 2007). According to when the herbal remedies may have recognizable therapeutic effects compared with synthetic drugs, derived from the plants are frumenty considered to be less toxic described by Momtaz and Abdollahi, (2010). Therefore, the search for the more effective and safer anti hyperglycemic agents becomes an area of active research.

The effective botanical products are available in capsules and tablets, as well as in other forms, such as water extracts, (also called decoctions or infusions), tinctures (hydro alcoholic extracts) and glycerites

(glycerin – extracted preparations that are alcohol free). All vary in potency (21) Searching for new antidiabetic drugs from natural plants is still attractive because they contain substances which take alternative and safe effect on diabetes mellitus (Domola *et al.*, 2010). Most of plants contain glycosides, alkaloids, terpenoids, flavonoids, carotenoids, etc that are frequently implicated as having antidiabetic effect (Grover *et al.*, 2002).

Status of Diabetes mellitus disease in the world:

Diabetes is one of the most common chronic diseases in the world. The number of people with diabetes is increasing dramatically due to population growth, aging, urbanization, and increasing prevalence of obesity and physical inactivity that is finally associated with major health and socio-economic problems (El-Haouari *et al.*, 2007; Ayyanar *et al.*, 2008). The extent and severity of these problems are reflected in extra mortality and at-risk people. For example, according to the last International Diabetes Federation (IDF) report, about 2566000 people (6% total population) are suffering from diabetes in Iran and its prevalence is increasing like other developing countries expecting to reach 5114900 in 2025 (Sicree *et al.*, 2007; Fazeli *et al.*, 2008). Current estimates demonstrate that the world prevalence of diabetes will increase to 7.7% (439 million) adults by 2030. According to Shaw *et al.* (2010) statement between 2010 and 2030, there will be a 69% increase in number of diabetic patients in developing countries and 20% in developed countries. So, with regard to the issue of socioeconomic burden of diabetes, discovery of more effective and less side effect therapies are necessary. In the recent years good data have been obtained from traditional medicines indicating usefulness of many herbal medicines (Hasani - Ranjbar *et al.*, 2008, 2009). For a very long time, plants have been an important role part of treatment of many diseases.

More than 400 traditional plant treatments for diabetes have been recorded, but only a small number of these have received scientific and medical evaluation to assess their efficacy depicted by Fathi-Azad *et al.* (2005). Hypoglycemic action from some treatments has been confirmed in animal models, and various hypoglycemic compounds have been identified It contains flavonoids (0.7-1.8%), silicic acid (1-4%), potassium-ions (0.6%), nitrates (1.5-3%), volatile oil,

histamine, serotonin, acetylcholine, formic acid and leukotrienes (LTB₄, LTC₄, LTD₄) reported by Bijan *et al.* (2003). The blood sugar lowering effect of *Urtica dioica* has been mentioned in old script such as those written by Avicenna. There have been other reports indicating the benefits of using the infusion or the extract of the leaves or other parts of this plant for the use in diabetes (Khouri. *et al.*, 2006; Khairandish *et al.*, 2009). Moreover, it is used internally and externally as protective activity in beta cells (Golalipour and Khouri, 2007), stereological studies in liver cells (Golalipour *et al.*, 2007), then morphological as well as histological alterations in renal and liver depicted by (Golalipour *et al.*, 2009a and b) and proliferation of pancreas in diabetic rat (Golalipour *et al.*, 2010).

Effects of polyherbal formulations: Polyherbal formulations have plant-based pharmacological agents which may exert synergistic, potentiative, agonistic antagonistic actions by virtue of its diverse active principles within themselves. These pharmacological principles work together in a dynamic way to produce maximum therapeutic efficacy with minimum side effects (Tiwari and madhusudana Rao, 2002). According to Ayurvedic texts, a combination of substances is used to get the enhanced described action and eliminate unwanted side effects (Pulok *et al.*, 2006).

Ashok *et al.*, (2010) investigated, antidiabetic and antioxidant values of a polyherbal formulation, Gly-13-C (*Pterocarpus marsupium*, *Azadirachta indica*, *Swertia chirata*, *Curcuma longa*, *Gymnema sylvestre*, *Emblica officinalis*, *Eugenia jambolana*, *Salacia reficulata*, *Trigonella foenum graecum*, *Vince rosea*, *Aegle marmelos* and two minerals viz., processed stannum and processed asphaltum in streptozotocin induced diabetic male Wistar rats. He also demonstrated that Gly-13-C provided better glycemic control and significantly protected liver tissue against oxidative damage than reference drug.

Karnim plus, polyherbal formulation containing the *Momordica charantia*, *ocimum sanctum*, *Azadirachta*

indica, *Picrorrhiza sanctum* and *Zingiber officinale* has been selected by Martin-Gallan *et al.* (2003). It was helpful in reducing the lipid profile seen in some diabetics in whom hyperglycemia and hypercholesterolemia co exist quite. Hence this product has been used as one of the alternative remedy for the treatment of diabetes.

Sitasawad *et al.* (2000) reported that MD-19 is a polyherbal formulation has been screened for antidiabetic activity in alloxan induced *diabetes mellitus*. MD-19 is a combination of medicinal plants with mineral namely *Momordica muricata*, *Tinospora cordifolia*, *Trigonella foenum graecum*, *Caesalpinia bonducella*, *curcuma longa*, along with shilajit (mineral). Treatment of diabetic rats with MD-19 restored the biochemical parameters significantly and it is also proved to be effective in nephrotoxicity and anti hyperglycemic activity (Kusano and Abe, 2000).

Fulzele *et al.*, 2002) projected on DRF/AY/5001, an indigenous polyherbal formulation containing medicinal plants such as *Gymnema sylvestre*, *Syzygium cuimini*, *Pterocarpus marsupium*, *Momordica charantia*, *Emblica officinales*, *Terminalia betericica*, *Terminalia chebula* and shudh shilajit (Asphaltum). This formulation has been selected to evaluate the antidiabetic and its related mechanism(s) of action. "Diabrid" a herbal based formulation has been carried out by Gupta and Kumar (2008) to evaluate the hypoglycemic activity in mild and moderate diabetes. The herbal constituents of Diabrid contain four indigenous medicinal plants namely *Gymnema sylvestre*, *Momordica charantia*, *Eugenia jambolana* and *Trigonella Graeceium*.

Furthermore, Hasani-Ranjbar *et al.* (2008) investigated Diabet, a polyherbal formulation intended to use for diabetic patients has been screened for antioxidant activity. The formulation exhibited significant antioxidant activity showing increased levels of SOD, CAT, GPx, GSH and decreased level of LPO.

TABLE 2: POLYHERBAL MEDICINAL PLANTS EFFECTIVE ON *DIABETES MELLITUS*

Botanical Name	Parts used	Views	Medicinal Properties
<i>Curculigo orchioides</i>	Tuber		Piles, jaundice, gonorrhoea, bronchitis, diarrhoea, dyspnoea, gleet, hydrophobia, pain in the joints, diabetes, cancer etc.
<i>Scoparia dulcis</i>	Whole plant		Wounds, ulcer, kidney stone skin disease. It increases the activities of plasma insulin, pain killers, stomach troubles, diuretics, fever.
<i>Santalum album</i>	Root		Urogenital (internal) and skin (external) antiseptic, dysuria and also useful in tuberculosis of gall bladder.
<i>Tribulus terrestris</i>	Root		Piles, frequency of urination in diabetes, cystitis and urinary tract infection, enhances sex hormones.
<i>Terminalia chebula</i> Retz.	Fruit		Stomach disorders, piles, fistula, chronic fever, cough, coryza, sore throat, asthma, renal calculi, diabetes.
<i>Cuminum cyminum</i> Linn.	Seeds		Antiseptic, anti-spasmodic, antitoxic, bactericidal, diuretic, emmenagogue, nervine, tonic, osteoarthritis, headaches.
<i>Ipomoea turpetum</i> L.	Root		It is used in the treatment of diarrhea, acne, and diabetes.

<i>Trigonella foenum-graecum</i> L.	Seeds		Hypoglycaemic, laxative, parasiticide, restorative, uterine toxic, anticholesterolemic, anti-inflammatory, antitumor, antidiabetic.
<i>Michelia champaca</i> L	Flower		Kidney diseases, anti-diabetic, nausea and fever. The oil extracted from the flowers is locally used in headache, ophthalmia, rheumatism and gout.
<i>Glycyrrhiza glabra</i> Linn.	Root		Arteriosclerosis, adjusts blood salts, balances blood sugar, cancer, circulation, colds, flu, heart strengthener, hypoglycemia.
<i>Centella asiatica</i> (L) Urban.	Whole plant		Antibacterial, anti-viral, anti-inflammatory, anti-ulcerogenic, anxiolytic, a cerebral tonic, a circulatory stimulant, anti-diabetic.
<i>Andrographis paniculata</i> Nees.	Whole plant		Sore throat, hepatitis, dysentery, cholera, diabetes, influenza, bronchitis, swellings and itches, piles, diabetes
<i>Withania somnifera</i>	Root		Dehydration, bone weakness, muscle weakness, tension, premature ageing, Parkinson's disease, bone cancer, bipolar disorder, diabetes.
<i>Salacia oblonga</i>	Root		Antidiabetic, anticancer, antibacterial, antifungal.

Conclusion: This review paper has been presented cumulative Polyherbal antidiabetic medicinal plants used in the treatment of *Diabetes mellitus* which can provide a comprehensive review on various plant species from Indian biosphere and their constituents. Many Indian plants have been investigated for their beneficial use in different types of diabetes and reports occur in numerous scientific journals. The use of herbs as hypoglycemic is a major avenue in Indian perspectives particularly for treating diabetes. Several new bioactive Polyherbal drugs (Secondary metabolites) isolated from vast kinds of medicinal plants having hypoglycemic effects showed antidiabetic activity commonly with more vibrant as well as most powerful than commonly available oral hypoglycaemic agents such as Daonil, Tolbutamide and Chlorpropamide. Though, number of bioactive compounds extracted, isolated and utilized for varieties of medicinal plants have not been well characterized.

REFERENCES:

- Alarcon-Aguilar, F.J., Roman-Ramos, R., Flores-Saenz, J.L., Aguirre-Garcia, F., 2002. Investigation on the Hypoglycaemic Effects of Extracts of Four Mexican Medicinal Plants in Normal and Alloxan-diabetic Mice. *Phytotherapy Research*, 16, 383 – 386.
- Ayyanar, K. Sankarasivaraman and S. Ignacimuthu. 2008. Traditional Herbal Medicines Used for the Treatment of Diabetes among Two Major Tribal Groups in South Tamil Nadu, India, *Ethanobotanical Leaflets*, 12: 276-280.
- Bijan, F., D. Ahmadv and, S. Vardasbi, F.J. Majin and SH. Khaghani, 2003. Induction of insulin secretion by a component of *Urtica dioica* leave extract in perfused islets of langerhans and its *in vivo* effects in normal and streptozotocin diabetic rats. *Ethnopharmacology*, 89: 47-53.
- Brownlee, M. 2001. Biochemistry and molecular cell biology of diabetic complications. *Nature.*, 414:813-820.
- Chakraborty R and Rajagopalan R. 2002: Diabetes and insulin resistance associated disorders: disease and the therapy. *Current Science*, 83: 1533-1538.
- DeWitt, D. and Hirsch. 2009. Outpatient Insulin Therapy in Type-1 and Type 2 Diabetes Mellitus. *JAMA.*, 17: 2254-2264.
- Donga J.J., Surani V.S., Sailor G.U., Chauhan S.P., Seth A.K. 2011. A systematic review on natural medicine used for Therapy of diabetes mellitus of some Indian Medicinal plants. *An International Journal of Pharmaceutical Sciences*, 2, (1): ISSN: 0976-7908
- Domola, M.S., V. Vu, C.A. Robson-Doucette, G. Sweeney and M.B. Wheeler, 2010. Insulin mimetics in *Urtica dioica*: Structural and computational analyses of *Urtica dioica* extracts. *Phytotherapy Res.*, 24: S175-S182.
- El-Haouari, M., I. Jardin, H. Mekhfi, J.A. Rosado and Salido, G.H. 2007. *Urtica dioica* extract reduces platelet hyper agreeability in type-2 diabetes mellitus by inhibition of oxidant production, Ca²⁺ mobilization and protein tyrosine phosphorylation. *J. Applied. Biomed.*, 5: 105-113.
- Fathi-Azad, F., A. Garjani, N. Maleki and Ranjdost, S. 2005. Study of the hypoglycemic activity of the hydroalcoholic extract of *Urtica dioica* in normal and diabetic rat. *Pharm. Sci.*, 2: 65-69.
- Fazeli, S.A., A.M. Gharravi, S. Ghafari, M. Jahanshahi and Golalipour, M.J. 2008. The granul density of the *Dentate gyrus* following administration of *Urtica dioica* extract on young diabetic rats. *Folia Morphol*, 67: 196-204.
- Fazeli, S.A., A.M. Gharravi, S. Ghafari, M. Jahanshahi and M.J. Golalipour, 2010. Effects of *Urtica dioica* extract on CA3 hippocampal pyramidal cell loss in young diabetic rats. *Neural Regeneration Res.*, 5: 901-905.
- Fulzele., S.V., P.M Satturwar, S.B. Joshi and A.K. Dorle, 2002. Studies on the immunomodulatory activity of a herbal formulation. *Ind. Drugs.*, 39: 364-367.
- Golalipour, M.J. and Khouri, V. 2007. The protective activity of *Urtica dioica* leaves on blood glucose concentration and beta-cells in streptozotocin-diabetic rats. *Pak. J. Biol. Sci.*, 10: 1200-1204.
- Golalipour, M.J., A.M. Gharravi, S. Ghafari and Afshar, M. 2007. Effect of *Urtica dioica* on morphometric indices of kidney in streptozotocin diabetic rats-a stereological study. *Pak. J. Biol. Sci.*, 10: 3875-3879.
- Golalipour, M.J., M.A. Gharravi, S. Ghafari and. Azarhoush, A 2009a. Protective effect of *Urtica dioica* renal morphometric and histologic alterations in streptozotocin diabetic rats. *J. Babol Univ. Med. Sci.*, 10: 14-22.
- Golalipour, M.J., S. Ghafari and M.H. Farsi, 2009b. Effect of *Urtica dioica* L. extract on quantitative morphometric alterations of liver parenchyma cells in STZ diabetic rats. *Int. J. Morphol.*, 27: 1339-1344.
- Golalipour, M.J., S. Ghafari, V. Kouri and A.A. Kestkar, 2010. Proliferation of the β -cells of pancreas in diabetic rats treated with *Urtica dioica*. *Int. J. Morphol.*, 28: 399-404.
- Grover, J. K S., Yadav and V. Vats. 2002. Medicinal plants of India with anti-diabetic potential. *Journal of Ethnopharmacology*, 81 (1): 81-10.
- Gupta A. and N.S. Kumar, 2008. Immunomodulatory and antioxidant activity of a polyherbal formulation. *Int. J. Pharmacol.*, 4: 287-291
- Hahraki, M.R., H. Mirshekari, A.R. Shahraki, E. Shahraki and K.H. Divnband, 2009. Effect of *Urtica dioica* boiling on serum glucose, insulin and lipids in fructose-fed male rats. *J. Ofogh-e-Danesh Gonabad Univ. Med. Sci. Health Services*, 14: 10-15.
- Hasani-Ranjbar, S., B. Larijani and M. Abdollahi, 2008. A systematic review of Iranian medicinal plants useful in *Diabetes mellitus*. *Arch. Med. Sci.*, 4, 3: 285-292.
- Hasani-Ranjbar, S., B. Larijani and Abdollahi, M. 2009. A systematic review of the potential herbal sources of future drugs effective in oxidant-related diseases. *Inflamm. Allergy Drug Targets*, 8: 2-10.
- Jachak, S.M. and A. Saklani, 2007. Challenges and opportunities in drug discovery from plants. *Curr. Sci.*, 92: 1251-1257.
- Kameswara Rao B, Renuka Sudarshan P, Rajasekhar MD, Nagaraju N, Appa R. C. 2003. Hypoglycemic activity of *Terminalia pallida* fruit in alloxan induced diabetic rats. *J. Ethnopharmacology*, 85: 169-172.
- Kar, A., Choudhary, B.K., Bandyopadhyay, N.G. 2003. Comparative evaluation of hypoglycaemic activity of some Indian medicinal plants in alloxan diabetic rats. *Journal of Ethnopharmacology*, 84, 105 – 108.

28. Khairandish, P., M. Mohraz, B. Farzamfar, M. Abdollahi and Shahhosseiny, M.H. 2009. Preclinical and phase 1 clinical safety of Setarud (IMOD™), a novel immunomodulator. *DARU*, 17: 148-156.
29. Khouri. V. and Golalipour, M.J. 2006. Chronic effect of the hydroalcoholic extract of *Urtica dioica* leaves in regeneration of β -cells of hyperglycemic STZ rats. *J. Med. Plants*, 5: 23-30.
30. Kumarappan, C T., Rao, T N. and Mandal, S. C. 2007. Polyphenolic extract of *Ichnocarpus frutescens* modifies hyperlipidemia status in diabetic rats. *Journal of Cell and Molecular Biology*, 6(2): 175-187.
31. Kusano, S., and Abe, H. 2000. Antidiabetic activity of white skinned sweet potato (*Ipomoea batatas* L.) in obese Zucker fatty rats. *Biol. Pharm.Bull*, 23: 23-26.
32. Lakka, H M., Laaksonen, D E., Lakka, T A., Niskanen, L K., Kumpusalo, E., Tuomilehto, J. and Salonen, J T. 2002. The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *Jama* 288:2709-2716.
33. Malihi, F., A. Hosseini-Tabatabaei, H. Esmaily, R. Khorasani, M. Baeeri and Abdollahi, M. 2009. Improvement of inflammatory and toxic stress biomarkers by silymarin in a murine model of type one diabetes mellitus. *Cent. Eur. J. Biol.*, 4: 369-380.
34. Marra G, Cotroneo P., Pitocco, D., Manto, A., Di Leo, M A., Ruotolo, V., Caputo, S., Giardina, B., Ghirlanda, G. and Santini, S A. 2002. Early increase of oxidative stress and reduced antioxidant defenses in patients with uncomplicated type 1 diabetes a case for gender difference. *Diabetes Care.*, 25:370-375.
35. Martin-Gallan, P., Carrascosa, A., Gussinye, M. and Dominguez, C. 2003. Biomarkers of diabetes-associated oxidative stress and antioxidant status in young diabetic patients with or without subclinical complications. *Free Radic Biol Med.*, 34:1563- 1574.
36. Milani, E., S. Nikfar, R. Khorasani, M.J. Zamani and M. Abdollahi, 2005. Reduction of diabetes-induced oxidative stress by phosphodiesterase inhibitors in rats. *Comparative Biochem. Physiol.*, 140 C: 251-255.
37. Mobaseri, M., A. Bahrami, N.A. Zarghami, A. Aliasgarzadeh and M. Rahmati et al., 2010. Effect of total extract of *Urtica dioica* on insulin and C-peptide secretion from rat (RIN5F) pancreatic β cells and glucose utilization by human muscle cells. *Iran. J. Endocrinol. Metab.*, 11: 721-727.
38. Mohseni-Salehi-Monfared, S.S., B. Larijani and. Abdollahi, M. 2009. Islet transplantation and antioxidant management: A systematic review. *World J. Gastroenterol.*, 15: 1153-1161.
39. Mohseni-Salehi-Monfared, S.S., E. Habibollahzadeh, H. Sadeghi, M. Baeeri, Abdollahi, M. 2010. Efficacy of Setarud (IMOD™), a novel electromagnetically-treated multi-herbal compound, in mouse immunogenic type-1 diabetes. *Arch. Med. Sci.*, 6: 663-669.
40. Momtaz, S. and Abdollahi, M. 2010. An update on pharmacology of satureja species; from antioxidant, antimicrobial, antidiabetes and anti-hyperlipidemic to reproductive stimulation. *Int. J. Pharmacol.*, 6: 454-461.
41. Mukherjee, P.K., Maiti, K., Mukherjee, K., Houghton, P.J. 2006. Leads from Indian medicinal plants with hypoglycemic potentials. *Journal of Ethnopharmacology*, 106, 1 – 28.
42. Petlevski, R., M. Hadzija, M. Slijepcevic and Juretic, D. 2001. Effect of antidiabetes herbal preparation on serum glucose and fructosamine in NOD mice. *J. Ethnopharmacol.*, 75: 181-184.
43. Pulok K. M., Kuntal M., Kakali M. and Peter Houghton. J. 2006. Leads from Indian medicinal plants with hypoglycemic potentials. *Journal of Ethnopharmacology*, 106, (1): 1-28.
44. Rahimi, R., S. Nikfar, B. Larijani and M. Abdollahi, 2005. A review on the role of antioxidants in the management of diabetes and its complications. *Biomed. Pharmacotherapy*. 59: 365-373.
45. Ramos, R.R., F. Alarcon-Aguilar, A. Lara-Lemus and J.L. Flores-Saenz, 1992. Hypoglycemic effect of plants used in Mexico as antidiabetics. *Arch. Med. Res.*, 23: 59-64.
46. Ravi, K., Rajasekaran, S., Subramanian, S., 2005. Antihyperlipidemic effect of *Eugenia jambolana* seed kernel on streptozotocin induced diabetes in rats. *Food and Chemical Toxicology*, 43, 1433 – 1439.
47. Rolo, A P. and Palmeira, C M. 2006. Diabetes and mitochondrial function role of hyperglycemia and oxidative stress. *Toxicol Appl Pharmacol.*, 212:167-178.
48. Said, O., S. Fulder, K. Khali, H. Azaizeh, E. Kassis and B. Saad, 2008. Maintaining a physiological blood glucose level with Glucoselevel, a combination of four anti-diabetes plants used in traditional Arab herbal medicine. Evidence-Based Complementary *Alternative Med.*, 5: 421-428.
49. Sarkhail, P., S. Rahmanipour, S. Fadyevatan, A. Mohammadirad and G. Dehghan., 2007. Antidiabetic effect of *Phlomis anisodonta*: Effects on hepatic cells lipid peroxidation and antioxidant enzymes in experimental diabetes. *Pharmacol. Res.*, 56: 261-266.
50. Shaw, J.E., R.A. Sicree and Zimmet, P.Z. 2010. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res. Clin. Pract.*, 87: 4-14.
51. Sicree, R., J. Shaw and Zimmet, P. 2007. Diabetes and Impaired Glucose Tolerance: Delice Gan Diabetes Atlas. 3rd Edn., International Diabetes Federation, Brussels, Belgium.
52. Sitasawad, S.L., Shewade, Y., Bhonde, R. 2000. Role of bittergourd fruit juice in STZ-induced diabetic state *in vivo* and *in vitro*. *J. Ethnopharmacol*, 73: 71-79.
53. Yeh, G.Y., Eisenberg, D.M., Kaptchuk, T.J., Phillips, R.S., 2003. Systematic Review of Herbs and Dietary Supplements for Glycemic Control in Diabetes. *Diabetes Care* 26, 1277 – 1294.
54. Varvarovska, J., Racek, J., Stozicky, F., Soucek, J., Trefil, L. and Pomahacova, R. 2003. Parameters of oxidative stress in children with Type-1 diabetes mellitus and their relatives. *J. Diabetes Complications*, 17:110-117.
55. Lan, H., Yanfei, Q., Xianglu, R., Jianmin, J., Qinglin, Y., Johji, Y., Michael, M. and Yuhao, L. 2011. The Ayurvedic medicine *Salacia oblonga* attenuates diabetic renal fibrosis in rats: suppression of angiotensin II/AT1 signaling. *Complement Alternat Med.*, 45:234-236.
