



Received on 20 February, 2012; received in revised form 02 April, 2012; accepted 23 May, 2012

CYTOTOXIC ACTIVITY OF METHANOL AND DICHLOROMETHANE EXTRACTS SOFT HALICONA SPECIES

Narasimharaju Alluri*, L.H. Thameemulansari and C. UmaMaheswara Reddy

Sri Ramachandra college of Pharmacy, Sri Ramachandra University, Chennai, Tamil Nadu, India

ABSTRACT

Keywords:

Marine sponges,
Cytotoxicity,
MTT assay,
HeLa cell lines

Correspondence to Author:

Narasimharaju Alluri

Sri Ramachandra college of Pharmacy, Sri
Ramachandra University, Chennai, Tamil
Nadu, India

The study was aimed to evaluate the anticancer activity of the Soft Halicono species on the HeLa cell line. The Soft Halicono species methanol and Dichloromethane extracts were tested for its inhibitory effect on HeLa cell line. The percentage viability of the cell line was carried out by using colorimetric method. The cytotoxicity of Soft *Halicono species* on HeLa cell was evaluated by MTT assay. Both the methanol and Dichloromethane extract has significant cytotoxic effect on HeLa cell line in concentration range between 0.1µg/ml to 100µg/ml by using MTT assay. R2 value of Soft Halicono species on HeLa cell of methanol extract was 0.8316 and for dichloromethane extract was 0.6699 and percentage cell inhibition for methanol extract 61.40% and Dichloromethane extract was 46.98%.

INTRODUCTION: Sponges (phylum: Porifera) are most primitive of the multicelled animals that have existed for 700–800 million years. Of the approximately 15,000 sponge species, most occur in marine environments. Only about 1% of the species inhabits freshwater¹. They inhabit every type of marine environment, from polar seas² to temperate and tropical waters^{3, 4} and are often more abundant and diverse in the tropics than stony and soft corals⁵. These animals are frequently exposed to intense predation and /or tissue infection by microorganisms^{6, 7}.

Sponges feed on bacteria⁸ and are constantly exposed to large populations of water-borne microbes, including opportunistic pathogens and fouling microorganisms. Despite these constant threats and a lack of the complex morphological and cellular defense mechanisms used by higher animals to combat bacterial pathogenesis^{9, 10}, sponges are highly successful members of the benthos and suffer few obvious bacterial infections. The sponge class Demospongiae is known to produce the largest number and diversity of secondary metabolites isolated from marine invertebrates¹¹.

Although the functions of these secondary metabolites are largely unknown, there is some evidence that they provide chemical defenses against predators¹². It has also been suggested that sponge secondary metabolites may provide defenses against fouling and infection^{13, 14, 15}; however, this possibility has not been adequately explored.

A HeLa cell is an immortal cell line used in medical research. The cell line was derived from Cervical cancer cells taken from Henrietta Lacks, who died from her cancer in 1951. Initially, the cell line was said to be named after a "Helen Lane" in order to preserve Lacks's anonymity. The objective of this study is to investigate cytotoxic activity of methanol and Dichloromethane extracts from sponges on HeLa cell lines using MTT assay.

MATERIALS AND METHODS

Collection of sponges: Soft Halicono species of marine demosponges were collected from the areas of the Rameswaram and Gulf of Mannar. Sponge samples were collected during September and October 2009. The sponges were collected at 1 to 27 m depth by

snorkeling and SCUBA. Sponge sample was immediately frozen after collection and maintained at -20°C prior to extraction.

Chemicals & Reagents: MTT assay kit (Sigma chemical Co, USA), Dichloromethane and Methanol (In-house), All the chemicals used for the biochemical analysis were procured from Sisco research laboratory, India. HeLa cell lines were collected from Department of Biotechnology, SriRamachandra University, Chennai.

Extraction: Sponge specimen was allowed to thaw, cut into small pieces, and then taken in to a graduated Cylinder containing 1000 ml of dichloromethane and in another cylinder with 1000ml of methanol and covered the cylinder with aluminum foil. And kept in cool and dry place for 15 days. Sponge tissue and solvent were transferred to capped containers and agitated for 24 h. After extraction, the sponge tissue was removed from the container and solvents squeezed from the tissue. The extract was collected by using vacuum separator. And the solution containing the extract at the volumetric concentration of the original tissue.

Cell lines and Culture condition: The cytotoxicity of the Dichloromethane and methanol extracts was tested against HeLa cell lines. Cells were grown in RPMI-1640 medium supplemented with 10% fetal bovine serum, 2 mM L-glutamine, 100 Ag/ml streptomycin, and 100 U/ml penicillin, and were incubated at 37°C in a humidified atmosphere of 95% air and 5% CO_2 .

Cytotoxic Bioassay: The proliferation rates of HeLa cells after treatment with essential oils were determined by the colorimetric 3-(4, 5-dimethyl-2-thiazolyl)-2, 5-diphenyl-2H-tetrazolium bromide (MTT) assay. The yellow compound MTT is reduced by mitochondrial dehydrogenases to the water-insoluble purple formazan, depending on the viability of cells. HeLa cells were plated in 96-well plates (105 cells/well for adherent cells or 0.5_105 cells/well for suspended cells in 100 Al of medium). After 24 h, the extract (2–125 Ag/ml) dissolved in distilled water was added to each well and incubated for 3 days (72 h). Control groups received the same amount of distilled water. Tumor cell growth was quantified by the ability of living cells to reduce the yellow dye 3-(4, 5-dimethyl-2-

thiazolyl)-2, 5-diphenyl-2H-tetrazolium bromide (MTT) to a purple formazan product.

At the end of the incubation, the plates were centrifuged and then, the medium was replaced by fresh medium (200 Al) containing 0.5 mg/ml MTT. Three hours later, the MTT formazan product was dissolved in 150 Al DMSO, and absorbance was measured using a multiplate reader (Spectra Count, Packard). Drug effect was quantified as the percentage of control absorbance of reduced dye at 550 nm.

RESULTS & DISCUSSION: Percentage cell viability of cell lines was carried out by using colorimetric method. The Cytotoxic activity of methanol extract and Dichloromethane extract is presented in **table 1 & 2** and in **figure 1 & 2**. It was found that the cell inhibition increases with increase in concentration steadily up to $100\mu\text{g/ml}$ on HeLa cell line. Percentage cell inhibition of methanol extract is 61.40% and Dichloromethane extract is 46.98%. And R2 value for methanol is 0.8316 and Dichloromethane is 0.6699.

TABLE 1: DETERMINATION OF CYTOTOXICITY FOR METHANOL EXTRACT BY MTT ASSAY

Conc. ($\mu\text{g/ml}$)	% Cell Inhibition	% Cell Survival
0.1	11.27	88.72
1	19.71	80.28
10	37.26	62.74
100	61.40	38.60

TABLE 2: DETERMINATION OF CYTOTOXICITY FOR DICHLOROMETHANE EXTRACT BY MTT ASSAY

Conc. ($\mu\text{g/ml}$)	% Cell Inhibition	% Cell Survival
0.1	19.20	80.79
1	28.81	71.19
10	38.71	61.28
100	46.98	53.02




FIG. 1: METHANOL EXTRACT OF SOFT HALICONA SPECIES FOR HELA CELL LINE BY MTT ASSAY




FIG. 2: DICHLOROMETHANE EXTRACT OF SOFT HALICONA SPECIES FOR HELA CELL LINE BY MTT ASSAY

Now over all study evaluate the sponge has potential activity on HeLa cell. So these extracts have considerable for anticancer activity.

CONCLUSION: In conclusion, our study can be considered as the first report on the Cytotoxic property of Soft Haliconna species. The results of the Cytotoxic activity against HeLa cell lines in this study are very promising with regards to possible antineoplastic chemotherapy and from a very sound basis for future research.

ACKNOWLEDGEMENT: The authors are grateful to the management of Sri Ramachandra University for providing the necessary infrastructure to carry out this work in a successful manner.

REFERENCES:

1. Belarbi E.H, A. C. Go´mez, Y. Chisti, F. G.Camacho, and E. M. Grima. 2003. Producing drugs from marine sponges. *Biotechnol. Adv.* 21: 585–598.
2. Dayton PK, Robilliard GA, Paine RT, Dayton LB (1974) Biological accommodation in the benthic community at McMurdo Sound, Antarctic. *Ecol Monogr* 44:105.
3. Reisswig HM (1973) Population dynamics of three Jamaican Demospongiae. *Bull Mar Sci* 23:191-226.
4. Wenner EL, Knott DM. Van Dolah RF. Burrell VG (1983) Invertebrate communities associated with hard bottom habitats in the South Atlantic Bight Estuar Coast Shelf *Sci* 17: 143-158.
5. Targett NM, Schmahl GP (1984) Chemical ecology and distribution of sponges in the Salt River Canyon, St. Croix, U.S.V.1 NOAA Technical Memorandum OAR NURP-1, Rockville, MD, p 361-369.
6. Faulkner, D.J., Harper, M.K., Haygood, M.G., Salomon, C.E., Schmidt, E.W., 2000. Symbiotic bacteria in sponges: sources of bioactive substances. In: Fusetani, N. (Ed.), *Drugs from the sea*. Karger, Basel, 1-5.
7. Newbold, R.W., Jensen, P.R., Fencal, W., Pawlik, J.R., 1999. Antimicrobial activity of Caribbean sponge extracts. *Aquatic Microbial Ecol.* 19, 279-284.
8. Bergquist RP, Bedford JJ (1978) The incidence of antibacterial activity in marine Demospongiae: systematic and geographic considerations. *Mar Biol* 46:215-221.
9. Simpson TL (1968) The structure and function of sponge cells: new criteria for taxonomy of poecilosclerid sponges (Demospongia). *Bull Peabody Mus Nat Hist* 25:1-141.
10. Pawlik JR, Chanas B, Toonen R, Fencal W (1995) Defenses of Caribbean sponges against predatory reef fish. I. Chemical deterrence. *Mar Ecol Prog Ser* 127:183-194.
11. Chanas B, Pawlik JR, Lindel T, Fencal W (1996) Chemical defense of the Caribbean sponge *Agelas clathrodes* (Schmidt). *J Exp Mar Biol Ecol* 208:185-196.
12. Amade P, Chevolut L (1982) Antimicrobial activities of marine sponges from French Polynesia and Brittany. *Mar Biol* 70: 223-228.
13. Thompson JE, Walker RP, Faulkner DJ (1985) Screening and bioassay for biologically-active substances from forty marine sponges species from San Diego California, U.S.A. *Mar Biol* 88:11-21
14. Faulkner DJ (1998) Marine natural products. *Nat Prod Rep* 15: 113-158.
15. Pawlik JR (1993) Marine invertebrate chemical defenses. *Chem Rev* 93:1911-1922
