



Received on 23 May, 2010; received in revised form 29 July, 2010; accepted 28 August, 2010

## ANTIRHEUMATOID ACTIVITY OF AQUEOUS EXTRACT OF *PIPER LONGUM* ON FREUNDS ADJUVANT-INDUCED ARTHRITIS IN RATS

Subhash R. Yende\*<sup>1</sup>, Vrushali D. Sannapuri<sup>2</sup>, Niraj S. Vyawahare<sup>3</sup> and Uday N. Harle<sup>1</sup>

Department of Pharmacology, Guru Nanak College of Pharmacy<sup>1</sup>, Nari, Nagpur (MS) India  
Department of Pharmacology, Allana College of Pharmacy<sup>2</sup>, Pune (MS) India  
Department of Pharmacology, AISSMS College of Pharmacy<sup>3</sup>, Pune (MS), India

### Keywords:

Anti rheumatoid activity,  
Piper longum,  
Sub-plenter,  
Complete Freund's adjuvant,  
Plethysmometer

### ABSTRACT

Aqueous extract of the fruits of the plant *Piper longum* was studied for their Anti rheumatoid activity in Freund's Adjuvant Induced Arthritis Rats with the dose of 200 and 400 mg/kg p. o. The administration of extract reported significant reduction in paw swelling on 4<sup>th</sup>, 8<sup>th</sup>, 14<sup>th</sup> and 21<sup>st</sup> day after sub-plenter administration of Complete Freund's adjuvant. The paw swelling was measured as a volume displacement using digital Plethysmometer. Furthermore, these results supported by radiographic analysis of affected knees of rats. From the results observed in the present investigation, it may be concluded that the aqueous extract of *P. longum* possesses potentially useful anti-arthritic activity in Complete Freund's Adjuvant model.

### Correspondence to Author:

#### Subhash R. Yende

Department of Pharmacology,  
Guru Nanak College of  
Pharmacy, Nari, Nagpur (MS)  
India

E-mail: subhashyende@gmail.com

**INTRODUCTION:** Rheumatoid arthritis (RA) is an autoimmune disease characterized by the chronic inflammation of synovial joints which results in severe bone destruction<sup>1</sup>. It affects approximately 5 million people worldwide of which 50 % are unable to work beyond 10 years of diagnosis<sup>2</sup>. A number of anti-inflammatory and ant rheumatic drugs used in treatment of RA have been developed over the past few decades, but still there is urgent need for more effective drugs with lower side effects<sup>3</sup>. Non-steroidal anti-inflammatory drugs (NSAIDs) are representing the mainstay of therapeutic management of arthritis. It is now recognized, however, these drugs, although effective at relieving the symptoms of the disease, may do little to improve the condition, and in some instances may contribute to its progression<sup>4</sup>. This further suggests need of effective alternative therapy. Recent studies shows that commonly used medicinal plants and herbal preparations are a good source for alternative therapy.

*Piper longum* L. (Piperaceae), popularly known in India as Pippali, is used as traditional medicine in Asia, especially in Indian medicine and in Pacific islands<sup>5</sup>. Various Piper species, widely distributed in the tropical and subtropical regions of the world, have been used as a spice and also as a folk medicine<sup>6,7</sup>. *Piper longum* is a reported as good remedy for treating gonorrhoea, menstrual pain, tuberculosis, respiratory tract infections, chronic gut related pain<sup>8</sup>. Other reported beneficial effects of *P. longum* include analgesic and diuretic effects, relaxation of muscle tension, and alleviation of anxiety<sup>9,10</sup>. The extract of crude drug of *P. longum* is frequently used as anti-inflammatory and insecticidal agents<sup>7,10</sup>. Piperine was the first amide isolated from piper species and was reported to display central nervous system depression, antipyretic, and anti-inflammatory activity<sup>6</sup>. Piperine is a potent inhibitor of the

mixed function oxygenase system and non-specific inhibition of cytochrome P450 isoenzymes<sup>11</sup>. The constituents of piper species have inhibitory activity on prostaglandin and leukotriene biosynthesis in vitro<sup>7</sup> and antifibrotic activity<sup>12</sup>. However, to our knowledge, no basic scientific studies on antiarthritic activity of aqueous extract of *P. longum* has been reported, hence we have evaluated the antiarthritic activity of the plant using Complete Freund's adjuvant (CFA) induced arthritis in rat which mimics symptoms of rheumatoid arthritis in humans. CFA induces the arthritis by heat killed cells of *Mycobacterium tuberculosis* and it mimics the latter's immunological and biochemical features wherein self antigens are recognized as foreign bodies.

#### **EXPERIMENTAL:**

**Animals:** Male albino rats of Wistar strain weighing around 200-250 g were used for the experiment purpose. The animals were housed in solid-bottomed polypropylene cages and acclimatized to animal house conditions. The rats were fed with commercial standard diet and water *ad libitum*. The experiments were designed and conducted in accordance with the guideline of CPCSEA and approved by Institutional Animal Ethical Committee (IAEC).

**Preparation of Extract:** The dried powder (1 kg) of seeds of *Piper longum* was macerated with the distilled water for 48 hours. The solvent from extract was removed completely by evaporation under vacuum using rotary evaporator and dried to get fine powder.

**Induction of Arthritis:** Arthritis was induced by a single sub-plantar injection of 0.1 ml of Complete Freund's adjuvant (CFA)(Sigma Chemicals, USA) containing 1.0 mg dry heat-killed *Mycobacterium tuberculosis* per milliliter sterile paraffin oil into a foot pad of the left hind paw of male rats<sup>13</sup>. The

swelling in hind paws were periodically examined in each paw from the ankle using digital Plethysmometer (Panlabs, India).

**Experimental Setup:** Animals were divided into four groups of six animals in each group as follows (**Table 1**).

**TABLE 1: GROUPING OF ANIMALS**

Group	Treatment	Dose (for 14 days)
Vehicle	Distilled water	1ml/kg (p. o.)
Diclofenac sodium	Diclofenac sodium	13.5 mg/kg (p. o.)
PL200	<i>P. longum</i> extract	200 mg/kg (p. o.)
PL400	<i>P. longum</i> extract	400 mg/kg (p. o.)

**Paw Volume Displacement:** On 0<sup>th</sup> day, the left hind paw volume of all rats as a volume displacement was measured using digital plethysmometer and on 1<sup>st</sup> day arthritis was induced in all rats using CFA. The aforementioned drug treatment was started on 1<sup>st</sup> day and continued for 14 days. The assessment of antiarthritic activity was carried out by measuring change in paw volume edema on 4<sup>th</sup>, 8<sup>th</sup>, 14<sup>th</sup> and 21<sup>st</sup> day after induction. The percent inhibition of paw volume of treated rats against vehicle treated rats was evaluated.

**TABLE 2: EFFECT OF AQUEOUS EXTRACT OF *P. LONGUM* AND CFA ON MEAN CHANGE IN POW VOLUME IN RAT**

Group	Mean changes in paw volume (Mean ± SEM)				% inhibition of paw swelling on 21 <sup>st</sup> day
	4 <sup>th</sup> day	8 <sup>th</sup> day	14 <sup>th</sup> day	21 <sup>st</sup> day	
Vehicle (only CFA treated)	3.29±0.03	2.83±0.15	2.51±0.04	2.46±0.08	0
Diclofenac sodium	2.56±0.12**	1.93±0.12**	1.64±0.09**	1.23±0.03**	55.00
PL200	2.42±0.07**	2.29±0.08*	2.11±0.10*	2.09±0.05**	15.04
PL400	2.64±0.05**	2.19±0.15*	1.78±0.05**	1.32±0.07**	46.32

The values are Means ± S.E.M. (n=6), \*  $P < 0.05$ , \*\*  $P < 0.01$ , compared with vehicle treated group (one-way ANOVA followed by Dunnett's test)

Swelling of soft tissue adjacent or around the joint is associated with arthritis and reduction of this swelling is an expected parameter or action


**Radiographic Analysis:** High resolution radiograph of the effected knees were taken using a Agfa machine Siemens system. The region between epiphysial growth plates of the femur and tibia (including the patella) was selected for analysis and representative rats hind paws from each group were photographed.

**Statistical Analysis:** Results were expressed as Mean±SEM. The data was analyzed using one way Analysis of Variance (ANOVA) followed by Dunnett's test.

**RESULTS:** From the results it was found that the aqueous extract of *P. longum* in doses of 400 mg/kg as well as 200 mg/kg shows significant ( $p < 0.01$ ) reduction in paw swelling. In 400 mg/kg treated group the peak effect ( $p < 0.01$ ) was observed on 4<sup>th</sup>, 14<sup>th</sup> and 21<sup>st</sup> day, while in 200 mg/kg treated group effect was observed on 4<sup>th</sup> and 24<sup>th</sup> day when compared with vehicle treated group. On 21<sup>st</sup> day after induction of arthritis the *P. longum* 400 mg/kg shows 46.32% inhibition in paw swelling and Diclofenac sodium (13.5 mg/kg) shows 55.00% inhibition in paw swelling (**Table 2**).

of drugs used in arthritis. The swelling of soft tissue was found to be more in the control animals group as compared to the Diclofenac sodium

treated and the *P. longum* 400 mg/kg treated group. The reduction of swelling was observed prominently in the diclofenac sodium treated group and followed by *P. longum* 400 mg/kg treated group and lastly in *P. longum* 200 mg/kg treated group (Fig. 1).



**FIG. 1: RADIOGRAPHIC ANALYSIS OF THE EFFECTED KNEES OF RATS**

**A**, Vehicle treated; **B**, Diclofenac Sodium treated; **C**, PL200 treated; **D**, PL400 treated

**DISCUSSION AND CONCLUSION:** CFA induced arthritis in rats is probably the best and the most widely used model since it has a close similarity to human rheumatoid disease. The determination of magnitude of swelling of the

injected hind paw is the most objective measurement that can be made to assess the antiarthritic activity. The change in the paw volumes in the 21 days after the inoculation of Freund's adjuvant into the hind paw is illustrated as above. In the antiarthritic groups the injected paws showed an immediate acute inflammatory response reaching a maximum on the fourth day<sup>14</sup>. The edema volume of the injected paw (primary lesion) developed rapidly and reached a peak in 18 hours after injection and persisted at this level for up to 90 hours<sup>15</sup>. A chronic phase of inflammation reached a plateau from the fourteenth to twenty first days<sup>14</sup>.

In adjuvant-induced arthritis model rats developed a chronic swelling in multiple joints with influence of inflammatory cells, erosion of joint cartilage and bone destruction and remodeling which have close similarities to human rheumatoid disease<sup>16</sup>. These inflammatory changes ultimately result in the complete destruction of joint integrity and functions in the affected animal. Also, the CFA administered rats showed soft tissue swelling around the ankle joints during the development of arthritis, which was considered as edema of the particular tissues<sup>17</sup>.

The determination of paw swelling is apparently simple, sensitive and quick procedure for evaluating the degree of inflammation and the therapeutic effects of drugs. Chronic inflammation involves the release of number of mediators like cytokines (IL-1 $\beta$  and TNF- $\alpha$ ) and interferon's. These mediators are responsible for the pain, destruction of bone and cartilage that can lead to severe disability<sup>18</sup>. TNF- $\alpha$  - induced free radical generation like H<sub>2</sub>O<sub>2</sub> activates inflammatory signalling pathway, including NF- $\kappa$ B in vascular cells<sup>19</sup>, and regulating the expression of cell adhesion molecules on endothelial cells and hence play an important role in various

inflammatory diseases<sup>20</sup>. It is reported that chloroform extracts of *P. longum* inhibited the TNF- $\alpha$ -induced expression of intercellular adhesion molecule-1 (ICAM-1) furthermore, extract inhibited the adherence of neutrophils to endothelial monolayer by inhibiting the TNF- $\alpha$ -induced expression of ICAM-1, Vascular cell adhesion molecule-1 (VCAM-1) and E-selectin in a dose- and time- dependent manner. Also, chloroform extracts of *P. longum* significantly inhibited the TNF- $\alpha$ -induced activation of NF-kB<sup>21</sup>.

However, in present study standard drug and aqueous extract of *P. longum* significantly suppressed the swelling of the paws. Reduction of paw swelling in the *P. longum* treated rats from the third week onwards may be due to immunological protection rendered by the plant extract. Piperine was the first amide isolated from piper species and was reported to display central nervous system depression, antipyretic, and anti-inflammatory activity<sup>6</sup>. Piperine is a potent inhibitor of the mixed function oxygenase system and non-specific inhibition of P450 isoenzymes<sup>11</sup>. Also piperine has potent immunomodulatory activity<sup>22</sup>. From the results observed in the current investigation, it may be concluded that the aqueous extract of *P. longum* possesses potentially useful anti-arthritic activity since it was active in Complete Freund's Adjuvant model.

## REFERENCES:

1. Suematsu, A. *et al*: Scientific basis for the efficacy of combined use of antirheumatic drugs against bone destruction in rheumatoid arthritis. *Modern Rheumatology*, 2007; 17: 17–23.
2. Hu Y, Green N, Gavrin LK, Janz K, Kaila N, Li HQ, Thomason JR, Cuzzo JW, Hall JP, Hsu S: Inhibition of Tpl2 kinase and TNF $\alpha$  production with quinoline-3-carbonitriles for the treatment of rheumatoid arthritis. *Bioorganic and Medicinal Chemistry Letter* 2006; 16: 6067–6072.
3. Badger AM, Lee JC: Advances in antiarthritic therapeutics. *Drug Discovery Today* 1997; 2: 427-435.
4. Collier S, Ghosh P: Evaluation of the effects of antiarthritic drugs on the secretion of proteoglycans by lapine chondrocytes using a novel assay procedure. *Annals of Rheumatoid Disease* 1989; 48: 372-381.
5. Shoba G, Joy D, Joseph T, Majeed M, Rajendran R Srinivas P: Influence of piperine on the pharmacokinetics of curcumin on animal and human volunteer. *Planta Medica* 1998; 64: 353-356.
6. Parmar VS, Jain SC, Bisht KS, Poonam RJ, Jha A, Tyagi OD, Prasad AK, Wengel J, Olsen CE, Boll PM: Phytochemistry of the genus *Piper*. *Phytochemistry* 1997; 46: 597-673.
7. Stohr JR, Xiao PG, Bauer R: Constituents of Chinese *Piper* species and their inhibitory activity on prostaglandin and leukotriene biosynthesis *in vitro*. *Journal of Ethnopharmacology* 2001; 75: 133-139.
8. Singh YN: Kava an overview. *Journal of Ethnopharmacology* 1992; 37: 18–45.
9. Singh YN, Blue Mental M: Kava, an overview. *Herbal Gram* 1997; 39: 34–55.
10. Ghoshal S, Lakshmi V: Potential antiamebic property of the roots of *Piper longum* Linn. *Phytotherapeutic Research* 2002; 16: 689-691.
11. Atal CK, Dubey RK, Singh JJ: Biochemical basis of enhanced drug bioavailability by piperine, evidence that piperine is a potent inhibitor of drug metabolism. *Journal of Experimental Therapeutics* 1985; 232: 258-262.
12. Christinaa AJM, Saraswathya GR, Heison SJ, Kothaia R, Chidambaranathana N, Nalinia G Therasal RL: Inhibition of CCl<sub>4</sub> induced liver fibrosis by *Piper longum* Linn. *Phytomedicine* 2006; 13: 196–198.
13. Mizushima, Y, Tsukada W, Akimoto T: A modification of rat adjuvant arthritis for testing anti-rheumatic drugs. *Journal of Pharmacy and Pharmacology* 1972; 24: 781–785.
14. Fahim AT, Fattah AA, Agha AM, Gad MZ: Effect of pumpkin seed oil on the level of free radicals scavengers induced during adjuvant arthritis in rats. *Pharmacological Research* 1995; 31: 73-79.
15. Sofia RD, Knobloch LC, and Vassar HB: Inhibition of primary lesion of adjuvant induced polyarthritis in rat (18-hour arthritis test) for specific detection of clinically effective antiarthritic drugs. *Journal of Pharmacology and Experimental Therapeutics* 1975; 193: 918-931.
16. Singh S, Majumdar DK: Effect of fixed oil of *Ocimum sanctum* against experimentally induced arthritis and joint edema in laboratory animals. *International Journal of Pharmacognosy*; 34: 218-222.
17. Carl MP: Experimental joint disease observations on adjuvant-induced arthritis. *Journal of Chronic Disease* 1963; 16: 863-874.
18. Eric GB, Lawrence JL: Rheumatoid Arthritis and its therapy. The textbook of therapeutics drug and disease management. Williams and Wilkins Company, Baltimore, 16th Ed; 1996: 579-595.
19. Garg AK Agrawal BB: Reactive oxygen intermediates in TNF signaling. *Molecular Immunology* 2002; 39: 509-517.
20. Rahman I, MacNee W: Role of transcription factors in inflammatory lung diseases. *Thorax* 1998; 53: 601–612.
21. Singh N, Kumar S, Singh P, Raj HG, Prasad AK, Parmar VS, Ghosh B: *Piper longum* Linn. Extract inhibits TNF-alpha-induced expression of cell adhesion molecules by inhibiting NF-kappa B activation and microsomal lipid peroxidation. *Phytomedicine* 2008; 15(4): 284-291.
22. Sunila ES, Kuttan G: Immunomodulatory and antitumor activity of *Piper longum* Linn. and piperine. *Journal of Ethnopharmacology* 2004; 90: 339-346.