



Received on 22 December, 2015; received in revised form, 08 April, 2016; accepted, 24 April, 2016; published 01 May, 2016

## TRADITIONAL INDIAN MEDICINAL PLANTS WITH POTENTIAL WOUND HEALING ACTIVITY: A REVIEW

Sapna Saini, Anju Dhiman and Sanju Nanda\*

Department of Pharmaceutical Sciences, M.D. University, Rohtak-124001, Haryana, India

### Key words:

Hemostasis, Indigenous medicine, Inflammation, Traditional medicine and Wound healing.

### Correspondence to Author:

**Dr. (Mrs.) Sanju Nanda**

Associate Professor, (Pharmaceutics)  
Department of Pharmaceutical  
Sciences, M.D. University, Rohtak-  
124001, Haryana, India

**E-mail:** sn\_mdu@rediffmail.com

**ABSTRACT:** In last few decades, there has been a great progress in understanding the biochemical and cellular events of normal wound healing. Healing of wounds, either accidental or surgical interventions, involves complex activities of blood cells, tissues, soluble mediators, cytokines and several growth factors. This increased cellular activity of damaged tissue enhances metabolic demands and active drug therapy. The main objective of treating a wound is to either shorten the time required for healing process or to minimize the undue effects. Plants due to presence of various valuable active phytoconstituents have immense potential for management and treatment of wounds over the years. All the Traditional systems of medicine, Ayurveda, Siddha and Unani describe applications of drugs of plant, mineral and animal origin to treat and heal wounds. Herbal drugs induce healing and regeneration of lost tissue by number of mechanisms. Due to their traditional applicability, affordability and safety plants gained a reputed position in the world of wound management and repair but scientific evidence for their wound healing potentials are very few. The present review is an attempt to highlight various Indian ethno-medicinal plants which are to be scientifically proved for the treatment of wounds. Beside this review also emphasis on normal wound healing process, pharmacological activities and role of plants in wound management and parameters used to assess wound healing.

**INTRODUCTION:** Skin covers the external surface of the body. Skin contributes in thermoregulation, serve as water repellent, and synthesize a number of useful compounds like vitamin D and most importantly it act as a protective barrier between the external environment and internal tissue<sup>1</sup>. According to medical dictionary; breaching of skin or underlying tissue due to any accident, act of violence or surgery is called as wound<sup>2</sup>. Following an injury, the skin has a tremendous capacity to heal.

When skin get injured, the repair process entails removal of the damaged tissue and laying down of a new extracellular matrix (ECM) over which epidermal continuity can be reestablished<sup>3</sup>. Wound repair must occur in a physiologic environment to promote tissue repair and regeneration. However, several clinically significant factors are known to impede wound healing including hypoxia, infection, tumors, metabolic disorders such as diabetes mellitus, the presence of debris and necrotic tissue, certain medications, and a diet deficiency in protein, vitamin, and minerals etc<sup>4</sup>.

Current estimates indicate that approximately 6 million people suffer from chronic wounds worldwide<sup>5</sup>. There are very few Indian studies on the epidemiology of wounds. The prevalence of

<p><b>QUICK RESPONSE CODE</b></p> 	<p><b>DOI:</b> 10.13040/IJPSR.0975-8232.7(5).1809-19</p>
<p>Article can be accessed online on: <a href="http://www.ijpsr.com">www.ijpsr.com</a></p>	
<p>DOI link: <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.7(5).1809-19">http://dx.doi.org/10.13040/IJPSR.0975-8232.7(5).1809-19</a></p>	

wounds in the population studied was 15.03 per 1000. The prevalence of chronic wounds in the community was reported as 4.5 per 1000 population whereas that of acute wounds was nearly doubled at 10.5 per 1000 population<sup>6</sup>. Today, the principles of topical wound therapy involve elimination of necrotic tissue, control of bacterial loads, management of wound exudates, maintenance of open proliferative wound edges, and provision of a moist and protected wound surface<sup>7</sup>. Medical treatment of wound includes administration of drugs either locally (topical) or systemically (oral or parenteral) or both in an attempt to aid wound repair<sup>8</sup>. Antimicrobial dressings including disinfectants, antiseptics and antibiotics are applied topically to exert a broad spectrum of non-selective antibacterial action<sup>9</sup>.

The most commonly used antiseptic products in clinical practice include povidone-iodine complex, chlorhexidine, alcohol, triclosan, hydrogen peroxide, boric acid, silver nitrate, silver sulfadiazine and sodium hypochlorite<sup>10</sup>. In clinical practices, topical antibiotics are commonly prescribed in wounds, cuts and burns for their potential use in localized cutaneous infections. However frequent and overuse of topical antibiotics can cause problems like development of resistant organisms<sup>11</sup>.


Therefore attention should be directed towards discovering an agent that accelerate wound healing either when it is progressing normally, or when it is suppressed by various agents like corticosteroids, anti-neoplastics, or non-steroidal anti-inflammatory agents<sup>12</sup>. Natural product research is often based on ethno-botanical information and many of the drugs used today were developed from medicinal plants employed in indigenous societies<sup>13</sup>.

This article provides a general description on normal wound healing process. Plant phytoconstituents that play a role in wound management is also explained in this review. It also provides knowledge regarding the parameters used to assess wound healing and also enlist various traditional Indian medicinal plant used for wound healing.

### Normal wound healing process:

Wound is defined as loss or breaking of cellular and anatomic or functional continuity of living tissue due to physical, chemical, thermal, microbial or immunological exploitation to the tissues<sup>14, 15</sup>. Wound healing disorders present a serious clinical problem and are likely to increase when they are associated with diseases such as diabetes, hypertension, and obesity. Additionally, increasing life expectancies will cause more people to face such disorders and therefore wounds become an unavoidable event of life<sup>16</sup>. Wound healing is complex multistep physiological process that involves multitude of cells and events. It is a dynamic, interactive process which involves soluble mediators, blood cells, extracellular matrix and parenchymal cells. It is dependent on a number of inter-related factors<sup>17-19</sup>.

In normal conditions, the epidermis (outermost layer of the skin) and dermis (inner layer) forms a protective barrier against the external environment. If this protective barrier is broken due to any kind of injury, a set of complex biochemical events of wound healing takes place in a highly sequenced manner to repair the damage<sup>20</sup>. Normal wound healing is mostly characterized by four continuous, overlapping and precisely planned phases: hemostasis phase (formation of blood clot), inflammation phase, proliferation or granulation phase, and remodeling or maturation phase (formation of scar) shown in **Fig. 1**<sup>21, 22</sup>.



**FIG. 1: NORMAL WOUND HEALING PHASES**

Optimal wound healing involves the following the events: (1) rapid hemostasis; (2) appropriate inflammation; (3) mesenchymal cell differentiation, proliferation, and migration to the wound site; (4)

suitable angiogenesis; (5) prompt re-epithelialization (re-growth of epithelial tissue over the wound surface); and (6) proper synthesis, cross-linking, and alignment of collagen to provide

strength to the healing tissue. Various cellular and bio-physiological events occur during wound healing process are shown in **Table 1**<sup>23</sup>.

**TABLE 1: NORMAL WOUND HEALING PROCESS**

Phases	Cellular and Bio-physiological events
Hemostasis	1. Vascular constriction 2. Platelets aggregation, degranulation and fibrin formation (Thrombus)
Inflammation	1. Neutrophil infiltration 2. Monocyte infiltration and differentiation to macrophages 3. Lymphocyte infiltration
Proliferation	1. Re-epithelialization 2. Angiogenesis 3. Collagen synthesis 4. Extracellular matrix formation
Remodeling	1. Collagen remodeling 2. Vascular maturation and regression

#### **Role of traditional medicines in wound healing:**

It has been estimated by WHO that at least 80% of the world population, mainly in the developing countries, still dependent on herbal medicines for their primary health care needs. Use of traditional medicine is based on its accessibility, affordability and its firm embedment within the faith systems of people<sup>24, 25</sup>. Traditional knowledge is a valuable asset for any country as it plays a vital role in making the nation more progressive and transforming its society<sup>26</sup>. Traditional medicine is a collection of therapeutic experience of various physicians on indigenous/traditional system of medicine<sup>27</sup>. According to WHO, "Traditional medicine refers to health practices, approaches, knowledge and beliefs incorporating plant, animal and mineral based medicines, spiritual therapies, manual techniques and exercises, applied singularly or in combination to treat, diagnose and prevent illnesses or maintain well-being."

Traditional medicine is called in various ways such as alternative medicine, complementary medicine, natural medicine, herbal medicine, phyto-medicine, non-conventional medicine, indigenous medicine, folk medicine, ethno medicine etc<sup>28</sup>. People in the rural places where modern health care services remains inaccessible are still use their knowledge, skill and practices on various plant products for treating various ailments<sup>29</sup>. It has been estimated that 70% of wound healing Ayurvedic drugs are

plant based, 20% of mineral based and remaining 10% consists of animal products as their base

material<sup>30</sup>. Traditional wound management involves disinfection, debridement and providing a moist environment to encourage the establishment of a suitable environment for the natural healing process<sup>31</sup>.

#### **Pharmacological activities of plants supporting wound healing:**

##### **Anti-inflammatory activity:**

Inflammation phase in normal wound healing process is essential, because it generates neutrophils that are responsible for microbial clearance in the region of the wound (phagocytosis) and also playing a role in antigen presentation. In this phase production of inflammatory cytokines and growth factors also occur. Any pathological process that interferes with this self-limited physiological process can result in a non healing wound because of net destruction of soluble growth factors and matrix elements<sup>32, 33</sup>. Various traditional texts and animal studies indicate the anti-inflammatory effect of plant extract thus can be used as wound healer drugs<sup>34, 35</sup>.

##### **Antioxidant activity:**

Plant produces various anti oxidative compounds to counteract reactive oxygen species (ROS) in order to survive. ROS include free radicals such as superoxide anion radicals, hydroxyl radicals and non free-radical species such as hydrogen peroxide

etc are exacerbating factors in cellular injury and aging process<sup>36</sup>. Various antioxidant natural remedies derived from plant such as polysaccharides from *Angelica sinensis*, *Aloe vera* gel, *Eucommia ulmoides* Oliver leaf extract possess wound healing by scavenging free radical and promoting collagen synthesis<sup>37</sup>.

#### Antimicrobial activity:

Microorganisms are believed to play a significant role in impaired healing of chronic wounds and the development of infection-related complications<sup>38</sup>. Bacterial infection can cause sepsis which impedes the repair process<sup>39</sup>. A number of polyherbal preparations containing antimicrobial action have been scientifically proved to possess wound healing activity e.g. polyherbal gel formulated from *Terminalia arjuna*, *Centella asiatica* and *Curcuma longa* reported to possess antimicrobial and wound healing activities<sup>40</sup>.

#### Analgesic activity:

Pain is a common experience for patients with wounds. At a physiological level, wound pain

arises from tissue damage (nociceptive pain) or from dysfunction of the nervous system (neuropathic pain)<sup>41</sup>. Pain may cause delay in wound healing process by dysregulating neuroendocrine and immune functions which play critical role in wound repair mechanism<sup>42</sup>. Thus, herbal preparations possessing analgesic and anti-inflammatory activities can be employed for pain management of wounds.

#### Role of phytoconstituents in wound healing activity:

Phytoconstituents present in the plant extracts may interfere with one or more phases of the wound healing process in a positive manner in proper sequence and at the right time frame to show improved efficacy as shown in **Table 2**<sup>43</sup>. Various substances possessing wound healing activity has been isolated from plants e.g. tannins from *Terminalia arjuna*, polysaccharides from *Opuntia ficus-indica*, asiaticoside, asiatic acid and madecassic acid from *Centella asiatica* and curcumin from *Curcuma longa* etc<sup>44</sup>.

**TABLE 2: PHYTOCONSTITUENTS AND THEIR ROLE IN WOUND HEALING**

Phytoconstituents	Role
Tannins	Promote wound healing due to their astringent and antimicrobial property, act as free radical scavengers
Flavanoids	Possess antioxidants and free radical scavenging effects, astringent and antimicrobial activity, improve vascularity
Saponins	Antioxidant and antimicrobial activity.
Sterols and Polyphenols	Free radical scavenging and antioxidant activity
Tri-terpenoids	Astringent and antimicrobial activity

#### Parameters used to assess wound healing activity:

For the study of wound healing activity many *in-vitro* and *in-vivo* models have been used. *In-vitro* study of wound healing can be done by keratinocytes assay, wound healing migration assay and cell proliferation assay or fibroblasts assay<sup>45</sup>. *In-vivo* models such as incision models, excision models, dead space models and burn models can be performed. Estimation of tensile strength of skin, hydroxyproline content and measurement of wound area, percentage of contraction and period of epithelialization are some more preferable

parameters used for evaluation of wound healing activity of any compound<sup>46-48</sup>.

#### Medicinal plants possessing wound healing activity:

India is eighth major centre of origin and diversification of domesticated taxa due to its glorious part of traditional medicinal system and is one of the world's twelve mega diversity countries<sup>49</sup>. Therefore, it has a rich traditional knowledge for management and treatment of wounds since thousands of years. Indian religious book 'Ramayana' described the use of plant based medicine from Himalayas for treating Lord Rama's Brother Lakshman in the battlefield of Lanka.

Herbal products are in huge demand in developed countries too due to undue side effects of synthetic products and comparatively safe parameters of herbal products. Various scientifically evident plants used by local communities of India for management of wound healing are given as:

***Acalypha indica* Linn. (Euphorbiaceae):**

*Acalypha indica* L. is a weed plant widely distributed throughout the plains of India. It has been reported to be useful in treating pneumonia, asthma, rheumatism and several other skin ailments. *A. indica* leaves and flowers showed antioxidant activity due to presence of flavanoids, kaempherol, glycosides, mauritianin, ciltoria and nictiflorin. The ethanolic extract of dried leaves of *A. indica* has been used for bedsores and wounds treatment<sup>50,51</sup>. Raja *et al.* showed the anti-bacterial activity of aqueous and acetone extract of leaves of plant<sup>52</sup>.

***Aloe vera* (Liliaceae):**

*Aloe vera* is indigenous medicinal plant found throughout India. There are over 100 active constituent are found in *A. vera* plant which possesses astringent, haemostatic, antidiabetic, antiulcer, antiseptic, antibacterial, anti-inflammatory, antioxidant, anticancer, antidiarrhoeal, and wound healing properties<sup>53</sup>. The mucilage content present in leaves of this plant that is also called as aloe gel is used for various cosmetics and medical applications<sup>54</sup>. *A. vera* leaves pulp was found to have better and faster wound healing effect than standard drug Povidone Iodine ointment (5 % w/w) on excision wound model<sup>55</sup>.

***Asparagus racemosus* (Liliaceae):**

*Asparagus racemosus* is an indigenous medicinal climber found all over India; especially in Northern India. Plant value is known for its saponin content which is a precursor of many pharmacologically active steroids<sup>56</sup>. Prabhat *et al.* studied the wound healing activity of aqueous root extract of *A. racemosus* on dead space model on albino rat. The extract showed a facilitating effect on wound healing by a significant increase in hydroxyproline, hexosamine and hexuronic acid in granulating tissue<sup>57</sup>.

***Azadirachta indica* (Meliaceae):**

*Azadirachta indica* commonly known as Neem is a well known versatile plant of India for more than 2000 years possess wide spectrum of medicinal activities. Nimbidine, nimbin, sodium nimbidate and nimbolide main active constituent extracted from oil of its seed kernels showed antibacterial, anti-pyretic and anti-inflammatory properties<sup>58</sup>. Purohit *et al.* evaluated the wound healing activity of ethanolic extract of leaves of *A. indica* in male albino rats<sup>59</sup>.

***Bryophyllum pinnatum* (Lam.) Kurz. (Crassulaceae):**

*Bryophyllum pinnatum* also known as *Bryophyllum calycinum* is a perennial herb used as folkloric medicine in India and many other countries. The main phytochemical constituents present in plant are phenols, flavanoids, alkaloids, tannins, glycosides, saponins, coumarins and sitosterols<sup>60</sup>. Leaves of plant possess disinfectant and antibacterial properties used for boils, swelling, insect-bite, burns and wounds<sup>62</sup>.

***Calotropis gigantea* (L.) R. Br. (Asclepiadaceae):**

In ancient Ayurvedic medicine the plant *Calotropis gigantea* is known as "Sweta Arka." It is a native to India, China, Malaysia and in many other countries. The main active phytoconstituents isolated from plant are flavonoids, triterpenoids, phenolic compounds, tannins, alkaloids, steroids, glycosides, saponins, terpenes, enzymes, alcohol, resin, fatty acids and esters of calotropeols<sup>63</sup>. Deshmukh *et al.* evaluated the wound healing activity of topical application of *Calotropis gigantea* in excision wound model. Percentage of wound contraction was found to be increased due to elevating hydroxyproline content in incision wound and dead space wound model<sup>64</sup>.

***Centella asiatica* (Linn.) Urban (Apiaceae):**

*Centella asiatica* or Indain Pennywort is distributed throughout the plains of India. It is known as Brahmi in Unani medicine and Mandookaparni in Ayurvedic medicine system<sup>65</sup>. The main therapeutic active biomarker compounds reported in plant are saponins containing triterpens namely, asiatic acid, madecassic acid, asiaticoside and madecassoside<sup>66</sup>. It has been reported that 1% *C.*

*asiatica* extract cream improves chronic wound healing<sup>67</sup>. Shukla *et al.* studied the wound healing activity of topical application of asiaticoside in normal as well as in diabetic animals. The rate of wound healing in normal animals significantly enhanced due to increase in collagen synthesis and tensile strength of the wound tissues<sup>68</sup>. Liu investigated the effect of madecassoside, on wound healing involved several mechanisms including antioxidative activity, collagen synthesis and angiogenesis affect<sup>69</sup>.

#### ***Curcuma longa* Linn. (Zingiberaceae):**

*Curcuma longa* is an oldest spice and very useful herb of Ayurvedic medicine. It has been used throughout India since thousands of years. It is also referred to as 'Indian saffron' and turmeric<sup>70</sup>. Major components of turmeric are named curcuminoids, which include mainly curcumin (diferuloyl methane), demethoxycurcumin, and bisdemethoxycurcumin. Curcumin, a potent antioxidant is believed to be the most important bioactive component of the herb turmeric and posses the properties like antioxidant, anti-inflammatory, anti-platelet, cholesterol lowering, antibacterial and anti-fungal effects<sup>71</sup>. Purohit *et al.* investigated that ethanolic extract of *C. longa* rhizomes have better and faster wound healing effect than standard drug Povidone Iodine ointment on excision wound model<sup>72</sup>.

#### ***Elephantopus scaber* Linn. (Asteraceae):**

*E. scaber* is a weed plant that grows under shady places. Main bioactives compounds isolated from ethanolic extract of plant are sesquiterpene lactones such as elescaberin, isodeoxyelephantopin, deoxyelephantopin, isoscabertopin and scabertopin<sup>73</sup>. Singh *et al.* evaluated the wound healing potential of aqueous leaf extract and deoxyelephantopin, a compound isolated from ethanolic extract of leaf. Wound healing found to be more effective in animals treated with deoxyelephantopin<sup>74</sup>.

#### ***Euphorbia neriifolia* Linn. (Euphorbiaceae):**

*Euphorbia neriifolia* L. grows luxuriously in north, central and south India<sup>75</sup>. Bigonia *et al.* evaluated the wound healing activity of *E. neriifolia* leaf on excision and dead space wound model. *E. neriifolia* augmented wound contraction and epithelisation by

increased protein and hydroxyproline content<sup>76</sup>. Gour *et al.* evaluated the anti-inflammatory and analgesic effect of hydroalcoholic leaf extract plant<sup>77</sup>.

#### ***Ficus racemosa* Linn. (Moraceae):**

*Ficus racemosa* Linn. large deciduous tree distributed all over India especially in north India<sup>78</sup>. Murti *et al.* evaluated the wound healing activity of aqueous and ethanolic extract of roots of *F. racemosa* on incision and excision wound model. Aqueous extract of root increased the percentage closure due to enhanced epithelialization and collagen synthesis<sup>79</sup>.

#### ***Jatropha curcas* Linn. (Euphorbiaceae):**

*Jatropha curcas* Linn. commonly known as physic nut or jungali arandi. Plant extract is used in the treatment of allergies, burns, cuts, wound inflammation, leprosy, leucoderma and smallpox<sup>80</sup>. Nwala *et al.* formulated an herbal ointment of *J. curcas* leaf extract and evaluated its wound healing activity. Ointment was found to be good wound care product<sup>81</sup>. Similarly, Esimone *et al.* showed the potential wound healing activity of methanolic leaf extract of *J. curcas* on excision wound model<sup>82</sup>.

#### ***Lantana camara* Linn. (Verbenaceae):**

*Lantana camara* is a flowering ornamental plant found throughout in India. It is also known as lantana, wild sage, Surinam tea plant or Spanish flag. In India tea of leaves is used as remedy against cough and pounded leaves are applied as a paste to cuts, ulcers, swellings and wound healing<sup>83, 84</sup>. Nayak *et al.* found the better burn wound healing activity of the leaf extract of *L. camara* in rats as compare to control group<sup>85</sup>.

#### ***Ocimum sanctum* Linn. (Labiaceae):**

*Ocimum sanctum* Linn. a commonly known as 'Tulsi' is widely distributed plant throughout India and different parts of the world. It possesses anti-inflammatory, analgesic, immunostimulatory, free radical scavenging and antimicrobial activity. The free radical scavenging activity of plant flavonoids help in the healing of wounds. Asha *et al.* evaluated the topical wound healing activity of aqueous extract of leaves of *O. sanctum*<sup>86</sup>. Goel *et al.* studied the wound healing activity of cold aqueous extract of *O. sanctum* along with its effect on tumor

necrosis factor- $\alpha$  (TNF- $\alpha$ ) by using excision wound model in *Wistar albino* rats. Wound healing was found to be faster in *O. sanctum* extract treated rats as compare to control group due to elevating TNF-

$\alpha$  production<sup>87</sup>. Some other plants used traditionally and also scientifically validated are tabulated in **Table 3**.

**TABLE 3: SCIENTIFICALLY EVIDENT PLANTS USED FOR WOUND HEALING ACTIVITY**

Botanical name/Common name (Family)	Part used	Extract used	Wound healing model used
<i>Acorus calamus</i> /Bach plant (Acoraceae)	Green leaves	Ethanollic	Excision and incision <sup>88</sup>
<i>Allium sativum</i> L./Garlic (Liliaceae)	Bulb	Aqueous and Ethanollic	Excision, incision and dead space <sup>89</sup>
<i>Adhatoda vasica</i> N./Adalsa (Acanthaceae)	Leaves	Methanollic, chloroform and diethyl ether	Excision <sup>90</sup>
<i>Alternanthera brasiliiana</i> Kuntz./ Gudaari saag (Amaranthaceae)	Leaves	Methanollic	Excision, incision and chorioallantoic membrane <sup>91</sup>
<i>Andrographis paniculata</i> /Kalmegh (Acanthaceae)	Whole plant	Pet ether and Ethanollic	Excision <sup>92</sup>
<i>Areca catechu</i> L./Betel nut (Arecaceae)	Areca nut	Ethanollic	Burn wound <sup>93</sup>
<i>Butea monosperma</i> Lam. /Palash (Fabaceae)	Stem bark	Ethanollic	Excision, incision and dead space <sup>94</sup>
<i>Cassia fistula</i> L./Amaltas (Caesalpinioideae)	Leaves	Ethanollic	Incision <sup>95</sup>
<i>Catharanthus roseus</i> L./Vinca rosea (Apocyanaceae)	Flower	Ethanollic	Excision, incision and dead space <sup>96</sup>
<i>Carica papaya</i> L./Papita (Caricaceae)	Fruit latex	Dried latex	Burn wound <sup>97</sup>
<i>Cordia dichotoma</i> Forst. F./Lasura (Boraginaceae)	Fruit	Ethanol, petroleum ether, butanol, ethyl acetate	Excision, incision and dead space <sup>98</sup>
<i>Desmodium triquetrum</i> L./Balolia (Fabaceae)	Leaves	Ethanol	Excision, incision and dead space <sup>99</sup>
<i>Embelia ribes</i> Burm. /Vayavidanga (Myrsinaceae)	Leaves	Ethanol	Excision, incision and dead space <sup>100</sup>
<i>Ficus religiosa</i> /Peepal (Moraceae)	Leaves	Hydroalcoholic	Excision and incision <sup>101</sup>
<i>Gentian lutea</i> Linn/Gentian (Gentianaceae)	Rhizomes	Alcoholic and petroleum ether	Excision and Incision <sup>102</sup>
<i>Glycyrrhiza glabra</i> L./mulhatti (Leguminosae)	Root	Aqueous	Excision <sup>103</sup>
<i>Gymnema sylvestri</i> /gurmar (Asclepiadaceae)	Leaves, stem and flower	Aqueous	Excision <sup>104</sup>
<i>Heliotropium indicum</i> /Siriyaari (Boraginaceae)	Whole plant	n-butanol	<i>In-vitro</i> model <sup>105</sup>
<i>Indigofera enneaphylla</i> L./Latahai (Leguminosae)	Whole plant	Pet ether, ethyl acetate and ethanol	Excision and incision <sup>106</sup>
<i>Jasminum auriculatum</i> /juhi (oleaceae)	Whole plant	Pet ether, ethyl acetate and ethanol	Excision and incision <sup>106</sup>
<i>Kaempferia galangal</i> / chandramula (Zingiberaceae)	Juice of leaves	Juice of leaves	Excision <sup>107</sup>
<i>Lycopodium serratum</i> Thunb./ club moss (Huperziaceae)	Rhizomes	Alcoholic	Excision, incision and dead space <sup>108</sup>
<i>Mimosa pudica</i> L./chuihui (Fabaceae)	Whole plant	Ethanollic	Excision, incision and dead space <sup>109</sup>
<i>Mimusops elengi</i> L./mulsari, bakula (Sapotaceae)	Leaves	Ethanollic	Excision and burn wound <sup>110</sup>
<i>Michelia champaca</i> /Champaca	Stem bark	Methanollic	Excision, incision

(Magnoliaceae) <i>Mirabilis jalapa</i> L./ Gulabakshi	Flower	Ethanollic	and dead space <sup>111</sup> Excision and incision <sup>112</sup>
(Nyctaginaceae) <i>Piper betle</i> L./Pan (Piperaceae)	Tuberous root	Hydromethanolic	Excision <sup>113</sup>
<i>Quercus infectoria</i> G. Olivier /Majuphal (Fagaceae)	Leaves and stem bark	Ethanollic	Excision <sup>114</sup>
<i>Rubia cordifolia</i> /Manjistha (Rubiaceae)	Leaves	Ethanollic	Excision, incision and dead space <sup>115</sup>
<i>Sesamum indicum</i> L./Sesame (Pedaliaceae)	Root	Ethanollic	Excision <sup>116</sup>
<i>Sida acuta</i> /Jangali methi (Malvaceae)	Sesamum seed and oil	--	Excision, incision, dead space and burn model <sup>117</sup>
<i>Terminalia chebula</i> Retz./ Haritaki, harad (Combretaceae)	Whole plant	Methanolic	Excision and incision <sup>118</sup>
<i>Tridax procumbens</i> L./Ghamra (Asteraceae)	Fruit	Ethanollic	Excision and incision <sup>119</sup>
<i>Weddelia chinensis</i> Merrill/ Bhringraj (Asteraceae)	Whole plant	Ethanollic	Excision and incision <sup>120</sup>
<i>Ziziphus nummularia</i> L./ Jharber, Beri (Rhamnaceae)	Leaves	Ethanollic	Excision, incision and dead space <sup>121</sup>
	Leaves	Ethanollic	Excision <sup>122</sup>

**CONCLUSION:** Utilization of plants for medicinal purposes in India has been documented long back in ancient literature because they are essential for human survival. Traditional medicinal system is widely distributed in India. A major proportion of population mostly belonging to rural areas is still dependent on traditional system of medicines for their various health needs. It is evident that any system of healthcare solely will not capable to cope-up with the health care demands of public. Therefore, traditional and cultural medical knowledge has a catalyzing effect in meeting health care demands. There are a number of plants which have been reported for their wound healing activity due to presence of valuable phytoconstituents. Most of these studies involve random screening of plant or extracts for wound healing activity. Since in actual experiments and usage, all the plant products as cited in the table have shown efficacious results, there are increased needs to isolate and investigate each active ingredient that has a positive role in the healing process.

The combination of traditional and modern knowledge can produce better drugs for wound healing with fewer side effects. Such type of traditional knowledge on plant can form a basis for

clinical, medicinal, pharmacological and novel drug delivery system for wound healing products.

**CONFLICT OF INTEREST:** Declared none

#### REFERENCES:

1. Tortora GJ and Grabowski SR: Principle of Anatomy and Physiology. John Willey & Sons Inc, USA, Edition 12, 2003: 1-27.
2. Gupta LC, Gupta K and Gupta A: New Concise Medical Dictionary. AITBS, Delhi, Fourth Edition, 2008.
3. Garg HG and Longaker MT: Scarless Wound Healing. Marcel Dekker, New York, 2000: 1-22.
4. Mackay D and Miller AL: Nutritional support for wound healing. Alternative Medicine Review 2003; 8:359-377.
5. Kumar B, Vijayakumara M, Govindarajan R and Pushpangadan P: Ethanopharmacological approaches to wound healing-exploring medicinal plants of India. Journal of Ethanopharmacology 2007; 114:103-113.
6. Gupta N, Gupta SK, Shukla VK and Singh SP: An Indian community-based epidemiological study of wounds. Journal of Wound Care 2004; 13:323-325.
7. James Q and Rosso D: Wound care in dermatology office: where are we in 2011? Journal of American Academy of Dermatology 2011; S1-S7.
8. Bairy KL: Wound healing potential of plant product. Journal of Natural Remedies 2012; 2:11-20.
9. Vowden P, Vowden K and Carville K: Antimicrobials made dressing easy. Wounds International 2011; 2:1-6.
10. Kumar JK, Kumar HRC, GunashakaranV, RameshY, Babu KP, Narasimha PN, Venkatewarlu A and Reddy LP: Application of broad spectrum antiseptic povidone iodine as powerful action: a review. Journal of Pharmaceutical Science and Technology 2009; 1:48-58.
11. Schwartz RA and Al-Mutairi N: Topical antibiotics in dermatology: an update. Gulf Journal of Dermatology and Venereology 2010; 17:1-19.
12. Raina R, Prawez S, Verma PK and Pankaj NK: Medicinal plants and their role in wound healing. Vet Scan 2008; 3:1-7.

13. Heinrich M: Ethanobotany and natural products: The search for new molecules, new treatment of old diseases or a better understanding of indigenous cultures. *Current Topics in Medicinal Chemistry* 2003; 3:29-42.
14. Lazarus GS, Cooper DM, Knighton DR, Margolis DJ, Pecoraro RE, Rodeheave G and Robson MC: Definitions and guidelines for assessment of wounds and evaluation of healing. *Archives of Dermatology* 1994; 130:489-493.
15. Pawar RS and Toppo FA: Plant that heal wounds: a review. *Herba Polonica* 2012; 58: 47-65.
16. DiPietro LA and Burns AL: *Wound Healing Method and Protocol*. Humana Press, Totowa NJ, 3-15.
17. Fianagan M: The physiology of wound healing. *Journal of Wound Care* 2000; 9:299-300.
18. Wiksman LB, Solomonik I, Spira R and Tennenbaum T: Novel insights into wound healing sequence of events. *Toxicologic Pathology* 2007; 35:767-779.
19. Singer AJ and Clark R: Cutaneous wound healing. *New England Journal of Medicine* 1999; 341:738-746.
20. Nagori BP and Solanki R: Role of medicinal plants in wound healing. *Research Journal of Medicinal Plant* 2011; 5:392-405.
21. Orsted HL and Keast D: Basic principles of wound healing. *Wound care Canada* 2011; 9:4-12.
22. Diegelmann RF and Evans MC: Wound healing: an overview of acute, fibrotic and delayed healing. *Frontiers in Bioscience* 2004; 9:283-289.
23. Guo S and DiPietro LA: Factors affecting wound healing. *Journal of Dental Research* 2010; 8:219-229.
24. Galabuzi C, Agea JD, Fungo BL and Kamoga RMN: Traditional medicine as an alternative form of health care system: a preliminary case study of Nangabo sub-county, Central Uganda. *African Journal of Traditional Complementary and Alternative Medicine* 2010; 7:11-16.
25. Kamboj VP: Herbal medicine. *Current Sciences* 2000; 78:35-51.
26. Singh A, Singh P, Singh G and Pandey AK: Plant used in primary health practices in Vindhya region of eastern Uttar Pradesh, India. *International Journal of Herbal Medicine* 2014; 2:31-37.
27. Pal SK and Shukla Y: Herbal medicine: current status and future. *Asian Pacific Journal of Cancer Prevention* 2003; 4:281-288.
28. Payyappallimana U: Role of traditional medicine in primary health care an overview of perspectives and challenges. *Yokohama Journal of Social Sciences* 2009; 14:51-77.
29. Ramya S, Alaguchamy N, Maruthappan VM, Sivaperumal R, Sivalingam M, Krishnan A, Govindraji V and Jayakumararaj R: Wound healing ethanomedicinal plants popular among Malayali tribes in Vattal hills Dharmapuri, TN, India. *Ethanobotanical Leaflet* 2009; 13:1257-1271.
30. Mittal A, Sardana S and Pandey A: Herbal boon for wounds. *International Journal of Pharmacy and Pharmaceutical Sciences* 2013; 5:1-12.
31. Purna SK and Babu M: Collagen wound dressing a review. *Burns* 2000; 26:54-62.
32. Pierce GF: Inflammation in nonhealing diabetic wounds the space-time continuum does matter. *American Journal of Pathology* 2001; 159:399-403.
33. Lan CCE, Wu CS, Haung SM, Wu IH and Chen GS: High-glucose environment enhanced oxidative stress and increased interleukin-8 secretion from keratinocytes-new insights into impaired diabetic wound healing. *Diabetes* 2013; 62:2530-2538.
34. Mascolo N and Autore G: Biological screening of Italian medicinal plants for anti-inflammatory activity. *Phytotherapy Research* 1987; 1:28-31.
35. Akihisa T, Yasukawa K, Oinuma H, Kasahara Yoshimasa, Yamanouchi S, Takido M, Kumari K and Tamura T.: Triterpene alcohol from the flowers of compositae and their anti-inflammatory effects. *Phytochemistry* 1996; 43:1255-1260.
36. Shahriar M, Hossain MI, Sharmin FA, Akhter S, Haque MA and Bhuiyan MA: *In-vitro* antioxidant and free radical scavenging activity of *Withania somnifera* root. *Iosr Journal of Pharmacy*. 2013; 3:38-47.
37. Sen CK, Khanna S, Gordillo G, Bagchi D, Bagchi M and Roy S: Oxygen, oxidants and antioxidants in wound healing an emerging paradigm. *Annals of New York Academy of Sciences* 2002; 957:239-249.
38. Mistic AM, Gardner SE and Grice EA: The wound microbiome: modern approaches to examining the role of microorganisms in impaired chronic wound healing. *Advances in Wound Care* 2014; 3:502-510.
39. Robson MC: Wound infection: a failure of wound healing caused by an imbalance of bacteria. *Surgical Clinics of North America* 1997; 77:637-650.
40. Patel NA, Patel M and Patel RP: Formulation and evaluation of polyherbal gel for wound healing. *International Research Journal of Pharmaceuticals* 2011; 1:15-20.
41. Mudge E and Orsted H: Wound infection & pain management made easy. *Wounds International* 2010; 1:1-6.
42. McGuire L, Heffner K, Glaser R, Needleman B, Malarkey W, Dickinson S, Lemeshow S, Cook C and Melvin WS: Pain and wound healing in surgical patients. *Annals of Behavioral Medicine* 2006; 31:165-172.
43. Soni H and Singhai AK: A recent update of botanicals for wound healing activity. *International Research Journal of Pharmacy* 2012; 3:1-7.
44. Ghosh PK and Gaba A: Phytoextract in wound healing. *Journal of Pharmacy and Pharmaceutical Sciences* 2013; 16:760-820.
45. Krishnamoorthy JR, Sumitra S, Ranjith S, Gokulshankar S, Ranganathan S, Mohanty BK and Prabhakaran G: An *in-vitro* study of wound healing effect of a poly-herbal formulation as evidenced by enhanced cell proliferation and cell migration. *Egyptian Dermatology Online Journal* 2012; 8:1-7.
46. Sharma Y, Jeyabalan G, Singh R and Semwal A: Current aspects of wound healing agents from medicinal plants: a review. *Journal of Medicinal Plants Studies* 2013; 1:1-11.
47. Nagelschmidt M and Struck H: Hydroxyproline fractions in serum and urine of rats during wound healing. *Research in Experimental Medicine* 1977; 170:211-215.
48. Gal P, Toporcer T, Vidinsky B, Marry M, Novotny M, Kilik R, Smetana KJ, Gai T and Sobó T: Early changes in the tensile strength and morphology of primary sutured skin wounds in rats. *Folia Biologica* 2006; 52:109-115.
49. Sahu PK, Masih V, Gupta S, Sen DL and Tiwari A: Ethanomedicinal plants used in the healthcare system of tribes of Dantewada, Chattisgarh India. *American Journal of Plant Sciences* 2014; 5:1632-1643.
50. Jagatheeswari D, Deepa J, Ali HSJ and Ranganathan P: *Acalypha indica* L. an important medicinal plant: a review of its traditional uses, and pharmacological properties. *International Journal of Research in Botany* 2013; 3:19-22.
51. Mullick A, Mandal S and Bhattacharjee R: *In-vitro* assay of antioxidant and antibacterial activity of leaf extract and leaf derived callus extract of *Acalypha indica*. *International Journal of Pharmacy and Biological Sciences* 2013; 3:504-510.
52. Raja RV and Savitha S: Wound healing properties of medicinal plants (*Acalypha indica* and *Azadirachta indica*). *Journal of Bioscience Technology* 2013; 4:525-530.
53. Chatterjee P, Chakraborty B and Nandy S: *Aloe vera* plant: review with significant pharmacological activities. *Mintage Journal of Pharmaceutical & Medical Sciences* 2013; 1:21-24.
54. Hashemi SA, Abdollah Madani SA and Abediankenari S: The review on properties of *Aloe vera* in healing of cutaneous wounds. *BioMed Research International*. 2015; 1-6.
55. Purohit SK, Solanki R and Soni M: Experimental evaluation of *Aloe vera* leaves pulp as topical medicament on wound

- healing. International Journal of Pharmacology Research 2012; 2:110-112.
56. Sachan AK, Das DR, Dohare SL and Shuaib M: *Asparagus racemosus* (Shatavari): an overview. International Journal of Pharmacy and Chemical Sciences 2012; 1:937-941.
  57. Prabhath KG, Kumar S, Rajput R, Vijayalakshmi, Udupa AL and Gupta S: Effect of *Asparagus racemosus* (Liliaceae) Willd. on dead space model wound healing. Journal of Pharmacy Research 2011; 4:2772-2774.
  58. Biswas K, Chattopadhyay I, Banerjee RK and Bandyopadhyay U: Biological activities and medicinal properties of neem (*Azadirachta indica*). Current Science 2002; 82:1336-1345.
  59. Purohit SK, Solanki R, Soni R and Mathur V: Evaluation of wound healing activity of ethanolic extract of *Azadirachta indica* leaves in male albino rats. Asian Journal of Pharmacy and Technology 2013; 3:73-75.
  60. Al-Snafi AE: The chemical constituents and pharmacological effects of *Bryophyllum calycinum*: a review. International Journal of Pharmaceutical Sciences and Research 2013; 4:171-176.
  61. Kambooj A and Sluja AK: *Bryophyllum pinnatum* (Lam.) Kurz. phytochemical and pharmacological profile: a review. Phcognoy Reviews 2009; 3:364-374.
  62. Khare CP: Indian medicinal plants an illustrated dictionary. Springer Publishers, 2007.
  63. Kumar PS, Suresh E and Kalavathy S: Review on the potential herb of *Calotropis gigantea* (L) R. Br. Scholars Academic Journal of Pharmacy 2013; 2:135-143.
  64. Nalwaya N, Pokharna G, Deb L and Jain NK: Wound healing activity of latex of *Calotropis gigantean*. International Journal of Pharmacy and Pharmaceutical Sciences 2009; 1:176-181.
  65. Jamil SS, Nizami Q and Salam M: *Centella asiatica* (Linn.) Urban o' a review. Natural Product Radiance 2007; 6:158-170.
  66. Hasim P, Sidek H, Helan MHM, Sabery A, Palanisamy UD and Ilham M: Triterpene composition and bioactivities of *Centella asiatica*. Molecules 2011; 16:1310-1322.
  67. Kosalwatna S, Shaipanich C and Bhanganada K: The effect of one percent *Centella asiatica* cream on chronic ulcers. Siriraj Hospital Gazette 1988; 40:456-461.
  68. Shukla A, Rasik AM, Jain GK, Shankar R, Kulshrestha DK and Dhawan BN: *In-vitro* and *in-vivo* wound healing activity of asiaticoside isolated from *Centella asiatica*. Journal of Ethnopharmacology 1999; 65:1-11.
  69. Liu M, Dai Y, Li Y, Luo Y, Huang F and Gong Z: Madecassoside isolated from *Centella asiatica* herbs facilitates burn wound healing in mice. Planta Medica 2008; 74: 809-815.
  70. Vasavda K, Hedge PL and Harini: Pharmacological activities of turmeric (*Curcuma longa* Linn.): a review. Journal of Homeopathy and Ayurvedic Medicine 2013; 2:1-4.
  71. Naz S, Jabeen S, Ilyas S, Manzoor F, Aslam F and Ali A: Antibacterial activity of *Curcuma longa* varieties against different strains of bacteria. Pakistan Journal of Botany 2010; 42:455-462.
  72. Purohit SK, Solanki R, Mathur V and Mathur M: Evaluation of wound healing activity of ethanolic extract of *Curcuma longa* rhizomes in male albino rats. Asian Journal of Pharmacy Research 2013; 3:79-81.
  73. Ho WY, Ky H, Yeap SK, Rahim RA, Omar AR, Chai LH and Alitheen NB: Traditional practice, bioactivities and commercialization potential of *Elephantopus scaber* Linn. Journal of Medicinal Plant Research 2009; 3:1212-1221.
  74. Singh SDJ, Krishna V, Mankani KL, Vidya SM and Manohara YN: Wound Healing activity of the leaf extracts and deoxyelephantopin isolated from *Elephantopus scaber* Linn. Indian Journal of Pharmacology 2005; 37:238-242.
  75. Kumara SM, Pokharen N, Dahal S and Anuradha M: Phytochemical and antimicrobial studies of leaf extract of *Euphorbia nerifolia*. Journal of Medicinal Plants Research 2011; 5:5785-5788.
  76. Bigoniya P and Rana AC: Wound healing activity of *Euphorbia nerifolia* leaf ethanolic extract in rats. Journal of Natural Remedies 2007; 7:94-101.
  77. Gaur K, Rana AC, Nema RK, Kori ML and Sharma CS: Anti-inflammatory and analgesic activity of hydro-alcoholic leaves extract of *Euphorbia nerifolia* Linn. Asian Journal of Pharmacy and Clinical Research 2009; 2:26-29.
  78. Paarakh PM: *Ficus racemosa* Linn. an overview. Natural Product Radiance 2009; 8:84-90.
  79. Kumar MK: Enhancement of wound healing with roots of *Ficus racemosa* L. in albino rats. Asian Pacific Journal of Tropical Biomedicine 2012; 2: 276-280.
  80. Sharma S, Dhamija HK and Parashar B: *Jatropha curcas*: a review. Asian Journal of Research in Pharmaceutical Sciences 2012; 2:107-111.
  81. Nwala CO, Akaninwor JO and Monanu MO: Phytochemical screening and wound healing activities of extracts of *Jatropha curcas* leaf formulated in a simple ointment base. International Journal of Engineering Science Invention 2013; 2:72-75.
  82. Esimone CO, Nworu CS and Jackson CL: Cutaneous wound healing activity of a herbal ointment containing the leaf extract of *Jatropha curcas* L. (Euphorbiaceae). International Journal of Applied Research in Natural Products 2009; 1:1-4.
  83. Reddy NM: *Lantana camara* Linn. chemical constituents and medicinal properties: a review. Scholars Academic Journal of Pharmacy 2013; 2:445-448.
  84. Saxena M, Saxena J and Khare S: A brief review on therapeutic values of *Lantana camara* plant. International Journal of Pharmacy and Life Sciences 2012; 3:1551-1554.
  85. Nayak BS, Raju SS and Ramsubhag A: Investigation of wound healing activity of *Lantana camara* L. in *Sprague dawley* rats using a burn wound model. International Journal of Applied Research in Natural Products 2008; 1:15-19.
  86. Asha B, Nagabhushan A and Shashikala GH: Study of wound healing activity of topical *Ocimum sanctum* Linn. in albino rats. Journal of Chemical and Pharmaceutical Research 2011; 3:122-126.
  87. Goel A, Kumar S, Singh DK and Bhatia AK: Wound healing potential of *Ocimum sanctum* Linn. with induction of tumor necrosis factor- $\alpha$ . Indian Journal of Experimental Biology 2010; 48:402-406.
  88. Jain N, Jain R, Jain A, Jain DK and Chandel HS. Evaluation of wound-healing activity of *Acorus calamus* Linn. Natural Product Research 2010; 24:534-541.
  89. Lotlekar Resha, Marihal SC and Purnima A: Screening of *Allium sativum* Linn. bulb for wound healing and antioxidant activities. International Journal of Pharmaceutical Sciences 2011; 3:1292-1298.
  90. Vinothapooshan G and Sundar K: Wound healing effect of various extracts of *Adhatoda vasica*. International Journal of Pharma and Bio Sciences 2010; 1:530-536.
  91. Barua CC, Talukdar A, Begam SA, Sharma DK, Pathak DC and Barua AG: Wound healing activity of methanolic extract of leaves of *Alternanthera brasiliiana* Kuntz. using *in-vivo* and *in-vitro* model. Indian Journal of Experimental Biology 2009; 47:1001-1005.
  92. Mohanty A, Pradhan DK, Mishra MR, Sahoo JK, Mishra A, Nandy BC, Meena K and Mokade L: Preliminary phytochemical screening and wound healing activity of *Andrographis peniculata*. Journal of Chemical and Pharmaceutical Research 2010; 2:649-654.
  93. Bharat M, Verma DK, Shanbhag V, Rajput RS, Nayak D and Amuthan A: Ethanolic extract of oral *Areca catechu* promotes burn wound healing in rats. International Journal of Pharmaceutical Sciences Review and Research 2014; 25:145-148.
  94. Muralidhar A, Babu KS, Shanker RT, Reddanna P and Latha J: Evaluation of wound healing properties of bioactive

- fractions from the extract of *Butea monosperma* (Lam). stem bark. International Journal of Phytomedicine 2011; 3:41-49.
95. Nurismail E: Antimicrobial and wound healing activities of three *Cassia* species. Master of Science. Thesis, Faculty of Medicine and Health Sciences. Universiti Putra Malaysia. Malaysia, 2003.
  96. Nayak BS, Pereira LMP: *Catharanthus roseus* flower extract has wound-healing activity in *Sprague Dawley* rats. BMC Complementary and Alternative Medicine 2006; 6:1-6.
  97. Gurung S and Basnet NS: Wound healing properties of *Carica papaya* latex: *in-vivo* evaluation of mice burn model. Journal of Ethanopharmacology 2009; 121:338-341.
  98. Kuppast IJ and Nayak PV: Wound healing activity of *Cordia dichotoma* Forst. F. fruits. Natural Product Radiance 2006; 5:99-102.
  99. Shirwaikar A, Jahagirdar S and Udupa AL: Wound healing activity of *Desmodium triquetrum* leaves. Indian Journal of Pharmaceutical Sciences 2003; 65:461-464.
  100. Kumara SHM, Krishna V, Shankarmurthy K, Abdul RB, Mankani KL, Mahadevan KM, Harish BG and Raja NH: Wound healing activity of embelin isolated from the ethanol extract of leaves of *Embelia ribes* Burm. Journal of Ethanopharmacology 2006; 109:529-534.
  101. Roy K, Shivakumar H and Sarkar S: Wound healing potential of leaf extracts of *Ficus religiosa* on *Wistar albino* strain rats. International Journal of PharmaTech Research 2009; 1:506-508.
  102. Farzaei MH, Abbasabadi Z, Shams-Ardekani MR, Abdollahi M and Rahimi R: A comprehensive review of plants and their active constituents with wound healing activity in Iranian medicine. Wounds 2014; 26:197-206.
  103. Oloumi MM, Derakhshanfar A and Nikpoor A: Healing potential of liquorice root extract on dermal wounds in rats. Journal of Veterinary Research 2007; 62:147-154.
  104. Omale J and Sunday AB: Evaluation of acute dermal irritation and wound contraction by *Gymnema sylvestre* and *Datura metel* extracts in rats. American Journal of Biomedical and Life Sciences 2014; 2:83-88.
  105. Dodehe Y, Barthelemy A, Calixte B, Jean DN, Allico JD and Nelly F: *In-vitro* wound healing effect of *n*-butanol fractions from *Heliotropium indicum*. Journal of Innovative Trends Pharmaceutical Sciences 2011; 2:1-7.
  106. Sivagamy M, Jeganathan NS, Manavalan R and Senthamarai R: Wound Healing Activity of *Indigofera enneaphylla* Linn. International Journal of Advances in Pharmacy, Biology and Chemistry. 2012; 1:211-214.
  107. Mittal A, Sardana S and Pandey A: *Jasminum auriculatum* – an overview. International Journal of Pharmaceutical Innovations 2011; 1:1-6.
  108. Mohammad Z, Voskula R, Karra A, Reddy CR and Tirupathi A: Wound healing activity of ethanolic extract of *Allium sativum* on alloxan induced diabetic rats family (Liliaceae). International Journal of Science Inventions Today 2013; 2:40-57.
  109. Manjunatha BK, Krishna V, Vidya SM, Mankani KL, Manohara YN: Wound healing activity of *Lycopodium serratum*. Indian J Pharm Sci. 2007; 69: 283-287.
  110. Singh MP, Bhargava S, Bhaduarua RS and Sharma CS: Wound healing potential of alcoholic extract of *Mimosa pudica* Linn. leaves. Pharmacologyonline 2010; 2:32-38.
  111. Gupta N and Jain UK: Investigation of wound healing activity of methanolic extract of stem bark of *Mimusops elengi* Linn. African Journal of Traditional Complementary and Alternative Medicine 2011; 8:98-103.
  112. Gowda A, Shanbhag V, Bangalore ER, Shenoy S, Prabhu K and Narayanareddy M: The wound healing property of ethanolic extract of *Michelia champaca* flowers in diabetic rats. International Journal of Basic and Clinical Pharmacology 2014; 3:1036-1042.
  113. Gogoi J, Nakhuru KS, Chattopadhyay P, Rai AK and Veer V: Hypertrophic scar formation on application of terpenoid fraction of tuberous root of *Mirabilis jalapa* L. on excision wound model in *Wistar albino* rats. International Scholarly Research Notices 2014; 1-9.
  114. Lien LT, Tho NT, Ha DM, Hang PL, Nghai PT and Thang ND: Influence of phytochemicals in *Piper betle* Linn. leaf extract on wound healing. Burns & Trauma. 2015; 3:1-8.
  115. Umachigia SP, Jayaveerab KN, Kumar CKA, Kumar GS, Vrushabendra BM and Kumar DVK: Studies on wound healing properties of *Quercus infectoria*. Tropical Journal of Pharmaceutical Research 2008; 7:913-919.
  116. Karodi R, Jadhav M, Rub R and Bafna A: Evaluation of the wound healing activity of a crude extract of *Rubia cordifolia* L. (Indian madder) in mice. International Journal of Applied Research in Natural Products 2009; 2:12-18.
  117. Kotade K and Mohammed A: Wound healing activity of *Sesamum indicum* L. seed and oil in rats. Indian Journal of Experimental Biology 2008; 46:777-782.
  118. Akilandeswari S, Senthamarai R, Valarmathi R and Prema S: Wound healing activity of *Sida acuta* in rats. International Journal of Pharm Tech Research 2010; 2:585-587.
  119. Choudhary GP: Wound healing activity of the ethanolic extract of *Terminalia chebula* Retz. International Journal of Pharma and Bio Sciences 2011; 2:48-52.
  120. Talekar YP, Das B, Paul T, Talekar DY, Apte KG and Parab PB: Evaluation of wound healing potential of aqueous and ethanolic extracts of *Tridax procumbens* Linn. in *Wistar* rats. Asian Journal of Pharmacy and Clinical Research. 2012; 5(4): 141-45.
  121. Verma N, Khosa RL, Garg VK. Wound healing activity of *Wedelia chinensis* leaves. Pharmacologyonline. 2008; 2: 139-145.
  122. Yosufoglu HS. Topical anti-inflammatory and wound healing activities of herbal gel of *Ziziphus nummularia* L. (F. Rhamnaceae) leaf extract. International Journal of Pharmacology. 2011; 7(8): 862-867.

**How to cite this article:**

Saini S, Dhiman A and Nanda S: Traditional Indian Medicinal Plants with Potential Wound Healing activity: A Review. Int J Pharm Sci Res 2016; 7(5): 1809-19. doi: 10.13040/IJPSR.0975-8232.7(5).1809-19.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)