



Received on 15 April 2022; received in revised form, 24 June 2022; accepted 14 September 2022; published 01 December 2022

FORMULATION AND DEVELOPMENT OF SOLID SELF-EMULSIFYING DRUG DELIVERY SYSTEM FOR CIPROFLOXACIN

P. S. Gawade^{*}, H. A. Joshi and R. N. Patil

Department of Pharmaceutics, Shivnagar Vidya Prasarak Mandal's College of Pharmacy, Malegaon (BK), Baramati, Pune - 413115, Maharashtra, India.

Keywords:

Ciprofloxacin, S-SEDDS, Solubility enhancement, Adsorption technique

Correspondence to Author: Dr. Priyanka Shivalal Gawade

Assistant Professor,
Department of Pharmaceutics,
Shivnagar Vidya Prasarak Mandal's
College of Pharmacy, Malegaon (BK),
Baramati, Pune - 413115,
Maharashtra, India.

E-mail: priyanka.gawadesvpm@gmail.com

ABSTRACT: The purpose is to formulate the solid self emulsifying drug delivery system (S-SEDDS) to improve poor aqueous solubility and dissolution of ciprofloxacin. A solubility study was carried out, and then Soyabean oil (as oil phase), Tween 20 (as surfactant), and Span 80 (as co-surfactant) were selected for preparation of emulsion by titration method. Adsorption to solid carrier technique was used for solid SEDDS, areosil (Colloidal silica) used as adsorbent. In that formulation, I was having carbopol 934 and another formulation II HPMC K 4 M. The flow properties of all formulation were within the acceptable range; therefore, they can be easily filled into capsule. IR spectral and DSC thermograph analysis showed that there was chemical interaction between drug and polymer. *In-vitro* release studies were carried out in phosphate buffer pH 7.4. Release profile of pure drug formulation showed 38%, formulation I showed highest 54% cumulative drug release and formulation II showed lowest *i.e.* 14% drug release. Student's t-test was determined for *In-vitro* drug release data for different polymeric formulations and showed a p-value of 0.001891 (<0.01) 99.99% significant.

INTRODUCTION: The oral route is the easy, most convenient route for administration and the major route of drug delivery for the chronic treatment of many diseases¹. Nowadays, new chemical entities (NCE) have poor water solubility and low oral bioavailability². Poor water-soluble drug formulation is very difficult for pharmaceutical scientists as formulating new drug discovery methods and oral delivery are related with low bioavailability³⁻⁵. Lipid-based drug delivery systems (LBDDS) have more importance in the present due to their ability to improve the solubility and bioavailability of drugs with low aqueous solubility⁶⁻⁷.

The absorption of a drug having lipid-based formulation depends on a few factors, including particle size, degree of emulsification and rate of dispersion and precipitation⁸. Lipid-based formulations may include oil preparations or suspensions, emulsions, and self-emulsifying drug delivery systems (SEDDS)⁹. SEDDS is mixture of having equal physical properties of oils, surfactants, co-surfactants and co-solvents, formulations to improve the oral absorption of hydrophobic drug compounds.

SEDDS is formulated in solid form for oral administration like soft or hard gelatin capsules, lquisolid tablets and form fine stable oil-in-water (o/w) emulsions when contact with aqueous media and gentle agitation of the gastrointestinal fluids¹⁰. LBDDS have excellent prospects for improving bioavailability of BCS class II and IV drugs¹¹. Some marketed lipid based formulations are efavirenz (Sustiva), saquinavir (Fortovase),

<p>QUICK RESPONSE CODE</p> 	<p>DOI: 10.13040/IJPSR.0975-8232.13(12).5108-19</p>
<p>This article can be accessed online on www.ijpsr.com</p>	
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.13(12).5108-19</p>	

ritonavir (Norvir), clofazamine (Lamprene) and cyclosporine A (Sandimmune and Sandimmun Neoral) ^{6, 10}. In the present study, ciprofloxacin is selected for formulation. It is an anti-infective agent (Antibiotics) having Quinolones subclass and belongs to BCS Class IV, and because of its low solubility and permeability, it becomes difficult to formulate ¹²⁻¹³.

At present in the market, it is available in the form of its hydrochloride salt. The work aims to improve the solubility and increase its dissolution by formulating it in a Solid self-emulsifying drug delivery system. In SMEDDS of Ciprofloxacin using oleic acid (oil), tween-80 and pluronic F₁₂₇ (surfactant) and ethanol (co-surfactant) were selected, which improves drug solubility, absorption rate, and dissolution ¹⁴.

MATERIALS AND METHODS:

Materials: Ciprofloxacin was procured from Yarrow Chemicals Mumbai., Tween-20 was procured from Merk Pvt Ltd, Mumbai, Span-80, Methanol, carbopol 934, Aerosil and Talc were procured from Loba Chemi, Mumbai. Hydroxy Propyl Methyl Cellulose K-4M was procured from Colorcon, Asia Pvt Ltd, Goa. Olive oil and Soyabean oil were procured from Cosmo Chem. Empty hard gelatin capsules were procured from Yarrow Chemicals, Mumbai.

Method:

Solubility studies ¹⁵: Solubility studies of ciprofloxacin were performed; by the excess amount (100 mg) of solid ciprofloxacin was added to 5 ml oils (Soyabean oil and Olive oil) and various concentrations of Smix (15%, 20%, 25%, 30% and 35%). These solutions fill in 10 ml vial

using rotary shaker and at rotation speed 25 RPM for 48 hours. The contents of each vial were filtered through a 0.45 μ m whatman filter paper using sintered glass grade 1 and filtrate was diluted with respective solutions. The absorbance were measured UV- Spectrophotometer. The concentration of ciprofloxacin was quantified by calculating concentration of dissolved drug by using calibration curve equation.

Partition Coefficient ¹⁶⁻¹⁷: Partition coefficient of ciprofloxacin was determined by Shake-Flask method using n-octanol as organic phase and water as aqueous phase. Accurately weighed 25 mg and 50 mg of ciprofloxacin transferred to 10 ml of water and 10 ml of n-octanol was added to each separating funnel. Flask was shaken for vigorously for 15 minutes, and stay as it is to stable. The contents from each separating funnel were filtered through a 0.45 μ m whatman filter paper using sintered glass grade 1 and filtrate was diluted with respective solutions. The absorbance of solutions was measured UV spectrophotometer.

Pseudo Ternary Phase Studies ¹⁸⁻²¹: Ternary mixtures with varying composition of Tween-20 as surfactant, Span-80 as co-surfactant and Soyabean oil were prepared. Surfactant and co-surfactant were mixed in different ratios [4:1(A), 3:2(B), 2:3(C), 4:1(D), 1:1(E)].

For each phase diagram, oil and specific surfactant to co-surfactant ratio (1A-5A, 1B-5B, 1C-5C, 1D-5D and 1E-5E) were titrated. Twenty-five combinations of oil and Smix were made to find out the maximum emulsion area in the phase diagrams. Selected oil and Smix ratio was taken for further preparation of S-SEDDS.

TABLE 1: BATCHES OF EMULSION ON BASIS OF SURFACTANT MIXTURES CONCENTRATIONS

S. no.	Oil Phase (Soybean Oil)	Surfactant Mixtures Concentrations (Tween-20: Span-80) in ml				
		A (8:2)	B (6:4)	C (4:6)	D (2:8)	E (5:5)
Distilled Water Added (ml)						
1.	1	2.5	2.8	5.4	5.7	3.5
2.	2	3.2	2	5	6	3.7
3.	3	4	3.6	5.3	6.3	4
4.	4	4.4	3.9	5.8	6.5	5
5.	5	3	4.2	4.9	6.9	5.4

Preparation of S-SEDDS ²²⁻²⁴: Ciprofloxacin drug was solubilized in oil phase and then added in that Tween 20 and Span 80 mixtures (Smix) and methanol as co-solvent with titration method. That

mixture was mixed properly to get emulsion formed by the cyclo mixer. The emulsion was kept overnight for evaporated solvent at room temperature. For solid preparation of SEDDS,

adsorption to solid carrier technique was applied. Aerosol as an adsorbent was added to that emulsion, and then the emulsion gets fine powder

form. Further, that powder was evaluated and prepared capsule formulation of Ciprofloxacin SEDDS.

TABLE 2: COMPOSITIONS OF S-SEDDS FORMULATIONS

Formulation I			Formulation II		
S. no.	Ingredients	Quantity Taken	S. no.	Ingredients	Quantity Taken
1.	Ciprofloxacin SEDDS Powder	250 mg	1.	Ciprofloxacin SEDDS Powder	250 mg
2.	Carbopol 934	100 mg	2.	HPMC K 4 M	100 mg
3.	Aerosil	3 g	3.	Aerosil	3 g
4.	Talc	20 mg	4.	Talc	g

Formulation Characterization:

Infrared Spectroscopy²⁶: FT-IR spectrum of drug, polymers, excipient, and physical mixtures were recorded as potassium bromide (KBr). Over the region 4000-400cm⁻¹ for its authentication and to study characteristics peaks of drug and polymer using FT-IR spectrophotometer (JASCO Model FT/IR-4100, JASCO FT/IR 6100, Bruker). The observed peaks were compared with the characteristic peaks from the reported IR spectrum; the sample was authenticated.

Differential Scanning Calorimetry²⁶⁻²⁷: The DSC study was carried out for drugs, polymers, and physical mixtures and formulations. The DSC thermograms were recorded on a Hitachi 7020 instrument. Each sample (2-10mg) was heated in crimped aluminium pans at a scanning rate of 5°C/min from 0 to 300°C. An empty aluminium pan was used as a reference.

Micromeritics Properties²⁸⁻³⁰: The powder properties include bulk density (BD), tap density (TD), angle of repose (AR), Hausner ratio (HR), Carr's index (CI) were determined.

Drug Content Uniformity³¹: 100 mg sample powder of optimized solid formulation of ciprofloxacin was dissolved in 100 ml Phosphate buffer pH 7.4. Then this solution was kept for a night for complete solubility.

After 24 hrs 1 ml solution was withdrawn and diluted up to 10 ml with Phosphate buffer p^H 7.4 and absorbance of obtained sample was determined with UV spectrophotometer at wavelength of 278 nm.

In-vitro Drug Release Studies³³⁻³⁵: Drug release studies from S-SEDDS formulations were performed with 900 ml of phosphate buffer pH 7.4

as a medium at 37±0.5°C. Ciprofloxacin-loaded S-SEDDS capsules (equivalent to 250 mg of ciprofloxacin) were placed in a dissolution apparatus (Tab machines DRS-8) at the basket speed of 100 rpm. At predetermined time intervals (15, 30, 45, 60, 120, 180, 240, 300, 320, and 400 min), an aliquot (5ml) of samples were collected, filtered, and analyzed for the content of ciprofloxacin by spectrophotometrically at λ_{max} of 278 nm.

Statistical Analysis³⁶: Statistical analysis was applied for In-vitro drug release data for different polymeric formulations at a significance level <0.01 for student's t-tests.

RESULT AND DISCUSSION:

Solubility Studies³⁷⁻³⁹: Ciprofloxacin was insoluble in water. Among various lipids studied, the maximum solubility of ciprofloxacin was observed in Soyabean oil (0.18 mg/ml).

However, the minimum solubility was found in olive oil (0.17 mg/ml). Similarly, amongst various surfactants and co-surfactants, the maximum solubility of ciprofloxacin was observed in Tween 20 (0.17 mg/ml), and the minimum solubility was found in Span 80 (0.10 mg/ml).




FIG. 1: SOLUBILITY STUDIES OF CIPROFLOXACIN IN VARIOUS MEDIUMS

Partition Coefficient¹⁶⁻¹⁷:**TABLE 3: PARTITION COEFFICIENT OF CIPROFLOXACIN IN WATER/ N-OCTANOL PHASE**

S. no.	Drug added (mg)	n-Octanol (ml)	Water (ml)	Log P
1	25	10	10	0.0513
2	50	10	10	0.3698

Pseudo Ternary Phase Studies³⁷⁻³⁹: Fig. 2A, B, C, D and E represent the phase diagram constructed to identify the area of stable emulsion in the presence of ciprofloxacin. Initially, based on the results of maximum solubility, the pseudo-ternary diagram was plotted between Soyabean oil and Tween 20: Span 80. The said system yielded a

significant emulsion region. Further, titration of Soyabean oil was conducted with various combinations of Tween 20 and Span 80 (i.e., 4:1, 3:2, 2:3, 1:4, and 1:1). It was subsequently observed that a combination of Soyabean oil, Tween 20 and Span 80 (4:1) yielded the maximum emulsion region for the formulation of SEDDS.












FIG. 2: TERNARY PHASE DIAGRAM OF SYSTEM A (SOYABEAN OIL/SMIX (4:1)/WATER); SYSTEM B (SOYABEAN OIL/SMIX (3:2)/WATER); SYSTEM C (SOYABEAN OIL/SMIX (2:3)/WATER); SYSTEM D (SOYABEAN OIL/SMIX (1:4)/ WATER); SYSTEM E (SOYABEAN OIL/ SMIX (1:1)/ WATER). SHADED AREAS REPRESENT THE REGION OF EFFICIENT SELF-EMULSIFICATION.

Infrared Spectroscopy⁴⁰⁻⁴³:








FIG. 3: IR SPECTRA OF CIPROFLOXACIN, SOYABEAN OIL, TWEEN 20, SPAN 80, CARBOLPOL 934, HPMC K 4 M, PHYSICAL MIXTURE CIPROFLOXACIN AND CARBOPOL 934 (1:0.25), PHYSICAL MIXTURE CIPROFLOXACIN AND HPMC K 4 M (1:0.25), FORMULATION I AND FORMULATION II SHOWED A, B, C, D, E, F, G, H, I AND J RESPECTIVELY

Identification of Drug, Polymer and Other Excipients: The FTIR spectrum of ciprofloxacin showed characteristics peak of O-H stretch at 3689.16 cm^{-1} , anhydride C=O stretch at 1815.65 cm^{-1} , C-F stretch at 1061.66 cm^{-1} and N-H wagging at 715 cm^{-1} . The FTIR spectrum of Soyabean oil showed characteristics peak of C=O stretch at 1298.82 cm^{-1} , ester C-O-C stretch at 1097 cm^{-1} and C=C stretch at 1578.45 cm^{-1} . The FTIR spectrum of Tween 20 showed characteristics peak of C=O stretch at 1700.61 cm^{-1} , ester C-O-C stretch at 1097.3 cm^{-1} , O-H stretch at 2760.6 cm^{-1} , C-H stretch at 2960.2 cm^{-1} . The FTIR spectrum of Span 80 showed characteristics peak of C=O stretch at 1698.98 cm^{-1} , ester C-O-C stretch at 1103.08 cm^{-1} , O-H stretch at $32.92.86\text{ cm}^{-1}$, C-H stretch at 2961.16 cm^{-1} and CH=CH stretch 2929.34 cm^{-1} . The FTIR spectrum of Carbopol 934 showed characteristics peak of C=O stretching at 1708.89 cm^{-1} , C-C stretching at 1448.43 cm^{-1} , O-H bending at 1245.99 cm^{-1} and C-H bending at 1311.36 cm^{-1} . The HPMC K 4 M FTIR spectrum showed characteristics peak of O-H bond in alcohols at

3489.56 cm^{-1} , C-H stretching at 2832 cm^{-1} , and C-O bond in ethers at 1744.3 cm^{-1} .

FTIR of Physical Mixtures: The FTIR spectrum of physical mixture Ciprofloxacin powder and Carbopol 934 (1:0.25) showed a peak of O-H stretching at 3852.11 cm^{-1} , C=O stretching at 1569 cm^{-1} and N-H wagging at 711 cm^{-1} . In this physical mixture absence of C-F stretch and =C-H bending. The FTIR spectrum of the physical mixture of Ciprofloxacin powder and HPMC K 4 M (1:0.25) showed peak of O-H stretching at 3435.56 cm^{-1} , =C-H bending at 607 cm^{-1} , C-F stretching at 1073.19 cm^{-1} and C-H stretching at 3029.62 cm^{-1} . In this physical mixture absence of anhydride C=O stretch and amide N-H wagging.

FTIR of Formulations: The FTIR spectrum of formulation I showed a peak of C=O stretching at 1740.44 cm^{-1} , =C-H bending at 609.39 cm^{-1} , N-H wagging at 711.604 cm^{-1} and C-O stretching 1159.01 cm^{-1} . In this formulation absence of O-H stretch and C-F stretch. The FTIR spectrum of

formulation II showed peak of =C-H bending at 620.966 cm^{-1} , C-H stretching at 2900.41 cm^{-1} , C-F stretching at 1028.54 cm^{-1} , C-O stretching at 1633.41 cm^{-1} , C-O-C stretch at 1097.3 cm^{-1} , C=C

stretching at 1470.46 cm^{-1} and C-C stretch at 1633.41 cm^{-1} . In these formulation absences of C=O stretch and N-H wagging.

Differential Scanning Calorimetry⁴⁴⁻⁴⁷:




FIG. 4: DSC THERMOGRAM OF CIPROFLOXACIN, CARBOPOL 934, HPMC K 4 M, PHYSICAL MIXTURE OF CIPROFLOXACIN AND CARBOPOL 934 (1:0.25), PHYSICAL MIXTURE OF CIPROFLOXACIN AND HPMC K 4 M (1:0.25), FORMULATION I AND FORMULATION II SHOWED A, B, C, D, E, F AND G RESPECTIVELY

DSC Study of Drug and Polymers: DSC thermogram of ciprofloxacin showed an

endothermic peak at 236.9°C indicated the drug's melting point with amorphous nature. DSC

thermogram of Carbopol 934 showed endothermic peaks at 222.3°C and 254°C. In that 254°C was the relevant endothermic peak of its melting point. DSC thermogram of HPMC K 4 M was showed endothermic peak at 87.68°C.

DSC Study of Physical Mixtures: DSC thermogram of physical mixture Ciprofloxacin powder and Carbopol 934 (1:0.25) showed an endothermic peak at 122°C, indicating the formation of a homogeneous mixture between drug and polymer. DSC thermogram of the physical mixture of Ciprofloxacin powder and HPMC K 4 M (1:0.25) was not show any endothermic peak. The drug peak completely emerged, indicating the formation of a homogeneous mixture between drug and polymer and would be in amorphous form.

Micromeritics Properties ²⁸

TABLE 4: MICROMERITICS PROPERTIES OF POWDER

Batch	Mean bulk density (g/ml) ± SD	Mean tapped density (g/ml) ± SD	Mean angle of repose ^o ± SD	Mean percent compressibility Index % ± SD	Mean hausners ratio ± SD
F	0.4327±0.0030	0.5278±0.0014	33.51±0.2274	17.50±0.1594	1.2230±0.0103

Bulk Density and Tapped Density: The bulk density and tap density of formulations was found to be 0.4327 g/ml and 0.5278 g/ml respectively.

Angle of Repose: The angle of repose of formulations ranges from 29.42° to 33.86°, indicating acceptable flow properties.

Percent Compressibility Index: The compressibility index of formulations is in the

DSC Study of Formulations: The DSC thermogram of formulation I showed an endothermic peak at 138.8°C indicating the formation of a homogenous mixture between drug and polymer and also indicates amorphous nature in between 160°C to 252°C regions. Another obtained endothermic peak at 253.8°C indicates that some quantity of polymer remains uncomplex in the formulation. For that, we could be increased the ratio of drug and polymer. The DSC thermogram of formulation II did not show any endothermic peak. The drug peak completely emerged, indicating the formation of a homogeneous mixture between drug and polymer in the amorphous nature. The exothermic peak was at 322.3°C because of decomposition.

range of 12.121%-18.182%, indicating the good flow properties.


Hausner's Ratio: Hausner's ratio of all formulations <1.25 indicate good flow ability. Capsule preparations comply with the test if not more than one individual content is outside the limits of 85 to 115 percent of the average content and none was outside the limits of 75 to 125 percent of the average content.

Drug Content Uniformity ⁴⁸:

TABLE 4: DRUG CONTENT UNIFORMITY OF FORMULATION

S. no.	Batch code	Absorbance	% Drug content
1.	Formulation I	0.187	96.84
2.	Formulation II	0.175	90.15

In-vitro drug Release Study ⁴⁹⁻⁵⁰:






FIG. 5: A, B AND C SHOWED ZERO-ORDER KINETIC MODELING FOR FORMULATION, MATRIX (HIGUCHI) KINETIC MODELING FOR FORMULATION, AND KROSMAYER - PEPPAS KINETIC MODELING FOR FORMULATION, RESPECTIVELY

Dissolution data were fitted in zero order, Higuchi (Matrix) model and Krosmayer-Peppas model. The best-fitted model for formulation I and pure drug formulation was the Higuchi model, and formulation II was fitted in the Zero-order model. The release profile of pure drug formulation showed 38%, formulation I showed the highest 54% cumulative drug release, and formulation II showed the lowest i.e. 14% drug release.

Dissolution data also fitted into Peppas and Krosmayer model it shows n-value 0.7362, 0.8184, and 0.5095, respectively, for pure drug formulation, formulation I and formulation II respectively, which was greater than 0.1 shows non-fickian or anomalous transport diffusion release mechanism.

Statistical Analysis⁵¹: Student's t-test was determined for In-vitro drug release data for different polymeric formulations showed a p-value 0.001891 (<0.01) it was 99.99% significant.

CONCLUSION: S-SEDDS of ciprofloxacin was prepared from aerosil (colloidal silica) filled in hard gelatin capsules, while liquid SEDDS was composed of soyabean oil, tween 20 and span 80. In vitro release studies were carried out in phosphate buffer pH 7.4. The release profile of pure drug formulation showed 38%, formulation I showed the highest 54% cumulative drug release, and formulation II showed the lowest, i.e., 14% drug release. Formulation II follows the Zero-order model, while formulation I follow Higuchi (matrix) model. Dissolution data also fitted into Peppas and Krosmayer model it shows n-value 0.7362, 0.8184, and 0.5095, respectively, for pure drug, formulation I and formulation II, which was greater than 0.1 showing non-fickian or anomalous transport

diffusion release mechanism. So, finally from all results, it was concluded that the Solid self emulsifying drug delivery system is a suitable approach for solubility and dissolution enhancement of poorly water-soluble ciprofloxacin.

ACKNOWLEDGEMENT: I thank my college Shivanagar Vidya Prasark Mandal's College of Pharmacy, Malegaon (bk), Baramati, Pune, Maharashtra, for supporting my PG research work. The absorption spectrum of optimized polymers was recorded by potassium bromide dispersion technique using JASCO 6100 and 4100 spectrophotometers at Maharashtra Institute of Technology, Pune, and Pune University. The DSC thermograms were recorded on a Hitachi 7020 at Maharashtra Institute of Technology, Pune.

CONFLICTS OF INTEREST: Nil

REFERENCES:

- Reddy LH and Murthy RH: Lymphatic transport orally administered drugs. *Indian Journal of Experiments Biology* 2002; 40: 1097-1109.
- Robinson JR: Introduction: Semi-solid formulations for oral drug delivery. *Bulletin of Technology Gattefosse* 1996; 89: 11-13.
- Dey P, Maiti S, Sa B and Sen K: Self emulsification of poorly soluble and highly permeable drugs: An overview. *International Journal Pharmaceutics* 2009; 1: 67-72.
- Giri TK, Badwaik H, Alexander A and Tripathi DK: Solubility enhancement of Ibuprofen in the presence of hydrophilic polymer and surfactant. *International Journal of Applied Biology and Pharmaceutical Technology* 2010; 1: 793-800.
- Kanika S, Pawar Y, Bansal B and Arvind K: Self emulsifying drug delivery system: A strategy to improve oral bioavailability. *Current Research and Informatinal Sciences* 2010; 3: 42-49.
- Kalepu S, Manthina M and Padavala V: Oral lipid-based drug delivery systems-an Overview. *Acta Pharmaceutica Sinica B* 2013; 3(6): 361-372.

7. Agrawal S, Giri TK, Tripathi DK and Alenxander A: A Review on novel therapeutic strategies for the enhancement of solubility for hydrophobic drugs through lipid and surfactant based self micro emulsifying drug delivery system: a Novel approach. *American Journal of Drug Discovery and Development* 2012; 2(4): 143-183.
8. Pouton CW: Lipid formulations for oral administration of drugs: nanoemulsifying, self-emulsifying and self-microemulsifying drug delivery systems. *European Journal of Pharmaceutical Sciences* 2000; 11: 93-98.
9. Hauss DJ, Fogal SE, Ficorilli JV, Price CA and Roy T: Lipid-based delivery systems for improving the bioavailability and lymphatic transport of a poorly water-soluble LTB4 inhibitor. *Journal of Pharmaceutical Sciences* 1998; 164-169.
10. Gursoy RN and Benita S: Self-emulsifying drug delivery systems (SEDDS) for improved oral delivery of lipophilic drugs, *Biomedical Pharmacotherapy* 2004; 58: 173-182.
11. Mullertz A, Ogbonna A, Ren S and Rades T: New perspectives on lipid and surfactant based drug delivery systems for oral delivery of poorly soluble drugs, *Journal of Pharmacy and Pharmacology* 2010; 62: 1622-1636.
12. Jain V, Singodia D, Gupta GK, Garg D and Keshava S: Ciprofloxacin surf-plexes in sub-micron emulsions: A novel approach to improve payload efficiency and antimicrobial efficacy. *International Journal Pharmaceutics* 2011; 409: 237-244.
13. Refat MS, El-Megharbel SM and Adam AMA: Physicochemical studies on different vital kinds of metal-drug binding and their application. *International E publication* 2013; 12-50.
14. Bakshi M, Mahajan SC and Bhandari G: Formulation design and evaluation of self-micro emulsifying drug delivery system of ciprofloxacin. *International Journal of Pharmaceutical Research and Biological Sciences* 2013; 2(1): 29-53.
15. Baka E, Comer JEA and Novak KT: Study of equilibrium solubility measurement by saturation shake flask method using hydrochlorothiazide as model compound. *J of Pharmaceut and Biomedical Analysis* 2008; 46: 335-341.
16. Berthod A and Broach SC: Determination of liquid-liquid partition coefficients by separation methods. *Journal of Chromatography* 2004; 1037: 3-14.
17. Bhagav P, Deshpande P, Pandey S and Chandra S: Development and validation of stability indicating UV spectrophotometric method for the estimation of brimonidine tartrate in pure form, formulations and Preformulation studies. *Der Pharmacia Lettre* 2010; 2(3): 106-122.
18. Nagasreenu Kommana, Kanchan Bharti, Dandabhattula Bhavya Surekha, Sathish Thokala and Brahmshwar Mishra: Development, optimization and evaluation of losartan potassium loaded Self Emulsifying Drug Delivery System. *Journal of Drug Delivery Science and Technology* 2020; 60: 102026.
19. Shimul Halder, Amena Islam, Md. Abdul Muhit, Manik Chandra Shill and Syed Shabbir Haider: Self-emulsifying drug delivery system of black seed oil with improved hypotriglyceridemic effect and enhanced hepatoprotective function. *Journal of Functional Foods* 2021; 78: 104391.
20. Salim Saifullah, Tasmina Kanwal, Shafi Ullah, Muhammad Kawish, Shahida Muhammad Habib, Imdad Al, Abubakar Munir, Muhammad Imran and Muhammad Raza Shah: Design and development of lipid modified chitosan containing muco-adhesive self-emulsifying drug delivery systems for cefixime oral delivery. *Chemistry and Physics of Lipids* 2021; 235: 105052.
21. Haoshi Gao, Haoyue Jia, Jie Dong, Xinggong Yang, Haifeng Li and Defang Ouyang: Integrated *in-silico* formulation design of self-emulsifying drug delivery systems. *Acta Pharmaceutica Sinica B* 2021; 11(11): 3585-3594.
22. Julian David Friedl, Arne Matteo Jörgensen, Bao Le-Vinh, Doris Elfriede Braun, Martina Tribus and Andreas Bernkop-Schnürch: Solidification of self-emulsifying drug delivery systems (SEDDS): Impact on storage stability of a therapeutic protein. *Journal of Colloid and interface* 2021; 584: 684-697.
23. Indrani Maji, Srushti Mahajan, Anitha Sriram, Pravin Medtiya, Ravindra Vasave, Dharmendra Kumar Khatri, Rahul Kumar, Shashi Bala Singh, Jitender Madan and Pankaj Kumar Singh: Solid self emulsifying drug delivery system: Superior mode for oral delivery of hydrophobic cargos. *Journal of Controlled Release* 2021; 337: 646-660.
24. Min-Jong Choi, Jung Suk Kim, Heesun Yu, Mi Ran Woo, Ji Eun Choi, Kyungho Baek, Jong Oh Kim, Yong Seok Choi, Han-Gon Choi and Sung Giu Jin: Comparison of the physicochemical properties, aqueous solubility, and oral bioavailability of rivaroxaban-loaded high-pressure homogenised and Shirasu porous glass membrane emulsified solid self-nanoemulsifying drug delivery systems. *Journal of Molecular Liquids* 2022; 346: 117057.
25. Naumann C, Brumm T and Bayerl TM: Phase titration behavior of single phosphatidylcholine bilayers on a solid spherical support studied by DSC, NMR and FT-IR, *Biophysical Journal* 1992; 63: 1314-1319.
26. Wang Y and Voth GA: Unique spatial heterogeneity ionic liquids. *Journal of American Chemical Society* 2005; 127: 12192-12193.
27. Byrn S, Pfeiffer R, Ganey M, Hoiberg C and Poochikian G: Pharmaceutical solids: A strategic approach regulatory considerations. *Pharmaceutical Research* 1995; 12(7): 945-954.
28. Gibson M: Pharmaceutical preformulation and formulation: A practical guide from candidate drug selection commercial dosage form. *Newyork Second Edition* 2009; 367-373.
29. Lachman and Lieberman L: *The Theory and Practice of Industrial Pharmacy*. 3rd Ed Varghese Publishing House Bombay 1987; 293-317.
30. Martin A: *Physical Pharmacy*, B.I Waverly Pvt Ltd, 4th Edition 332.
31. Santosh Koirala, Prabin Nepal, Govinda Ghimire, Rojina Basnet, Ishwori Rawat, Aashma Dahal, Jitendra Pandey and Kalpana Parajuli-Baral: Formulation and evaluation of mucoadhesive buccal tablets of aceclofenac. *Heliyon* 2021; 7(3): 06439.
32. Hao Loua and Michael J. Hagemana: Particle Size Limits to Pass the Acceptance Value (AV)-Based USP Content Uniformity Test for Tablets. / *Journal of Pharmaceutical Sciences* 2021; 110: 3527-3534.
33. Yadav AV, Yadav VB and Shete AS: *Experimental Biopharmaceutics and pharmacokinetic*. Ed 1st Nirali Prakashan 2011; 1.1-1.64.
34. Brahamankar DM and Jaiswal SB: *Biopharmaceutics and pharmacokinetic a treatise*. Ed 2nd Vallabh Prakashan Delhi 2009; 431-434.
35. Rusha Sardhara, Kaushalendra Chaturvedi, Harsh S. Shah, Bhavani Prasad Vinjamuri, Antoine Al-Achi, Kenneth R. Morrise and Rahul V. Haware: Predictive Performance Comparison of Computed Linear and Quadratic Multivariate Models for In-Situ UV Fiber Optics Tablet Dissolution Testing. *European Journal of Pharmaceutical Sciences* 2021; 161: 105806.

36. Montgo DC: Design and analysis of experiments. USA, John Wiley sons and Inc. Eighth Edition 2013; 65-130.
37. Min-Jong Choi, Jung Suk Kim, Heesun Yu, Mi Ran Woo, Ji Eun Choi, Kyungho Baek, Jong Oh Kim, Yong Seok Choi, Han-Gon Choi and Sung Gyu Jin: Comparison of the physicochemical properties, aqueous solubility and oral bioavailability of rivaroxaban-loaded high-pressure homogenised and Shirasu porous glass membrane emulsified solid self-nanoemulsifying drug delivery systems. *Journal of molecular liquids* 2021; 346: 117057.
38. Moataz B. Zewail, Sanaa A. El-Gizawy, Mohamed A. Osman and Yusuf A. Haggag B: Preparation and *In-vitro* characterization of a novel self-nano emulsifying drug delivery system for a fixed-dose combination of candesartan cilexetil and hydrochlorothiazide. *Journal of Drug Delivery Science and Technology* 2021; 61: 102320.
39. Nagasreenu Kommana, Kanchan Bharti, Dandabhattula Bhavya Surekha, Sathish Thokala and Brahmeshwar Mishra: Development, optimization and evaluation of losartan potassium loaded Self Emulsifying Drug Delivery System. *J of Drug Deliv Sci and Tech* 2020; 60: 102026.
40. Stuart B. Infrared spectroscopy: Fundamentals and applications; An TS Analytical Techniques in the Sciences. Wiley and Sons Ltd 2004; 71-93.
41. Lampman GM, Pavia DL, Kriz GS and Vyan JR: Spectroscopy. Ed 4th Delhi Cengage Learning Publication 2010; 29.
42. <http://www2ups.edu/faculty/hanson/Spectroscopy/IR/IRInterpretation>.
43. <http://orgchem.colorado.edu>.
44. Brown ME: Handbook of thermal analysis and calorimetry. Elsevier science BV 2003; 1: 279-361.
45. Suye Li, Yanna Zhao, Lili Wang, Hengqian Wu, Yan Gao, Lingxuan Zhang and Zhengping Wang and Jun Han: A Evaluation of water induced phase transition of Fexofenadine Hydrochloride during wet granulation process using NIR and DSC techniques. *Microchemical Journal* 2021; 169: 106497.
46. Rahamatullah Shaikh, Saeed Shirazian, Sarah Guerin, Eoin Sheehan d, Damien Thompson and Gavin M: Walker and Denise M. Croker: Understanding solid-state processing of pharmaceutical cocrystals *via* milling: Role of tablet excipients. *International Journal of Pharmaceutics* 2021; 601: 120514.
47. Yuan Chen, Tarun Tejasvi Mutukuri, Nathan E. Wilson and Qi (Tony) Zhou: Pharmaceutical protein solids: Drying technology, solid-state characterization and stability: *Advanced Drug Delivery Reviews* 2021; 172: 211–233.
48. Indian Pharmacopoeia. Government of India Ministry of Health and Family Welfare. The Indian Pharmacopoeia Commission. Ghaziabad 2010; 1: 182-184.
49. Fan Zhang, Yinping Zhou, Ni Wu, Ranran Jia, Aijing Liu, Bo Liu, Zhou Zhou, Haitang Hu, Zhihui Han and Xiang Ye: *In-silico* prediction of bioequivalence of Isosorbide Mononitrate tablets with different dissolution profiles using PBPK modeling and simulation. *European Journal of Pharmaceutical Sciences* 2021; 157; 105618.
50. Tao Lu and Timo L.M. ten Hagen: A novel kinetic model to describe the ultra-fast triggered release of thermosensitive liposomal drug delivery systems. *Journal of Controlled Release* 2020; 324: 669-678.
51. Shumyla Mehrj and Yamini Sudha Sistla: Optimization of process conditions for the development of pectin and glycerol based edible films: Statistical design of experiments. *Electronic Journal of Biotechnology* 2022; 55: 27–39.

How to cite this article:

Gawade PS, Joshi HA and Patil RN: Formulation and development of solid self-emulsifying drug delivery system for ciprofloxacin. *Int J Pharm Sci & Res* 2022; 13(12): 5108-19. doi: 10.13040/IJPSR.0975-8232.13(12).5108-19.

All © 2022 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)