



Received on 06 June 2023; received in revised form, 25 October 2023; accepted, 23 November 2023; published 01 March 2024

A CUMULATIVE REVIEW ON THERAPEUTIC FLORAE USED TO TREAT DIABETES MELLITUS

Sree Mahalakshmi Pasumarthy* and Udaya Guttikonda

Department of Pharmacology, Narayana Pharmacy College, Nellore - 524003, Andhra Pradesh, India.

Keywords:

Diabetes mellitus, Insulin resistance, Blood glucose, Insulin secretagogues, oxidative stress, Phytoconstituents

Correspondence to Author:

Sree Mahalakshmi Pasumarthy

Assistant Professor,
Department of Pharmacology,
Narayana Pharmacy College, Nellore
- 524003, Andhra Pradesh, India.

E-mail: pasumarthysreemahalakshmi3@gmail.com

ABSTRACT: Diabetes Mellitus is a chronic ailment characterized by clinical manifestations like hyperglycaemia. The development of novel drugs is based on the understanding of insulin resistance and oxidative damage leading to secondary complications of diabetes such as Retinopathy, Nephropathy etc. In the modern era, there are numerous allopathic drugs available for the treatment of the ailment. Herbs include a variety of chemical components that are responsible for their therapeutic effects, including polyphenols, saponins, terpenoids, alkaloids, sesquiterpenes, and flavonoids. Herbal remedies, which include traditional medicines, have the ability to treat a wide range of disorders. The use of herbs to treat a condition provides extra benefits by targeting the root cause of the disease. They are less expensive, more effective at lower dose frequencies and have less side effects when compared to allopathic medicines. This review includes a wide range of various phytoconstituents derived from different medicinal plants for the treatment of diabetes mellitus.

INTRODUCTION: Diabetes Mellitus is a chronic metabolic condition known as hyperglycaemia that is brought on by a rise in blood glucose levels. Blood sugar levels will be higher than they would be under normal circumstances. It is mostly caused by a drop in insulin production or activity in the body. When we ingest foods (carbohydrates, lipids, fats, and proteins), glucose enters our cells and gives us the energy we need for daily activities¹. To provide energy, the blood transports glucose to every single cell. The hormone Insulin is created by pancreatic beta cells and is the final step in the glucose cycle.

Higher blood glucose levels result from inadequate insulin production or improper body utilisation by the pancreas. Medical consequences brought on by the condition include damage to the eyes and nerves, neurological problems, heart problems, and impairments in metabolic function.

Diabetes causes an imbalance between insulin resistance and pancreatic insulin production². 9.3% of people in the world today have diabetes. In the years after 2045, it will increase by up to 10.2%. Natural herbs are now crucial in the treatment of many diseases following the pandemic of COVID-19. When compared to allopathic medications, herbal remedies are more cost-effective, have fewer adverse effects, and require less frequent administration. In order to control diabetes, numerous herbs have been mentioned in various literature sources³. According to ADA, there are three forms of DM^{4, 5, 6}.

	<p style="text-align: center;">DOI: 10.13040/IJPSR.0975-8232.15(3).626-35</p>
	<p style="text-align: center;">This article can be accessed online on www.ijpsr.com</p>
<p>DOI link: https://doi.org/10.13040/IJPSR.0975-8232.15(3).626-35</p>	

Type-1	Type-2	Gestational Diabetes
<ul style="list-style-type: none"> • Rare in people • Body does not produce Insulin • Auto immunity occurs • Causes: bacteria, virus, vitamin.D deficiency • Children, adults are mainly affected 	<ul style="list-style-type: none"> • Common (90-95%) • Insulin is produced but it is not used by the body and develops resistance • Causes: Obesity, lifestyle • Adolescent people are mainly affected. 	<ul style="list-style-type: none"> • During pregnancy

FIG 1: TYPES OF DM

Risk factors involved in Diabetes Mellitus:




FIG 2: RISK FACTORS INVOLVED IN DM

DM is caused by various genetic, metabolic and environmental factors. White individuals living in the USA, the United Kingdom, and Japan are all significantly impacted. Imbalance in genes like GLUT2 (transportation of glucose into pancreas), GCGR (glucagon hormone), ABCC8 (releases

insulin), and TCF7L2 (controls glucose release and synthesis)⁷. Familial history is another important element because it hereditarily transmits the disease to the healthy individuals. Age, gender, stress, high cholesterol levels, an unhealthy lifestyle, and obesity are other risks⁸.

TABLE 1: MEDICINAL FLORAE USED INMANAGING DM

S. no.	Vernacular name	Botanical name	Plant part used	Active phytoconstituent	Major Findings	Reference
1	Custard apple	<i>Annona squamosa</i>	leaves	Proantho-cyanidin Rutin Squafosacin	Enhances homeostasis and glucose secretion. Additionally, the plant raises GLP-1 levels in the blood.	9

2	Bitter apple Fox melon Bitter - Water melon	<i>Citrullus colocynthis</i>	fruit seed	Cucurbitacin Elate ricin B	Potent novel insulin promoter to reduce hyperglycaemia by GLUT 4 translocation Reduces hyperglycaemia through glycogenolysis and lowering G-6-P levels	10 11
3	Alanda	<i>Ephedra foeminea</i>	Aerial parts	Limonene Kaempferol	The plant's aqueous extract possesses anti-radical properties.	12
4	Loquat	<i>Eriobotrya japonica</i>	leaf seed	Naringenin Procyanidin Rutin	Loquat leaf extract in aqueous form is rich in minerals and vitamins and is used to reduce insulin resistance and oxidative stress. The seed's ethanol extract exhibits glucose tolerance properties.	13 14
5	Jamun	<i>Eugenia jambolana</i>	seed	Peonidin Malvidin Petunidin Delphinidin	demonstrates HHK-1 gene-mediated hypoglycaemic inhibition and enhances liver, kidney functioning. Utilised to treat both short-term and long-term conditions.	15,16
6	Danshen Gegen	<i>Salvia miltiorrhiza +Radix Pueraria lobata</i>	Whole plant	Puerarin Tanshinone II A	By lowering oxidative stress, inflammation, and promoting metabolism, the combination of these two herbs is used to treat DM	17
7	Behund Thohar	<i>Euphorbia nerifolia</i>	stem	Neriifolin Euphol Cycloartenol Neriifoliene	By enhancing the GLUT 4 transporter, a hydroalcoholic extract of the stem possesses insulin secreting action. Free radical scavenging activity by inhibiting α -amylase	18
8	Alfalfa	<i>Medicago sativa</i>	flower leaf	Stachydrine Myristic acid Arginine	The plant's aqueous extract may be able to stimulate the production of insulin from mesenchymal stem cells, which are similar to pancreatic beta cells.	19
9	Baikal skull cap	<i>Scutellaria coptis</i>	Whole plant	Baicalin Wogonoside Coptisine Berberine	Regulates intestinal flora and anti- inflammatory activity by interacting with TLR4	20
10	Purple heart	<i>Tradescantia pallida</i>	leaves	Spathulenol Caryophyllene oxide α -copaene β -caryophyllene	When compared to pure drug extract, Niosomal leaf extract has a stronger anti-diabetic impact, according to an <i>in-vitro</i> glycosylation experiment.	21
11	Stinging needle	<i>Urtica dioica</i>	leaves Whole plant	Vitamin B1 Vitamin K Xanthophylls Sistosterin	Reduced insulin resistance and improved lipid metabolism are the effects of leaf extract. Along with treating diabetes, the UD extract also lessens neurological side effects including memory loss	22 23
12	Fenu Greek	<i>Trigonella foenum- graecum</i>	seed	Diosgenin Coumarin	The functions of many organs, including the pancreas, liver, and kidneys, are demonstrated by seed extract. Both oral and iv administration lower blood glucose levels	24
13	safflower	<i>Carthamus tinctorius</i>	Seed oil	Carthamin Guanosine Cynarside Daucossterol	In individuals with metabolic disorders, seed oil extract was utilised to reduce insulin resistance	25

14	Hing	<i>Ferula assafoetida</i>	Oleo gum resin	(Z)- β -ocimene β -pinene	Encouraging the development of β -cells in the pancreas, lowering oxidative stress and insulin resistance. Also used to treat chronic DM	26
15	Bael Indian quince	<i>Aegle marmelos correa</i>	Stem bark	Aegeline Imperatorin Citral Lupeol	The plant's stem, bark contains UFG and lowers blood sugar and oxidative stress. In the long run, UBD is a powerful chemical to treat diabetes	27
16	Coriander	<i>Coriandrum sativum</i>	seed	Linalool Limonene Γ -terpinene α -pinene	Hyperglycaemia can be treated using seed extract, which also functions as an insulin secretor	28
17	Mulberry	<i>Morus alba</i>	fruit	Caffeic acid Gallic acid Vanillic acid Myricetin	In an RCT, mulberry demonstrates its capacity to regulate blood sugar and insulin levels before dawn	29
18	Okra	<i>Abelmoschus esculentus</i>	Edible pod	3-O-gentiobioside	Fibres present in okra are responsible for slow absorption of glucose in GIT. It is not advised to use this vegetable and allopathic drugs at the same time	30
19	Gum Arabic tree	<i>Acacia nilotica</i>	Pods Leaf Bark	Ellagic acid Kaempferol Quercetin	The effects of diabetes are lessened by the leaf extract	31
20	Giloy	<i>Tinospora sinensis</i>	Stem	Berberine Caffeic acid Myricetin Ferulic acid	The stem extract is protective against diabetes by reducing chemical mediators and free radicals	32
21	Red onion	<i>Allium cepa</i>	Whole plant	Allicin Fisetin Quercetin Alliinase	It is utilised as a food supplement to lower blood sugar levels	33
22	Barberry	<i>Berberies vulgaris</i>	Root Bark	Berberine Oxyacanthine Berbamine Palmatine	Reduced triglyceride and cholesterol levels, controls body weight	34
23	Red silk cotton tree	<i>Bombax ceiba</i>	flower	Rutin Resveratrol Baicalein Hesperetin	Flower extract aids in removing free radicals from the body. For the treatment, receptors and DNA are targeted	35
24	Pirampu	<i>Calamus rotang</i>	leaves	α -asarone β -asarone Acoradin Galangin Eugenol	HbA1C, LDL, VLDL, and blood sugar levels are all reduced by ethanol leaf extract	36
25	Peri winkle	<i>Catharanthus roseus</i>	Leaves	Vinblastine Vincristine Ajmalicine Catharanthine Vindoline	Through the GLUT 2 and 4 genes, the leaf juice extract has an effect on insulin secretion similar to that of the pancreas.	37
26	Sweet lemon	<i>Citrus limetta</i>	Fruit peel	Hesperidin Naringin Limonene Γ -terpinene	Reduces oxidative stress, demonstrating anti-diabetic effects	38
27	Yam	<i>Dioscorea alata</i>	Tuber	Rosmarinic acid	Low GI. It functions by a number of mechanisms, including reducing hyperglycaemia, inflammation, and mitochondrial dysfunction	39
28	Amla	<i>Emblica</i>	Fruit	Rutin	Amla chelates arsenic and reduces	40

		<i>officinalis</i>		Quercetin Catechol Gallic acid Ellagic acid Glauanol Sitosterol Tiglic acid Limolin Jangomolide Garuganin	ROS and cytokines	
29	Fig tree	<i>Ficus glomerata</i>	leaf		Restores uric acid, creatinine, cholesterol levels and also hyperalgesia	41
30	Coffee plum	<i>Flacourtia jangomas</i>	Leaves stem		Enhances a number of chemical characteristics	42
31	Garuga	<i>Garuga pinnata</i>	bark		Haemoglobin, blood glucose, cholesterol, and triglycerides are all reduced by extract	43
32	Bruhati	<i>Solanum indicum</i>	fruit	Solasodine Solasonine Solamargine	Demonstrates glucose tolerance and acts as an antioxidant, anti-inflammatory, and anti-hyperglycaemic	44
33	Soya bean	<i>Glycine max</i>	Leaf	Genistein Glycitin	Unsaturated fatty acids present in soya promotes blood glucose levels and also regulates Insulin resistance	45
34	Indian turnsole	<i>Heliotropium indicum</i>	Whole plant	Indicine Echinitine Supinene	Improves blood glucose levels in the circulation	46
35	Shoe flower/ Chinese hibiscus	<i>Hibiscus rosasinensis</i>	flower	Orientin Quercetin β -sitosterol	Used to treat acute and subacute phases of hyperglycemia	47
36	Cogon grass	<i>Imperata cylindrical</i>	rhizome	Arundoin Cylindrin Simiarenol	Reduces the amount of LDL, cholesterol, and related adverse effects	48
37	Sweet potato	<i>Ipomoea batatas</i>	peel	Citrusinine Chlorogenic acid	The sweet potato's peel includes a number of active constituents that are thought to have anti-diabetic properties	49
38	Physic nut	<i>Jatropha curcas</i>	root	Γ -cadinene α -epicadinol Pulegone	Improves parameters such as RBC, WBC, PCV, lymphocytes, neutrophils	50
39	Miracle leaf	<i>Kalanchoe pinnata</i>	leaves	Bryophillin A	In the early stages of the condition, it is used as an anti-diabetic	51
40	Henna	<i>Lawsonia inermis</i>	leaf	Lawson	The plant contains beneficial components that can be used to treat diabetes, such as polyphenols	52
41	Mango	<i>Mangifera indica</i>	leaves	Mangiferin β -carotene Campesterol	The leaf's phenolic chemicals have an anti-hyperglycaemic action	53
42	Banana	<i>Musa paradisiaca</i>	Leaves Fruit peel	Phytol Vitamin.E B-sitosterol Stigmasterol Hexa-decenoic acid	lowers IL-6 and TNF- α levels. Increases PPAR- γ and GLUT-4 levels	54
43	Black cumin	<i>Nigella sativa</i>	Seed oil	Nigellitimine Allo-oceminol	Shows hypoglycaemic effect with a daily intake of 1.5–3 ml of seed oil	55
44	Creeping wood sorell	<i>Oxalis corniculata</i>	leaves	Isovitexine	The disease is treated by increasing the extract's dose and exposure	56
45	Piyasal	<i>Pterocarpus marsupium</i>	Heart wood	Pterostilbene Epicatechin Pterosupin	lowers the levels of the inflammatory mediators- TNF- α and IL-6	57
46	Tulsi	<i>Ocimum sanctum</i>	leaves	Eugenol β -bisabolen Elemene Estragole Borneol	It reduces blood glucose levels when given at a dose of 2gm/kg. Additionally, leaves have the ability to scavenge radicals (superoxide, hydroxyl)	58

47	Cardamom	<i>Elettaria cardamomum</i>	Whole spice	1,8-cineole α -terpinyl acetate Sabinene β -linalool	Cardamom supplementation in T2DM patients reduces oxidative stress and hyperglycemia.	59
48	Gurmar/ Podaparthi	<i>Gymnema sylvestre</i>	leaves	Gymnemic acids α , β - chlorophylls	Stimulates the secretion of insulin from the pancreas. preventing the stomach from absorbing glucose	60
49	Ginseng	<i>Panax ginseng</i>	roots	Ginsenoside Dammarane	According to research conducted in vitro and in vivo, the ginsenosides found in plant roots have anti-diabetic properties	61
50	Bhuinamla	<i>Phyllanthus simplex</i>	Whole plant	Vanillin β -daucosterol Rutin	Since it lowers oxidative stress, it has potential for reducing secondary complications of DM	62
51	Mountain knot grass	<i>Aerva lanata</i>	leaves	Ervine Aervoside Kaempferol Quercetin	As an anti-diabetic drug, the phenolic component of leaves inhibits the actions of α -glucosidase and α -amylase.	63
52	Garden sage	<i>Salvia officinalis</i>	leaves	Thujone Eucalyptol Caryophyllene	Used as a dietary additive to safeguard pancreatic β -cells	64
53	Common Chick weed	<i>Stellaria media</i>	Whole herb	β -carotene Genistein Coumarin	The herbal tea derived from the plant is thought to function as an anti-diabetic drug via the STAT-3 pathway	65
54	Tamarind	<i>Tamarindus indica</i>	Fruit pulp	β -amyirin Compesterol β -sitosterol	It has good inhibitory activity against α -amylase & α -glucosidase and also regulates the negative consequences of diabetes.	66
55	Arjun tree	<i>Terminalia arjuna</i>	Stem bark	Arjunic acid Arjungenin Luteolin Pellargonidin	lowers the amounts of the enzymes needed to absorb glucose, such as hexokinase, G-6-P, and F-6-P	67
56	Black gram	<i>Vinga mungo</i>	seeds	Carbohydrate Proteins Minerals Vitamins	Regulates the activity of the antioxidant enzymes SOD, CAT, and GPx	68
57	Aswagnadha	<i>Withania somnifera</i>	Root leaf	Withanolide Withaferin A Viscosalactone B	Maintains the levels of substances including lipids, ASP, ALT, G-6-P	69
58	Ginger	<i>Zingiber officinale</i>	rhizome	Gingerol Zingerone Shogaol Citral	Ginger juice improves metabolism and lowers triglyceride and cholesterol levels	70
59	Indian Sundarban mangrove	<i>Rhizospora mucronata</i>	leaves	Rhizophorin – A Phomo-xanthone Lupeol Coumarin B-sitosterol	Improves the body's metabolism to treat diabetic individuals	71
60	Devil's cotton	<i>Abroma augusta</i>	leaves	A bromine Abromasterol Digitonide	The extract is employed as a preventive diabetic medication	72

Based on the condition, the following are some of the ways it works as an anti-diabetic agent:

- Insulin secretors and releasers.
- Insulin mimetic substances.
- Enhances metabolism.
- Blocks the action of the enzymes such as α -glucosidase and α -amylase.

- Lowers the levels of chemical indicators including cholesterol, triglycerides, AST, ALT, ACP.
- Pancreatic stimulators for the release of insulin.
- Supplementation of kitchen spices in case of emergency purpose.
- Radical scavenging activity.
- Lowers the levels of TNF- α , IL-6 (anti-inflammatory).
- Decreases the absorption of glucose from pancreas.
- Enhances serum parameters such as RBC, WBC, lymphocytes, neutrophils.
- Low glucose index to act as an anti-hyperglycaemic.

Future Prospects for Diabetes Mellitus Treatment: Traditional medical practices including Ayurveda, Homoeopathy, Siddha, and Unani offer a wide range of flora for treating various ailments around the world. Plants have gained significant relevance since Covid-19 and people are increasingly turning to natural remedies for positive effects. The therapeutic efficacy of plants is attributed to the presence of a variety of active phytoconstituents, including alkaloids, saponins, flavonoids, polyphenols, triterpenes, and sesquiterpenes. We discovered the future targets for the therapy of diabetes, which is to improve insulin resistance, by studying all the mechanisms, applications, and pathophysiology of the illness. As a result, drugs derived from either chemicals or plants ought to be able to lessen the resistance to insulin. Insilco studies on various anti-diabetic phytoconstituents are in progress.

CONCLUSION: In inference, diabetes is the most prevalent illness in the modern world. Various applications of medicinal plants for the treatment of diabetes are explained in this cumulative review.

CONFLICT OF INTEREST: No Conflict of interest

ACKNOWLEDGEMENTS: Thanks to Narayana Pharmacy College, Nellore, A.P, India.

REFERENCES:

1. Rohan Verma: Introduction to Diabetes. Diabetes Management 2022.
2. Muzeeb Z. Banday, Aga S. Sameer and Saniya Nissar: Pathophysiology of diabetes: An overview. Avicenna Journal of Medicine 2020; 10: 174-188.
3. Haikael D. Martin, Rose Kasole and Judith Kimiywe: Traditional Medicine and its role in the management of Diabetes Mellitus: "patients and herbalist's perspectives". Evidence-based complementary and Alternative Medicine 2019.
4. American Diabetes Association. Diagnosis & Classification of Diabetes Mellitus. Diabetes care 2019; 37: 81-90.
5. Saeedi P, Petersohn I, Salpea P, Malanda B, Karuranga S, Unwin N, Colagiuri S, Guariguata L, Motala AA, Ogurtsova K, Shaw JE, Bright D and Williams R: IDF Diabetes Atlas Committee. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. Diabetes Res Clin Pract 2019; 157: 107843
6. Is Type 2 diabetes genetic. Health line 2023.
7. Unai Galicia – Gracia, Asier Benito-Vicente, Shifa Jebari, Asier Larrea-Sebal, Haziq Siddiqi and Kepa B: Uribe, Helena Ostolaza, Cesar Martin: Pathophysiology of Type 2 Diabetes. International Journal of Molecular Sciences 2020; 21: 6275.
8. Amit Sapra and Priyanka Bhandari: Diabetes. Stat Pearls 2023.
9. Ansari P, Hannan JMA, Seidel V and Abdel-Wahab YHA: Polyphenol-Rich Leaf of *Annona squamosa* Stimulates Insulin Release from BRIN-BD11 Cells and Isolated Mouse Islets, reduces (CH₂O)_n Digestion and Absorption, and Improves Glucose Tolerance and GLP-1 (7-36) Levels in High-Fat-Fed Rats Metabolites 2022; 12: 995.
10. Drissi F, Lahfa F, Gonzalez T, Peiretti F, Tanti JF, Haddad M, Fabre N and Govers R: A *Citrullus colocynthis* fruit extract acutely enhances insulin-induced GLUT4 translocation and glucose uptake in adipocytes by increasing PKB phosphorylation. Journal of Ethnopharmacology 2021; 270: 113772.
11. Arezoo Afshari, Fouzieh Salimi, Azin Nowrouzi, Masoumeh Babaie Khalili, Salar Bakhtiyari, Gholameza Hassanzadeh, Maryam Shabani, Ali Ahadi and Masoumeh Farhadinezhad: Differential expression of gluconeogenic enzymes in early and late stage diabetes: the effect of *Citrullus colocynthis* (*L.*) *scharid*. Seed extract on hyperglycemia and hyperlipidemia in Wistar-Albino rat's model. Clinical Phyto Science 2021.
12. Hajleh MNA, Khleifat KM, Alqaraleh M, Al-Hraishat E, Al-Limoun MO, Qaralleh H and Al-Dujaili EAS: Antioxidant and Antihyperglycemic Effects of *Ephedra foeminea* Aqueous Extract in Streptozotocin-Induced Diabetic Rats. Nutrients 2022; 14: 2338.
13. Khouya T, Ramchoun M, Elbouny H, Hmidani A, Bouhlali EDT and Alem C: Loquat (*Eriobotrya japonica* (Thunb) Lindl.): Evaluation of nutritional value, polyphenol composition, antidiabetic effect, and toxicity of leaf aqueous extract. Journal of Ethnopharmacology 2022; 296: 115473.
14. Liu Y, Zhang W, Xu C and Li X: Biological Activities of Extracts from Loquat (*Eriobotrya japonica* Lindl.): A Review. International Journal of Molecular Sciences. 2016; 17(12): 1983.

15. Vora A, Varghese A, Kachwala Y, Bhaskar M, Laddha A, Jamal A and Yadav P: *Eugenia jambolana* extract reduces the systematic exposure of sitagliptin and improves conditions associated with diabetes: A pharmacokinetic and a pharmacodynamic herb-interaction study. *Journal of Traditional Complementary Medicine* 2018; 9: 364-71.
16. Kishalay Jana, Tushar Kanti Bera, Debidas Ghosh: Anti diabetic effects of *Eugenia jambolana* in the streptozotocin induced diabetic male albino rat. *Biomarkers and Genomic Medicine* 2015; 7: 116-124.
17. Wanlin Niu, Junjie Miao, Xuejia Li, Qian Guo, Zujun Deng and Lirong Wu: Metabolomics combined with systematic pharmacology reveals the therapeutic effects of *Salvia miltiorrhiza* and *Radix Pueraria lobata* herb pair on type 2 diabetic rats. *Journal of Functional Foods* 2022; 89: 104950.
18. Sushma Devi, Navneet Kaur, Manish Kumar, Praveen Kumar: *In-vitro* and *In-vivo* evaluation of anti-diabetic potential of *Euphoria nerifolia* in streptozotocin induced diabetes in rats and its *in-vitro* anti-oxidant studies. *Food Chemistry Advances* 2023; 100199.
19. Mansourzadeh S, Esmaeili F, Shabani L and Gharibi S: Trans-differentiation of mouse mesenchymal stem cells into pancreatic β -like cells by a traditional anti-diabetic medicinal herb *Medicago sativa* L. *Journal of Traditional Complementary Medicine* 2022; 12: 466-476.
20. Chang-hua Zhang, Jun-Qing Sheng, Surendra Sarsaiya, Fu-xing Shu, Tong-tong Liu, Xiu-ying Tu, Guang-qiang Ma, GuoLiang Xu, Hong-xiang Zheng and Li fen Zhou: The anti-diabetic activities, gut microbiota composition, the anti-inflammatory effects of *Scutellaria-coptis* herb couple against insulin resistance – model of diabetes involving the toll-like receptor 4 signalling pathway. *Journal of Ethnopharmacology* 2019; 237: 202-214.
21. Fariha Imtiaz, Muhammad Islam, Hamid Saeed, Arbar Ahmed, Maryam Asghar, Bushra Saleem, Muhammad Asim Farooq, Daulat Haleem Khan and Leena: peltonen: Novel phytoniosomes formulation of *Tradescantia pallida* leaves attenuates diabetes more effectively than pure extract. *Journal of Drug Delivery Science & Technology* 2023; 83: 104399.
22. Akram Ahangarpour, Maryam Mohammadian and Mahin Dainat: Anti Diabetic effect of hydroalcoholic *Urtica dioica* leaf extract in male rats with fructose induced insulin resistance. *Iranian Journal of Medical Sciences* 2012; 37: 181-186.
23. Maryam Keshvari, Masoud Rahmati, Rahim Mirnasouri and Farzaneh: Effects of endurance exercise and *Urtica dioica* on the functional, histological and molecular aspects of the hippocampus in STZ-induced diabetic rats. *Journal of Ethnopharmacology* 2021; 258: 112801.
24. Marine E. Baset, Tasneem I. Ali, Hanan Elshamy, Ahmed M. El Sadek, Diana G. Sami, Marwa T. Badawy, Saras. Abou-Zekry Hana H. Heiba, Mona K. Saadeldin and Ahmed Abderlatif: Anti-diabetic effects of fenugreek (*Trigonella foenum-graecum*): A comparison between oral and intraperitoneal administration – an animal study. *International Journal of Functional Nutrition* 2020.
25. Maede Ruyvaran, Alizamani, Alireza Mohamadian, Mohammed M. Zarshenas and Majid Nimrouzi: Safflower (*carthamus tinctorius* L.) oil could improve abdominal obesity, blood pressure and insulin resistance in patients with metabolic syndrome: A randomized, double-blind, placebo-controlled clinical trial. *Journal of Ethnopharmacology* 2022; 282: 114590.
26. Latifi E, Mohammadpour AA, Fathi Hafshejani B and Nourani H: *Ferulesasafoetida* oleo gum resin ethanolic extract alleviated the pancreatic changes and antioxidant status in streptozotocin-induced diabetic rats: A biochemical, histopathological, and ultrastructural study. *Journal of Food Biochemistry* 2022; 46: 14191.
27. Vikas Kumar, Danish Ahmed, Amita Verma, Firoz Anwar, Mohammad Ali and Mohd Muzeeb: Umbelliferone β -o-galactopyranoside from *Aegle marmelos* (L.) Corr. An ethnomedicinal plant with anti-diabetic, anti-hyperlipidemic and anti-oxidative activity. *BMC Complement Alternative Medicine* 2013.
28. Das S, Chaware S, Narkar N, Tilak AV, Raveendran S & Rane P: Antidiabetic activity of *Coriandrum sativum* in streptozotocin induced diabetic rats. *International Journal of Basic & Clinical Pharmacology* 2019; 8(5): 925–929.
29. Hye In Jeong, Soobin Jang and Kyeong Hankim: *Morus alba* L. for blood sugar management: A systematic and meta-analysis. *Plant-Derived Bioactive compounds as an anti-diabetic agent* 2022.
30. Haque MA, Hossain MS, Sayed NMA, Islam MT, Khan MR, Ahmmed F, Zohora FT, Agagunduz D, Ming LC and Capasso R: *Abelmoschus esculentus* (L.) Moench Pod Extract Revealed Antagonistic Effect against the Synergistic Antidiabetic Activity of Metformin and Acarbose upon Concomitant Administration in Glucose-Induced Hyperglycaemic Mice *Biologics* 2022; 2: 128-138.
31. Manas Ranjan Saha, Priyankar Dey, Indrani Sarkar, Dilip De Sarker, Biswajit Haldar, Tapas Kumar Chaudhuri and Arnab Sen: *Acacia nilotica* leaf improves insulin resistance and hyperglycemia associated acute hepatic injury and nephrotoxicity by improving systemic antioxidant status in diabetic mice. *Journal of Ethnopharmacology* 2018; 210: 275-286.
32. Anindita Banerjee, Siddhartha Singh, Shilpi Kumari Prasad, Sourav Kumar, Oly Banerjee, Tapan Seal, Sandip Mukherjee and Bithin Kumar Maji: Protective efficacy of *Tinospora sinensis* against streptozotocin induced pancreatic islet cell injuries of diabetic rats and its correlation to its phytochemical profiles. *Journal of Ethnopharmacology* 2020; 248: 112356.
33. Masood S, Rehman AU and Bashir S: Investigation of the anti-hyperglycaemic and antioxidant effects of wheat bread supplemented with onion peel extract and onion powder in diabetic rats. *Journal of Diabetes and Metabolic Disorders* 2021; 20: 485–495.
34. Hemmati M, Serki E and Gholami M: Effects of an ethanolic extract of *Berberis vulgaris* fruits on hyperglycaemia and related gene expression in streptozotocin-induced diabetic rats. *International Journal of Phytomedicine and Phytochemistry* 2017.
35. Yin K, Yang J, Wang F, Wang Z, Xiang P, Xie X, Sun J, He X and Zhang X: A preliminary study of the chemical composition and bioactivity of *Bombax ceiba* L. flower and its potential mechanism in treating type 2 diabetes mellitus using ultra-performance liquid chromatography quadrupole-time-flight mass spectrometry and network pharmacology analysis. *Frontiers Nutrition* 2022; 9: 1018733.
36. Kumar KV, Lalitha KG and Kumar RS: Antidiabetic and Antihyperlipidemic Effects of *Calamus rotang* L leaves (Arecaceae) in Streptozotocin-Nicotinamide Induced Diabetic Model. *Journal of Pharmaceutical Research* 2021; 20: 80–87.
37. Al-Shaqha WM, Khan M and Salam N: Anti-diabetic potential of *Catharanthus roseus* Linn. and its effect on the glucose transport gene (GLUT-2 and GLUT-4) in

- streptozotocin induced diabetic wistar rats. BMC Complementary and Alternative Medicine 2015; 15.
38. Lu K and Yip YM: Therapeutic Potential of Bioactive Flavonoids from *Citrus fruit* Peels toward Obesity and Diabetes Mellitus. *Future Pharmacology* 2023; 3(1): 14-37.
 39. Sri Nabawiyathi Nurul Makiyah, Masaki Kita, Ika Setyawati and Sri Tasminatun: *Dioscorea alata* L. Tubers improve diabetes through Anti-Hyperglycaemia, Anti-Inflammation, Ameliorate Insulin resistance and mitochondrial dysfunction. *Indonesian Biomedical Journal* 2022; 14: 365-75.
 40. Singh MK, Dwivedi S, Yadav SS, Yadav RS and Khattri S: Anti-diabetic Effect of *Embllica-officinalis* (Amla) Against Arsenic Induced Metabolic Disorder in Mice. *Indian Journal of Clinical Biochemistry* 2020; 35: 179-187.
 41. Shaikh A, Ibrahim M and Khan M: Effect of *Ficus glomerata* leaf extract in streptozotocin-induced early diabetic complications and its characterization by LC-MS. *EXCLI Journal* 2020; 19: 33-47.
 42. Singh AK and Singh J: Evaluation of anti-diabetic potential of leaves and stem of *Flacourtia jangomas* in streptozotocin-induced diabetic rats. *Indian Journal of Pharmacology* 2010; 42: 301-305.
 43. Shirwaikar A, Rajendran K and Barik R: Effect of aqueous bark extract of *Garuga pinnata Roxb.* in streptozotocin-nicotinamide induced type-II diabetes mellitus. *Journal of Ethnopharmacology* 2006; 107: 285-290.
 44. Gadewar MM, GK, P. Mishra, P.C. Ashraf, G.M. Almashjary, M.N. Harakeh, S. Upadhye, V. Dey, A. Singh, P. Jha NKL and Jha SK: Evaluation of Antidiabetic, Antioxidant and Anti-Hyperlipidemic Effects of *Solanum indicum* Fruit Extract in Streptozotocin-Induced Diabetic Rats. *Current Issues in Molecular Biology* 2023; 45: 903-917.
 45. Choi MS, Ryu R, Seo YR, Jeong TS, Shin DH, Park YB, Kim SR and Jung UJ: The beneficial effect of soybean (*Glycine max (L.) Merr.*) leaf extracts in adults with prediabetes: a randomized placebo-controlled trial. *Food & Functional Journal* 2014; 5: 1621-1630.
 46. Subhan Ali Mohammad, Shaik Abdul Nabi, Saritha Marella, Krishna Tilak Thandaiah, Malaka Venkateshwarulu Jyothi Kumar and Chippada Appa Rao: Phytochemical screening and antihyperglycemic activity of *Heliotropium indicum* whole plant in Streptozotocin induced diabetic rats. *Journal of Applied Pharmaceutical Science* 2014; 4:065-071.
 47. Luana Alves Freitas Afiune, Thaís Leal-Silva, Yuri Karen Sinzato and Rafaienne Queiroz Moraes-Souza: Beneficial effects of *Rosa sinensis* L. flower aqueous extract in pregnant rats with Diabetes. *PLOS ONE* 2017.
 48. Ranjana Kohli, Madan L Kaushik, Jai Parkash Kaidan and Bhupendra Chauhan: Anti diabetic activity of rhizome extracts of *Imperata cylindrical* against alloxan – induced Diabetes Mellitus in rats. *International Journal of Research in Pharmaceutical Sciences* 2020; 11: 2704-2709.
 49. Akhtar N, Akram M, Daniyal M and Ahmad S: Evaluation of antidiabetic activity of *Ipomoea batatas* L. extract in alloxan-induced diabetic rats. *International Journal of Immunopathological Pharmacology* 2018; 32: 2058738418814678.
 50. Aladodo RA, Muhammad NO & Balogun EA: Effects of Aqueous Root Extract of *Jatropha curcas* on Hyperglycaemic and Haematological Indices in Alloxan-induced Diabetic Rats. *Fountain Journal of Natural and Applied Sciences* 2013; 2(1): 52-58.
 51. Ana L. Aguero-Hernández, Catalina Rosales-Lopez, Cristina Herrera, Andres Vargas-Picado and Rodrigo Munoz and Ana Abdelnour-Esquivel: Hypoglycemic effect of *Kalanchoe pinnata (Lam) Pers.* Leaf Extract. *Pharmacognosy Journal* 2020; 12: 557-561.
 52. Widyawati T, Pane YS and Yusoff NA: Effect of *Lawsonia inermis* Linn. Extracts on Blood Glucose Level in Normal and Streptozotocin – Induced Diabetic Rats. *Pakistan Journal of Nutrition* 2019; 18: 671-676.
 53. Villas Boas GR, Rodrigues Lemos JM, de Oliveira MW, Dos Santos RC, Stefanello da Silveira AP, Barbieri Bacha F, Ito CNA, Bortolotte Cornelius E, Brioli Lima F, Sachilarid Rodrigues AM, Belmal Costa N, Francisco Bittencourt F, Freitas de Lima F, Meirelles Paes M, Gubert P and Oesterreich SA: Aqueous extract from *Mangifera indica* Linn. (Anacardiaceae) leaves exerts long-term hypoglycemic effect, increases insulin sensitivity and plasma insulin levels on diabetic Wistar rats. *PLoS One* 2020; 15: 0227105.
 54. Abdel Aziz SM, Ahmed OM, Abd El-Twab SM, Al-Muzafar HM, Amin KA and Abdel-Gabbar M: Antihyperglycemic Effects and Mode of Actions of *Musa paradisiaca* Leaf and Fruit Peel Hydroethanolic Extracts in Nicotinamide/Streptozotocin-Induced Diabetic Rats. *Evidence Based Complementary and Alternative Medicine* 2020; 9276343.
 55. Rachman PNR and Akrom E Darmawan: The efficacy of black cumin seed (*Nigella sativa*) oil and hypoglycemic drug combination to reduce HbA1c level in patients with metabolic syndrome risk. *IOP Conference Series: Materials Science and Engineering* 2017; 259: 012018.
 56. Dutta A, Lahkar M & Handique C: Evaluation of antidiabetic activity of *Oxalis corniculata* in streptozotocin induced diabetic rats. *International Journal of Basic & Clinical Pharmacology* 2017; 5: 2178–2183.
 57. Dhayaney Vijayan and Sibi G: *Pterocarpus marsupium* for the Treatment of Diabetes and Disorders. *Journal of Complementary Science and Alternative Health Care* 2019; 9(1): 555754
 58. Raiya Adiba Antora and Rabeta Mohd Salleh: Antihyperglycemic effect of *Ocimum* plants: A short review *Asian Pacific Journal of Tropical Biomedicine* 2017; 7(8): 755-759,
 59. Aghasi M, Ghazi-Zahedi S, Koohdani F, Siassi F, Nasli-Esfahani E, Keshavarz A, Qorbani M, Khoshamal H, Salari-Moghaddam A and Sotoudeh G: The effects of green cardamom supplementation on blood glucose, lipids profile, oxidative stress, sirtuin-1 and iris in in type 2 diabetic patients: a study protocol for a randomized placebo-controlled clinical trial. *BMC Complementary and Alternative Medicine* 2018; 18: 18.
 60. Sejal M. Karwa, Dattaprasad N. Vikhe and Ravindra S. Jadhav: A Review on Phytochemical Investigation of *Gymnema sylvestre* leaves. *Research Journal of Science and Technology* 2022; 14(4): 261-8.
 61. Chen W, Balan P and Popovich DG: Review of Ginseng Anti-Diabetic Studies. *Molecules* 2019; 24(24): 4501.
 62. Shabeer J, Srivastava RS and Singh SK: Antidiabetic and antioxidant effect of various fractions of *Phyllanthus simplex* in alloxan diabetic rats. *Journal of Ethnopharmacology* 2009; 124: 34-38.
 63. Akanji MA, Olukolu SO and Kazeem MI: Leaf Extracts of *Aerva lanata* Inhibit the Activities of Type 2 Diabetes-Related Enzymes and Possess Antioxidant Properties. *Oxidative Medicine and Cellular Longevity* 2018; 2018: 3439048.

64. Askhani-Esfahani S, Noorafshan A, Ebrahimi A, Bahmani – Jahromi M and Imanieh M: *Salvia officinalis* protects pancreatic beta cells against streptozotocin – induced damage; A Stereological study. Jundishapur Journal of Natural Pharmaceutical Products 2021; 17(1): 109906.
65. Demjan V, Soja A, Kiss T, Fejes A, Gausz FD, Szucs G, Siska A, Foldesi I, Tengolics R, Darula Z, Csupor D, Pipicz M and Csont T: *Stellaria media* tea protects against diabetes-induced cardiac dysfunction in rats without affecting glucose tolerance. Journal of Traditional and Complementary Medicine 2021; 12: 250-259.
66. Krishna RN, Anitha R and Ezhilarasan D: Aqueous extract of *Tamarindus indica* fruit pulp exhibits antihyperglycaemic activity. Avicenna Journal of Phytomedicine 2020; 10: 440-447.
67. Thomson HAJ, Ojo OO, Flatt PR and Abdel-Wahab YHA: Aqueous Bark Extracts of *Terminalia Arjuna* Stimulates Insulin Release, Enhances Insulin Action and Inhibits Starch Digestion and Protein Glycation *in-vitro*. Austin J Endocrinol Diabetes 2014; 1(1): 1001.
68. Mthiyane FT, Dlodla PV, Ziqubu K, Mthembu SXH, Muvhulawa N, Hlengwa N, Nkambule BB and Mazibuko-Mbeje SE: A Review on the Antidiabetic Properties of *Moringa oleifera* Extracts: Focusing on Oxidative Stress and Inflammation as Main Therapeutic Targets. Front Pharmacol 2022; 13: 940572.
69. Shah RA, Khan S, Vakil M and Quereshi MT: Evaluation of Anti-Diabetic potential of *Withania Somnifera* by Glucose uptake assay on 3T3 F442A Fibroblast (Adipocyte cells). International Journal of Pharmaceutical Sciences and Research 2017; 8(1): 120-25.
70. Nurinda E, Kusumawardani N, Wulandari AS, Fatmawati A, Emelda E, Nisa H, Hasan NA, Iriyanti WF, Rohmah M, Lestari P and Aprilia V: Pharmacological Study: Synergistic Antidiabetic Activity of Cinnamon Bark and Zingiber Extract in Streptozotocin-Induced Diabetic Rats. Open Access Maced J Med Sci 2022; 10(8): 1-6.
71. Sachithanandam V, Lalitha P, Parthiban A, Mageswaran T, Manmadhan K and Sridhar RA: Review on Antidiabetic Properties of Indian Mangrove Plants with Reference to Island Ecosystem. Evidence Based Complementary Alternative Medicine 2019; 5: 430514.
72. Khanra R, Dewanjee S, K Dua T, Sahu R, Gangopadhyay M, De Feo V and Zia-Ul-Haq M: *Abroma augusta* L. (Malvaceae) leaf extract attenuates diabetes induced nephropathy and cardiomyopathy *via* inhibition of oxidative stress and inflammatory response. Journal of Translational Medicine 2015.

How to cite this article:

Pasumarthy SM and Guttikonda U: A cumulative review on therapeutic flora used to treat diabetes mellitus. Int J Pharm Sci & Res 2024; 15(3): 626-35. doi: 10.13040/IJPSR.0975-8232.15(3).626-35.

All © 2024 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)