



Received 24 February, 2010; received in revised form 09 April, 2010; accepted 25 April, 2010

EVALUATION OF ANTIOXIDANT, WOUND HEALING AND ANTI- INFLMMATORY ACTIVITY OF ETHANOLIC EXTRACT OF LEAVES OF FICUS RELIGIOSA

Rita M. Charde*¹, Hemant J. Dhongade¹, Manoj S. Charde² and A.V.Kasture³

J. L. Chaturvedi College of Pharmacy ¹, Electronic Zone Building, MIDC, Nagpur, Maharashtra, India

Kamala Nehru College of Pharmacy ², Buttibori, Nagpur, Maharashtra, India

Department of Pharmaceutical Sciences, Rashtrasant Tukadoji Maharaj Nagpur University ³, Nagpur, Maharashtra, India

Keywords:

Ficus religiosa,
antioxidant,
wound healing,
anti- inflammatory

ABSTRACT

The present study deals with evaluation of antioxidant, wound healing and anti-inflammatory activity of ethanolic extract of leaves of *Ficus religiosa*. The ethanolic extract prepared by maceration technique was subjected to screen for antioxidant activity using DPPH radical scavenging method and wound healing activity using incision, excision, histopathological and dead space wound model and the study was supported with evaluation of granuloma tissue to estimate hydroxyproline content and histopathological evaluation. The anti-inflammatory study was carried out by using carageenan induced rat paw odema method. The tested extract of different dilutions in range 200µg/ml to 1000 µg/ml shows activity in range of 6.34% to 13.35%. Significant increase in wound closure rate, skin breaking strength, granuloma breaking strength was observed. The hydroxyproline content was also increased with decrease in scar area. The initial healing action might be due to increased collagen deposition and better alignment, with the obtained results it can be concluded that *Ficus religiosa* extract has significant wound healing activity and initial healing may be due to presence of glycosides and tannins. The extract shows prominent anti-inflammatory activity as compared to that of standard (Ibuprofen gel). The extract shows good anti-inflammatory activity on carageenan induced rat paw odema method.

Correspondence to author:

Rita M. Charde

J. L. Chaturvedi College of Pharmacy,

Electronic Zone Building,
MIDC, Nagpur, Maharashtra,
India

E-Mail:
manojudps@rediffmail.com

INTRODUCTION: *Ficus religiosa* (Moreaceae), commonly known as peepal tree is distributed throughout India, Pakistan, China and other tropical countries. The bark is reported to possess wound healing activity¹, anti-inflammatory, analgesic, anti-lipid peroxidation activity², and tender shoots have purgative properties³. The leaf extract shows the presence of glycosides and tannins from the phytochemical analysis⁴.

Wounds are visible results of individual cell death or damage and can be classified by site, size, depth and causation – surgery, accident or circulatory failure⁵. Wound healing is a process which is fundamentally a connective tissue response. Initial stage of this process involves an acute inflammatory phases followed by synthesis of collagen and other extracellular macromolecules which are latter remolded to form scars^{6, 7}. Inflammation is the process which may be due to release of histamine, kinins, serotonin and prostaglandin⁸. Anti-inflammatory agents are the agents which normally inhibit the release of these inflammatory mediators. The present study deals with the use of traditionally claimed use of *Ficus religiosa*. The invitro antioxidant activity was screened by DPPH method. The wound healing study was evaluated by incision and excision and dead space wound model followed by histopathological study. Anti-inflammatory study was screened by carrageenan induced paw edema method.

MATERIAL AND METHODS:

Plant material and Preparation of Herbal

Extract: *Ficus religiosa* fresh leaves were collected from the area of university campus, and authenticated in Department of Botany, Rashtrasant Tukdoji Maharaj Nagpur University Campus, Nagpur. Initially these leaves were washed with fresh water to remove adhering dirt and foreign particles and dried at 35 - 40°C in an oven. The dried leaves were crushed and grinded to get powder and weighed. The weighed powder was then placed with ethanolic solution in a cylinder. 500g of *Ficus religiosa* powder in 1.0 liter of ethanolic solution were macerated for 7 days. The mensturm was removed and concentrated by vaccum distillation. Again the crude material was allowed to undergo maceration for 4 days followed by 2 days for complete extraction. The mensturm was collected and concentrated by vaccum distillation and then air dried in an evaporating dish till constant weight was obtained. The percent yield of *Ficus religiosa* leaf extract is 26.52 %.

Animals: In the present study male Wistar rats (150-200g) were used for the study. They were individually housed and maintained on normal standard diet (Gold Muhor Brand, Lipton India limited) and water *ad libitum*. Temperature was maintained at 23±1°C with 12hr light and dark cycle.

Antioxidant activity: The DPPH (1, 1-diphenyl- 2- picryl hydrazyl) method was used for the determination of in vitro

antioxidant activity for crude extract. DPPH is stable free radicals scavenge by antioxidant present in test solution. To 1.0 mL of 0.01 mM Solution of DPPH, 1.0mL of ethanolic solution and 0.95 mL of tris HCl buffer of 0.05 M was added. To this solution 50 µl of extract of specific strength was added. It was kept for 20 min for reaction at room temperature and then absorbance was recorded at 517 nm. Control solution was also carried out without extract. Percent scavenging of DPPH radical was calculated by comparing absorbance between the test and diluted control mixture.⁹

Wound Healing activity: Animals were wounded under light ether anesthesia, semi- aseptically. The animals were assigned in to three groups, each group containing six animals. First group was untreated group which was taken as control (Group- 1). In Second group (Group-2) wounds received topical application of *Ficus religiosa* extract, while animals in Group- 3 (Reference Standard) received treatment of Framycetine Sulphate Cream (FSC) in both excision and incision wound models. No other topical or systemic therapy was given to animals during the course of this study. The experimental protocols were approved by the Institute Animal Ethical Committee.

Excision wounds¹⁰: Hairs were removed from dorsal thoracic central region of anaesthetized rats. Full thickness from the demarked area was excised to produce wound measuring around 300 mm² as shown in Fig 1. Wound was cleaned with cotton swab soaked in alcohol. The test

extract and FSC cream was applied on the excised wound once daily for 24 days starting from the first day of wounding. Wound contraction was measured as percent reduction in wound area¹¹.



Fig. 1: Excision Wound Model

Resutured Incision wounds: Animals were anaesthetized and paravertebral incisions (2.5 -3.0 cm long) were made through the entire length of skin. After the incision was made, the parted skin was kept together and stitched with nylon thread at 0.5 cm apart with curved needle (No. 11).The Photographic representation of incision wound model with the stitches on the incised skin are shown in Fig 2. The sutures were removed on day 8 and healing (tensile) strength was measured on day 10¹².



FIG. 2: Incision Wound Model

Dead space wound model: Under light ether anesthesia, subcutaneous dead space wounds were inflicted in the region of axilla and groin, by making a pouch through a small nick in the skin. Granuloma formation was induced by implanting either sterile cotton pellets or grass piths. Two sterile cotton pellets weighing 10 mg (sterilized by autoclaving) were implanted in axilla by technique of D'Arcy et al as describe by Turner¹³, but the granuloma were removed on 10th day. The sutures were mobbed with an alcoholic swab and animals were placed into their individual cages after recovery from anesthesia. Physical, mechanical and histopathological changes in granuloma tissue were studied in this model. Excision of granuloma from the surrounding tissues were performed on the 10th post wounding day under light ether anesthesia cotton pellet granuloma excised from dead space wound were dried overnight at 60°C so as to obtained a constant dry weight, their weights were noted and expressed as mg/100 gm body weight¹⁴. The excised tissue was cut into two approximately equal halves. One half of the granuloma tissue was used for determination of hydroxyproline content¹⁵. The other part is kept in 10% formalin solution for histopathological studies to evaluate the effect of extract on collagen formation.

Histopathological study: The histopathology study was done to evaluate the healing promoters like keratinization, epithelization, collagenation, fibrosis and neovascularization, which were evidenced

to promote wound healing in the experimental animals. The section from 10 day old regenerated tissue of incision wound was taken and washed with normal saline solution to study the healing markers. After fixing the section in 10% formalin solution the tissues were dehydrated with 90% ethanol, cut into thin sliced section (7 µm thick), stained with haemotoxyline- eosin dye and observed under light microscope for keratinization, epithelization, fibrosis, collagen formation and neovascularization¹⁶. The interpretation of the results were numbered from 1 to 5 of which 5 stands for maximum similarity and 1 stands for least similarity form normal tissue around the wound area in the *Ficus religiosa* treated and untreated wounds¹⁷.

Anti-inflammatory Study: Anti-inflammatory activity of *Ficus religiosa* leaf extract was studied by carrageenan induced paw edema method. The animals were divided into three groups containing six animals in each group. Group- 1 (control) untreated group, Group- 2 topical application with *Ficus religiosa* extract, Group 3- topical application of Ibuprofen gel. Group 2 and 3 received topical application of *Ficus religiosa* extract and Ibuprofen gel for comparison of anti-inflammatory activity. One hour after the application of *Ficus religiosa* extract 0.1 ml of carrageenan (1%) was injected into sub planar region of hind paw of rat. Measurement of paw volume (ml) were made by mercury displacing techniques using plethysmometer immediately before and 1, 2, 3 and 4 hr

after carrageenan injection. Percentage inhibition of inflammation after 1, 2, 3 and 4 hr was calculated by newbould's method¹⁸.

Statistical Analysis: Statistical difference between the groups were evaluated using one way analysis of variance (ANOVA) followed by Turkey's Kramer Multiple comparison test ($P < 0.001$).

RESULT: The present study deals with evaluation of antioxidant, wound healing and anti-inflammatory activity of ethanolic extract of *Ficus religiosa* leaves. The method used to determine the peroxide value or scavenging activity is based on decomposition or scavenging of DPPH which is a very stable free radical. The comparative percent peroxide value of *Ficus religiosa* extract is shown in Fig. 3.




FIG. 3: Percent Peroxide Value for *Ficus Religiosa* Leaf Extract

Percent peroxide value for *Ficus religiosa* extract is in the range of 6.34% to 13.35% for the extract with 200 $\mu\text{g/ml}$ to 1000 $\mu\text{g/ml}$ strength extract. The wound healing study was screened by four models incision, excision; dead space wound model and histopathological study. The anti-inflammatory study was

performed by carrageenan induced rat paw edema method. The acute toxicity study of ethanolic extract of *Ficus religiosa* extract do not show any signs of toxicity up to 3 g/kg body weight. Since there was no mortality at higher dose 1/10th of maximum dose of extract tested for acute toxicity was screened for evaluation of wound healing activity i.e., 300 mg/kg. In the excision wound study the wounds are treated with FSC cream and *Ficus religiosa* extract shows complete healing in day 24 - 25. The results of these groups were compared with the healing activity of untreated group which took more than 30 days for wound closure and fall of eschar. The results are shown in Table 1.

In incision wound study the tensile strength was measured on day 10 of regenerated tissues. The tensile strength of wounds treated with FSC cream and *Ficus religiosa* extract is 342.21 ± 3.55 and 379.00 ± 6.32 respectively was compared with mean tensile strength \pm SEM of untreated group i.e., 281.30 ± 5.82 . The comparative tensile strength is shown in Fig. 4.




FIG. 4: Effect of *Ficus Religiosa* Leaf on Tensile Strength

TABLE- 1: EFFECT OF TOPICAL APPLICATION OF *FICUS RELIGIOSA* LEAF EXTRACT ON EXCISION WOUND

Post wounding Days	Wounding Area (mm ²)			F Values
	Group 1	Group 2	Group 3	
0	311.86±3.03 (0)	302.31±1.818 (0)	369.50±4.38 (0)	F (2,15) =124.89
3	272.51±5.06 (12.62)	264.37±1.294 (12.54)	308.61±1.68 (16.47)	F (2,15) =55.180
6	212.29±4.86 (27.76)	236.71±4.736 (20.21)	245.89±2.04 (33.45)	F (2,15) =6.357
9	184.11±4.28 (41.01)	220.49±1.313 (31.25)	217.23±4.62 (41.20)	F (2,15) = 29.320
12	160.72±4.49 (46.54)	173.91±1.437 (42.47)	157.87±3.94 (57.27)	F (2,15) =5.821
15	135.81±3.76 (56.45)	104.79±1.267 (63.33)	129.09±3.79 (65.06)	F (2,15) =26.454
18	126.46±3.04 (59.45)	73.09±1.379 (75.82)	92.63±2.86 (74.93)	F (2,15)=112.90
21	114.44±5.71 (63.30)	35.69±1.411 (88.19)	44.56±3.24 (87.94)	F (2,15)= 123.68
24	97.26±3.58 (68.81)	9.258±0.467 (96.94)	13.85±1.32 (96.25)	F (2,15)= 498.14
27	66.69±2.57 (78.61)	9.258±0.467 (96.94)	0.00 (100)	F (2,15)= 671.29


Values are mean ± SEM of animals (n=6) in each group. Number in parenthesis indicates percentage of wound contraction. All the values are significant at p<0.001 as compared to Group 1

In histopathological study some healing markers were evaluated like keratinization, epithelization, fibrosis, Collagenation and neovascularization. After the histopathological evaluation of slides the *Ficus religiosa* extract treated wound shows prominent promotion of keratinization, epithelization, fibrosis and collagen formation, the results are compared with untreated group and wounds treated with FSC cream was considered as reference standard. The comparative promotion of these markers are shown with the help of photomicrographs in Fig. 5 (a), (b) and (c). The promotion of healing markers is four times more than that of untreated wound. The results are shown in Table 2.


Similarly in dead space wound model there was significant increase in granuloma breaking strength, dry granuloma weight and hydroxyproline content of granulation tissues as compared to controls. The results are shown in Table 4.

The topical anti-inflammatory activity of *Ficus religiosa* extract was carried out. The percentage protection (inhibition) of edema for *Ficus religiosa* extract and Ibuprofen gel was found to be 46.82 (1hr), 21.30 (2hr), 23.31 (3hr), 59.44 (4hr) and 36.54 (1hr), 64.85 (2hr), 63.83 (3hr), 67.62 (4hr) respectively. The tested extract showed significant anti-inflammatory activity when the results are compared with untreated group and


decreased in inflammation by Ibuprofen gel was taken as reference standard. The results are highly significant ($p < 0.001$) and are shown in Table 3.



(A)



(B)



(C)

FIG. 5: HISTOPATHOLOGY OF 10 D OLD REGENERATED TISSUE

Photomicrographs of histology of regenerated tissues collected from a) Untreated control, b) *Ficus religiosa* treated wounds, and c) FSC (1% w/w) treated after 10 d. Magnification si 100 X. K - Keratinization, E - Epithelization, F- Fibrous tissue and N- Neovascularization

DISCUSSION: A herbal drug for the treatment of chronic diseases or as raw material from which more or less complex chemical compounds with particular biological activity are isolated, the deeply held beliefs in the efficiency of natural products/herbals should be proven by scientific investigation bringing the tradition and experience in the ancient knowledge with in the realm of sciences¹⁹.

In excision wound healing study from the observed values it were assumed that *Ficus religiosa* extract shows better and faster healing as compared to untreated group. During the initiation of the study day "0" the wound closure with *Ficus religiosa* extract was very slow as compared to untreated and FSC treated wounds. As the study progress the wound healing efficiency increases shows complete healing on day 18-21 which is same for group 3 (wounds treated with FSC cream), But the untreated group took more time to complete wound closure. The study was carried out till fall of eschar leaving no scar behind. This shows healing potential of *Ficus religiosa* extract with better and faster wound closure.

The result obtained for tensile strength shows higher tensile strength for *Ficus religiosa* extract treated wounds as compared to wounds of untreated group. The increase in tensile strength may be due to promotion of collagen formation which significantly contributing to effective and better wound healing.

TABLE- 2: HISTOPATHOLOGICAL EXAMINATION OF WOUNDS TREATED WITH *FICUS RELIGIOSA* LEAF EXTRACT AT THE END OF 10 DAYS

Parameters	Group 1	Group 2	Group 3	F values
Keratinization	0.48±0.03	3.90±0.081	3.66±1.22	F(2,15) = 481.06
Epithelization	1.23±0.55	3.91±0.07	4.16±0.04	F(2,15) = 755.58
Fibrosis	2.01±0.07	3.76±0.07	4.20±0.05	F(2,15) = 260.49
Collagenation	1.85±0.42	3.80±0.08	4.43±0.05	F(2,15) = 468.40
Neovascularization	0.50±0.06	3.68±0.06	4.42±0.12	F(2,15) = 563.88

Values are reading of mean ± SEM of six animals, Value 5 refers to maximum similarity and 0 refers to least similarity of wound from normal tissues. All the values are significant at $p < 0.001$. Group 1 – Untreated group, Group 2 – Treated with *Ficus religiosa* extract, Group 3 – Treated with framycetine sulphate cream

TABLE 3: TOPICAL ANTI-INFLAMMATORY ACTIVITY OF *FICUS RELIGIOSA* LEAF EXTRACT

Group	Dose	Percent inhibition of paw volume at different time intervals			
		1hr	2hr	3hr	4hr
Standard (Ibuprofen gel)	300	36.54	64.85	63.83	67.62
Test (<i>Ficus religiosa</i> extract)	300	46.82	21.30	23.31	9.44

TABLE 4: EFFECT OF *FICUS RELIGIOSA* LEAF EXTRACT ON DEAD SPACE WOUND MODEL

Dead Space Wound Models			
Parameters	Granuloma Breaking Strength	Dry Granuloma weight (mg %	Hydroxyproline
Content	(g)	of B.W.)	Studied
Control	278.89±2.603	43.08±2.381	6.89±0.4604
Ethanol extract	202.85±1.032	50.23±1.125	7.29±1.325
F values	1.425	1.230	1.658

The *Ficus religiosa* extract also promotes healing markers as compared to that of control group. In addition to this increase in granuloma breaking strength, dry granuloma weight and hydroxyproline content strongly emphasize the positive wound healing potential of ethanolic extract of *Ficus religiosa*. Accordingly hydroxyl proline (marker of collagen) was significantly increased in treated group as

compared to control and further histopathological studies of granulation tissue, which recorded increase in collagen content in treated group.

The determination of anti-inflammatory activity is based on plethysmographic measurement of edema produce by sub planer injection of carrageenan in hind paw of rat. The

increase edema in animal treated standard (Ibuprofen gel) and *Ficus religiosa* extract were composed with increase in edema of untreated control animals at constant interval of 1, 2, 3 and 4hrs. The percentage inhibition of odema at known interval in treated animals was used for the purpose of calculating the percent inhibition of edema of control. The present study revealed that the *Ficus religiosa* extract showed significant inhibition of edema. The maximum activity showed during 1st and 2nd hrs, the results are highly significant ($p < 0.001$) as compared to standard. The anti-inflammatory activity may be due to inhibition of release of histamine, serotonin and kinens in first hour after injection of carrageenan and also retard the release of prostaglandin and like substances²⁰, shows anti-inflammatory activity of *Ficus religiosa* extract.

It has been reported that tannins possess ability to increase the collagen content, which is one of the factor for promotion of wound healing²¹. As the leaves are rich in tannins, it may be responsible for wound healing activity. The extract also shows antioxidant activity i.e., free radical scavenging activity which might reduces lipid peroxidation, may not only prevent or slows down the onset of necrosis but also improve vascularity²². The experimental plant also shows antimicrobial activity as reported earlier these are the factors count for wound healing activity of *Ficus religiosa* extract.

Thus it may be concluded that the leaves of *Ficus religiosa* shows significant

antioxidant, wound healing and anti-inflammatory activity. Further studies are in progress to isolate the bioactive component of plant extract.

REFERENCES:

1. Choudhary GP: Evaluation of ethanolic extract of *Ficus religiosa* bark on incision and excision wounds in rats. *Planta Indica* 2006; 2(3),17-9.
2. Sreelekshmi R, Latha PG, Arafat MM, Shyamal S, Shine VJ, Anuja GI, Suja SR and Rajasekharan S: Anti-inflammatory, analgesic and anti-lipid peroxidation studies on stem bark of *Ficus religiosa* Linn. *Natural Product Radiance* 2007; 6(5), 377-81.
3. Warriar PK, Nambiar VPK, Ramankutty C: *Indian Medicinal Plants. Vol-1, Madras: Orient Longman Ltd; 1995, 38.*
4. Kalyon Roy, H. Shivakumar, Sibaji Sarkar: Wound Healing Potential of Leaf Extracts of *Ficus Religiosa* on Wistar albino strain rats. *International Journal of Pharm Tech Research*, Vol.1, No.3, pp 506-508.
5. Prasad D. Rao CM: Wound healing activity of cyperus rotundus linn. *Indian J Pharm Sci* 1995; 33:845.
6. Chitra P. Sajithal GB: Influence of Aloe vera on Collagen turnover in healing of dermal wounds in rats. *Indian J Exp Biology* 1998; 36:896.
7. Jaswanth A. Akilandeswari LV. Rukmani: Wound healing activity of Aegle marmelos. *Indian J Pharm Sci* 2001; 63:41.
8. Rang HP. Text book of pharmacology, International Student edition, Churchill living stone 1995: 246.
9. Rekka E. Kourounakis PN: Effect of hydroxyethyl rutosides and related compounds on lipid peroxidation and free radical scavenging activity Some structural aspects. *J Pharm Pharmacol* 1991; 43:486.
10. Morton JJP. Melon HM: *Archives Internationales de Pharmacodynamic et de Therapie* 1972; 176:117.
11. Suguna L. Siva Kumar P. Chandrakaran G : Effect of *Centella asiatica* extract in dermal wound healing in rats. *Indian J Exp Biology* 1996; 34:1208-1211.
12. Lee KH: Studies on mechanism of action of salycilates II, Effect of vitamin A on wound healing. *J. Pharmaceutical Sciences* 1968; 1238-1240.
13. Ehrlich HP, Hunt TK: The effects of cortisone and anabolic steroids on the tensile strength of healing wounds. *Ann Surg* 1969; 170:203-6.

14. Dipasquale G, Meli A: Effect of body weight changes on the formation of cotton pellet-induced granuloma. *J Pharm Pharmacol* 1965; 17:379-82..
15. Woessner JF: The determination of hydroxyproline in tissue samples. *Arch Biochem* 1961; 93:440. .
16. Yeo JH. Lee KW. Kim HC. Lyuny K: *Biology and Pharmaceutical Bulletin* 2000; 23:1220-1223.
17. Rao GV. Shiv Kuamr HG Parathasa RG: *Indian J Pharamcology* 1996; 28:249-253.
18. Thomas CA, Rama Sarma GV: Analgesic and anti-inflammatory activities of *Melia dubia* bark. *Indian Drugs* 1999; 6:203-5.
19. Duke JA. Bogen Schutz-Godwain M. Ducellier JD: *Handbook of Medicinal herbs* CRC press, London 2002.
20. Thripathy KP. Rajasekaran A: *Indian Drugs* 2001; 38:426.
21. Tenni R. J: *Biochem* 1988; 37:169-177
22. Sunitha S, Nagaraj M, Varalakshmi P: Hepatoprotective effect of lupeol and lupeol linoleate on tissue antioxidant defence system in cadmium-induced hepatotoxicity in rats. *Fitoterapia* 2001, 72(5): 516-523.3.