



Received on 31 January, 2010; received in revised form 18 March, 2011; accepted 06 April, 2011

SIMULTANEOUS DETERMINATION OF SALBUTAMOL AND AMBROXOL IN FIXED DOSE COMBINATION BY SPECTROPHOTOMETRY

Priyanka A. Patel, Manjusha N. Dole*, Sanjay D. Sawant, Priyanka S. Shedpure

Department of Pharmaceutical Chemistry, Smt. Kashibai Navale College of Pharmacy, Kondhwa, Pune, Maharashtra, India

ABSTRACT

Keywords:

Salbutamol sulphate,
Ambroxol hydrochloride,
Absorbance correction method,
First order derivative method

Correspondence to Author:

Manjusha N. Dole

Asst. Prof., Department of
Pharmaceutical Chemistry, Smt.
Kashibai Navale College of
Pharmacy, Kondhwa, Pune,
Maharashtra, India

Salbutamol (SAL) and Ambroxol (AMB) is used for the treatment of asthma and bronchitis. Two simple, economical, accurate, and precise methods for simultaneous estimation of Salbutamol Sulphate (SAL) and Ambroxol Hydrochloride (AMB) in tablet dosage form have been developed. The methods are absorbance correction method (I) and first order derivative method (II). In first method ambroxol hydrochloride (AMB) was determined directly from calibration plot by measuring absorbance at 300nm and salbutamol sulphate (SAL) was determined after correction for absorbance of ambroxol hydrochloride (AMB) at 223 nm. Second method is based on first order derivative spectroscopy to overcome the spectral interference from the other drug. Wavelength 252 nm was selected for SAL and 232 nm for AMB. Both the methods showed linearity in the concentration range of 2-10 μ g/ml for SAL and 2-20 μ g/ml for AMB. The accuracy and precision of the methods were determined and the methods validated statically. No significant difference was observed between the results obtained by the two methods.

INTRODUCTION: Salbutamol sulphate (SAL), chemically known as bis [(1RS)-2-[(1, 1-dimethylethyl) amino]-1-[4-hydroxy-3-(hydroxymethyl) phenyl] ethanol] sulphate, is beta-adrenoceptor agonist used for the relief of broncho-spasm in conditions such as asthma and chronic obstructive pulmonary disease. The drug is official in Indian pharmacopoeia¹⁻⁴. Ambroxol hydrochloride (AMB) is chemically, trans-4-((2-amino-3, 5-dibromobenzyl) amino) cyclohexanol hydrochloride. Ambroxol reduces bronchial hyper-reactivity and acts as a mucolytic and cough suppressant. Combination of SAL and AMB is used for the treatment of asthma and bronchitis⁵⁻⁷

Literature survey reveals that salbutamol in combination with other drugs has been estimated by UV spectrophotometric methods⁸⁻¹², RP-HPLC methods¹³⁻¹⁴, TLC method¹⁵. For simultaneous determination of Ambroxol in combination with other drugs, UV spectrophotometric methods^{8, 16-20}, RP-HPLC²⁰⁻²⁴, HPTLC²⁵ and LC-MS/MS²⁶ are reported. Only one spectrophotometric method has been reported for the simultaneous estimation of salbutamol and ambroxol in combination⁸. Therefore, in the present work successful attempt has been made to estimate both the drugs simultaneously by two simple UV spectrophotometric methods i.e., absorbance correction method and first order derivative method. The proposed methods were optimized and validated as per ICH guidelines.




FIG. 1: SALBUTAMOL (4-[2-(TERT-BUTYLAMINO)-1-HYDROXYETHYL]-2-(HYDROXYMETHYL) PHENOL)




FIG 2: AMBROXOL HYDROCHLORIDE (TRANS-4-((2-AMINO-3,5-DIBROMOBENZYL) AMINO) CYCLOHEXANOL)

MATERIAL AND METHODS:

Instrumentation: For the present study JASCO double beam UV/Visible spectrophotometer (Model V-630) was used with slit width fixed at 1.5nm, equipped with spectra manager software (Version 1.5). Pair of 1-cm matched quartz cells were used to measure the absorbance of solution. The samples were weighed on electronic analytical balance (Contech Model CB-50) (table 1).

TABLE 1: INSTRUMENTS

Name	Model	Manufacture
UV/Visible spectrophotometer	V-630	Jasco
Electronic analytical balance	CB-50	Contech

Materials: Gift samples of Salbutamol sulphate and Ambroxol hydrochloride were provided by Glenmark Pharmaceuticals Limited, Nasik, India. The pharmaceutical dosage form used in this study was Sal Mucolite tablets (Cheminnova Remedies Pvt. Ltd). Each uncoated tablet contains 2mg SAL and 30mg AMB.

Solvent: Methanol Spectroscopic grade (Thomas Baker)

Preparation of stock solutions: Standard stock solutions of both Salbutamol sulphate and Ambroxol hydrochloride were prepared by dissolving 10 mg of SAL and 10mg of AMB separately in 20ml of 0.1N HCL in 100ml volumetric flasks. Final volume was made up to 100ml with 0.1N HCL to get working standard solution of each containing 100µg/ml of both SAL and AMB.

Methods:

Absorbance correction method (Method 1): This method involves absorbance correction for SAL determination by subtracting absorbance of AMB from total absorbance of sample at 223nm (λ_{max} of SAL). AMB concentration was determined directly from calibration plot by measuring absorbance at 300nm where SAL shows zero absorbance. The equations obtained for the determination of concentration are,

$$C_{AMB} = A_{300nm}/ax_1 \dots\dots\dots [I]$$

$$C_{SAL} = A_{223nm} - (ax_2 * C_{AMB})/ay_2 \dots\dots\dots [II]$$

A_{300nm} and A_{223nm} = Absorbance of sample at 300nm and 223nm

C_{AMB} and C_{SAL} = Concentrations of AMB and SAL in sample matrix.

ax_1 and ax_2 = Absorptivities of AMB at 300 nm and 223 nm

ay_1 and ay_2 = Absorptivities of SAL at 300 nm and 223 nm

TABLE 2: ABSORPTIVITY VALUES OF SAL AND AMB

Drug	Absorptivity at 300nm	Absorptivity at 223nm
SAL	0.000 (ay_1)	0.0388(ay_2)
AMB	0.008975(ax_1)	0.03865(ax_2)




FIG 3: OVERLAY SPECTRA OF SAL AND AMB FOR ABSORBANCE CORRECTION METHOD

First Order Derivative method (Method II): This method is based on first order derivative spectroscopy to overcome the spectral interference from other drug. Zero order spectrums of both the drugs are converted to first order derivative spectra with the help of spectra manager software of the instrument (**Figure 4**).

It was observed that SAL showed $dA/d\lambda$ zero at 252nm in contrast to AMB that has considerable $dA/d\lambda$ at this wavelength. Further, AMB has $dA/d\lambda$ zero at 232nm while at this wavelength SAL has significant $dA/d\lambda$. Therefore, wavelengths 252nm and 232nm were employed for determination of AMB and SAL respectively without interference of other drug. The calibration curves were plotted at these wavelengths of concentrations against $dA/d\lambda$ separately. The equation of line obtained to determine concentrations of SAL and AMB is as follows;

$$C_{SAL} = dA/d\lambda_{252nm} - 0.000356/0.0035 \dots\dots\dots [III]$$

$$C_{AMB} = dA/d\lambda_{232nm} - (- 0.00046)/0.00195 \dots\dots\dots [IV]$$




FIG 4: OVERLAY SPECTRA OF SAL AND AMB FOR FIRST ORDER DERIVATIVE METHOD

Analysis of tablet formulation: For the estimation of drugs in the commercial formulations, ten tablets containing 2 mg of SAL and 30 mg of AMB were weighed and average weight was calculated. The tablets were crushed and powdered in glass mortar. For the analysis of drugs, quantity of powder

equivalent to 1 mg of SAL and 15 mg of AMB was transferred to 100 ml volumetric flasks and dissolved in sufficient quantity of 0.1N HCL. It was sonicated for 30mins and volume was made up to obtain a stock solution of 10 µg/ml of SAL and 150 µg/ml of AMB. This solution was then filtered through whatman filter paper # 42. Further dilutions were made from this stock solution to get required concentration. In First method, AMB concentration was determined directly from calibration plot by measuring absorbance at 300nm

where SAL shows zero absorbance and SAL was determined by subtracting absorbance of AMB from total absorbance of sample at 223nm (λ_{max} of SAL) by using equation I and II respectively. In second method, concentration of SAL and AMB was determined by measuring the $dA/d\lambda$ at wavelength 252nm (zero crossing of AMB) and 232nm (zero crossing of SAL) using equation III and IV respectively. Results of tablet analysis are shown in **Table 3**. The assay procedure was repeated six times (n=6).

TABLE 3: RESULT OF MARKETED FORMULATION ANALYSIS

Method	Label claim	% Label Claim*(Mean±SD)	%RSD
Absorbance correction method	SAL 2mg	98.97±0.00172	0.2839
	AMB 30mg	98.41±0.000476	0.36033
First order Derivative method	SAL 2mg	100.99±0.1567	0.2328
	AMB 30mg	99.87±0.09987	0.1974

*Average of six determination

Validation: The method was validated according to ICH guidelines to study linearity, accuracy and precision ²⁷.

Linearity: The linearity of measurement was evaluated by analyzing different concentrations of the standard solution of SAL and AMB. For both the methods, the Beer law was obeyed in the concentration range 2-10 µg/ml and 2-20 µg/ml for SAL and AMB respectively.

For both the method, the correlation coefficient was found to be >0.998.

Accuracy (Recovery studies): To ascertain the accuracy of proposed methods, recovery studies were carried out by standard addition method at three different levels (80%, 100% and 120%). Percent recovery for SAL and AMB, by both the methods, was found in the range of 98.20- 102% (**Table 4**).

TABLE 4: RESULTS OF RECOVERY STUDIES

Level Recovery	Drug	Conc. of Drug in µg/ml		Method I*			Method II*		
		Drug taken	Std drug added	%recovery	SD	%RSD	%recovery	SD	%RSD
80	SAL	1	0.8	99.44	0.0138	0.2516	99.88	0.0986	0.1476
100		1	1	98.68	0.0069	0.5742	101.5	0.2986	0.6563
120		1	1.2	99.25	0.0128	0.9447	101.6	0.1965	0.3232
80	AMB	15	12	99.57	0.0030	0.2709	99.79	0.2567	0.4123
100		15	15	98.57	0.0022	0.8510	99.80	0.1543	0.1986
120		15	18	99.86	0.0026	0.8825	100.33	0.1265	0.1756

Precision: The reproducibility of the proposed methods was determined by performing tablet assay at different time intervals on same day (Intra-day precision) and on three different days (Inter-day precision).

RESULTS AND DISCUSSION: Under experimental conditions described, linearity, assay of tablet, accuracy studies and precision were performed. In both the methods linearity was observed in the concentration range of 2-10µg/ml and 2-20 µg/ml for SAL and AMB respectively and correlation coefficient

was found to be > 0.998 . The results of commercial tablet formulation are presented in Table 1. Results of accuracy studies are presented in Table 2. Percent recovery for SAL and AMB by both the methods was found in the range of 98.20% to 102 %. S.D. and R.S.D. for six determinations of tablet sample, by both the methods, was found to be less than 2.0 indicating the precision of both the methods. Hence, it can be concluded that the developed spectrophotometric methods are accurate, precise and can be employed successfully for the estimation of salbutamol and ambroxol in bulk and formulation.

CONCLUSION: The two spectrophotometric methods were developed and validated as per ICH guidelines. The standard deviation and % RSD calculated for the proposed methods are within limits, indicating high degree of precision of the methods. The results of the recovery studies performed indicate the methods to be accurate. Hence, it can be concluded that the developed spectrophotometric methods are accurate, precise and can be employed successfully for the estimation of salbutamol and ambroxol in bulk and formulation.

ACKNOWLEDGEMENT: The authors express their gratitude to Glenmark Pharmaceutical Ltd. Nasik, India for Gift samples of pure drugs and Cheminnova Remedies Pvt. Ltd for tablet formulation. Special thanks to Dr. S.D. Sawant, Principal Smt. Kashibai Navale College of Pharmacy for providing excellent infrastructure facility to carry out this research work.

REFERENCES:

1. Indian Pharmacopoeia, published by the controller of publication: New Delhi: 2007; III: p.1687
2. Wilson and Gisvold's Textbook "Organic Medicinal and Pharmaceutical chemistry" edited by John H. Block and John M. Beale, Wolters Kluwer, London, UK: 2004; p. 96-99.
3. Barnes PJ, Page CP, Pharmacology and therapeutics of asthma and COPD, Springer-Verlag Berlin Heidelberg, published by Jaypee Brothers medical publishers: Germany; 2004, p.15
4. The Merk Index, An Encyclopedia of Chemicals, Drugs and Biologicals, 14th ed., published by Merk laboratories: 2006; p. 216
5. Indian Pharmacopoeia, published by the controller of publication, New Delhi: 2007; III: p. 143,250, 701.
6. The Merk Index, An Encyclopedia of Chemicals, Drugs and Biologicals, 14th ed.: published by Merk laboratories: 2006; p. 385
7. Barnes PJ, Page CP, Pharmacology and therapeutics of asthma and COPD, Springer-Verlag Berlin Heidelberg published by Jaypee Brothers medical publishers: Germany 2004; p.217-218
8. Srinivasan KK, Shirwaikar A, Joseph A, Jacob A, Prabu LS, Simultaneous estimation for the analysis of Salbutamol Sulphate and Ambroxol Hydrochloride in solid dosage forms by ultraviolet Spectrophotometry, Indian Drugs 2005; 42(9): 576-579.
9. Dave HN, Mashru RC, Thakkar AR, Simultaneous determination of salbutamol sulphate, bromhexine hydrochloride and etofylline in pharmaceutical formulations with the use of four rapid derivative spectrophotometric methods, Analytica Chimica Acta 2007; 597:113-120
10. Mukherji G, Aggarwal N, Quantitative estimation of salbutamol sulphate by derivative UV spectroscopy in the presence of albumin, International Journal of Pharmaceutics 1992; 86: 153-158
11. Mukherji G, Aggarwal N, Derivative UV-spectroscopic determination of salbutamol sulphate in the presence of gelatin, International Journal of Pharmaceutics 1991; 71:187-191
12. Parimoo P, Umapathi P, Ilango K, Simultaneous quantitative determination of salbutamol sulfate and bromhexine hydrochloride in drug preparations by difference spectrophotometry, International Journal of Pharmaceutics 1993; 100: 227-231
13. Colthup PV, Dallas FAA, Saynor DA, Carey PF, Skidmore LF and Martin LE, Determination of Salbutamol in human plasma and urine by high-performance thin-layer chromatography, J. Chromatography 1985; 345:111-118
14. Jacobson GA and Peterson GM, High-performance liquid chromatographic assay for the simultaneous determination of ipratropium bromide, fenoterol, salbutamol and terbutaline in nebulizer solution, J. Pharm. & Biomed. Anal. 1994; 12: 825-832
15. Dave H. N., Mashru R. C., Patel A. K., Thin Layer Chromatography Method for the Determination of Ternary Mixture Containing Salbutamol Sulphate, Bromhexine Hydrochloride and Etofylline, J. Pharm. Sci. & Res. 2010; 2(2): 143-148
16. Gowekar NM, Pande VV, Kasture AV, Tekade AR, Chandorkar JG., Spectrophotometric estimation of ambroxol and cetirizine hydrochloride from tablet dosage form. Pak. J. Pharm Sci. 2007; 20(3): 250-1.
17. Bhatia NM, Ganbavale SK, Bhatia MS, More HN, Kokil SU, RP-HPLC and Spectrophotometric estimation of ambroxol hydrochloride and cetirizine hydrochloride in combined dosage form. Indian J. Pharm Sci. 2008; 70: 603-608
18. Makarand A, Bonde CG, Development and validation of simultaneous UV spectrophotometric method for the determination of levofloxacin and ambroxol in tablets, International Journal of ChemTech Research 2009; 1 : 873-888
19. Hadad GM, Gindy AE, Waleed MM, Mahmoud, HPLC and chemometrics-assisted UV-spectroscopy methods for the simultaneous determination of ambroxol and doxycycline in capsule, Spectrochimica Acta Part A 2008; 70 : 655-663
20. Dinçer Z, Basan H, Göger NG, Quantitative determination of ambroxol in tablets by derivative UV spectrophotometric method and HPLC, J Pharm & Biomed. Anal. 2003; 1: 31(5): 867-872.

21. Nobilis M, Pastera J, Svoboda D and Kvstina J, High-performance liquid chromatographic determination of ambroxol in human plasma, *J. Chromatography* 1992; 581: 251-255
22. Meiling Qi, Wang P, Cong R, Yang J, Simultaneous determination of roxithromycin and ambroxol hydrochloride in a new tablet formulation by liquid chromatography, *J.Pharm. & Biomed. Anal.* 2004; 35 :1287–1291
23. Nagappan KV, Meyyanathan SN, Raja RB, Reddy S, Jeyaprakash MR, Birajdar AS et al., A RP-HPLC Method for Simultaneous Estimation of Ambroxol Hydrochloride and Loratidine in Pharmaceutical Formulation, *Research J. Pharm. and Tech.*, 2008; 4: 366-369
24. Maithani M, Raturi R, Vertika Gautam, Dharmendra Kumar, Anand Gaurav and Ranjit Singh, Simultaneous estimation of ambroxol hydrochloride and cetirizine hydrochloride in tablet dosage form by RP-HPLC method, *International Journal of Comprehensive Pharmacy* 2010; 1: 1-3
25. Jain PS., Stability-indicating HPTLC determination of ambroxol hydrochloride in bulk drug and pharmaceutical dosage form, *J chromatographic Sci.* 2010; 48(1): 45-48.
26. Hohyun Kim, Jeong-Yeon Yoo , Sang Beom Han, Hee Joo Lee, Kyung Ryul Lee, Determination of ambroxol in human plasma using LC-MS/MS, *J.Pharm. & Biomed. Anal.*, 2003; 32 : 209-/216
27. ICH Harmonized Tripartite Guidelines, Validation of analytical procedures: text & methodology, Q2 (R), Nov 2005.
