



Received on 19 September, 2015; received in revised form, 13 February, 2016; accepted, 19 December, 2016; published 01 March, 2016

PHYSICO-CHEMICAL AND PHARMACOLOGICAL PROSPECTIVE OF ROGHAN-E-NARJEEL (COCONUT OIL)

Mohd. Shamim Khan ^{*1} Qamrul Hasan Lari ² and Mahmood Ahmad Khan ³

Government Unani Dispensary ¹, Kota, Rajasthan, India

Department of Kulliyat ², State Takmeel-ut-Tib College, Lucknow, UP, India

Unani Expert ³, TKDL, Unit Neeri, CSIR, New Delhi, India

Key words:

Roghan-e-Narjeel,
lauric acid, anti-hyperlipidemic
activity, anti-microbial activity,
healing activity..

Correspondence to Author:

Dr. Mohd. Shamim Khan

Medical Officer,
Govt. Unani Dispensary,
Masjid Gali, BheemganjMandi,
Kota, Rajasthan - 324002, India


Email: drshamimmd@yahoo.co.in

ABSTRACT: Roghan-e-Narjeel (coconut oil) is extracted from dried Maghz-e-Narjeel (coconut kernels) which is derived from coconut (*Cocosnucifera* L.) tree. It is used extensively in tropical countries especially India, Srilanka and Philippines. The oil contains 92% of saturates consisting of medium chain fatty acids in the form of triglycerides containing lauric acid in large amount, and about 8% of unsaturates consisting of oleic and linoleic acids as triglycerides. The oil is colourless and has odour typical of the coconuts. It plays an important role in the treatment of ischemic heart disease, infections (viral, bacterial, fungal, protozoal), wounds and obesity. These effects of oil can be attributed to the presence of lauric acid, capric acid and caprylic acid. The aim of this review paper is to highlight the physico-chemical and pharmacological properties of Roghan-e-Narjeel (coconut oil) according to modern as well as unani aspect.

INTRODUCTION: Roghan-e-Narjeel (coconut oil) also known as coconut butter, is an edible oil extracted from dried Maghz-e-Narjeel (coconut kernels) which contains about 60-65% of the oil and derived from coconut (*Cocosnucifera* L.) tree ¹. ² Coconut oil is one of the primary sources of energy, in tropical countries like India, Srilanka, Philippines, and Indonesia ³. There are commonly four varieties of coconut oil available in the market. Firstly *pure coconut oil* is extracted from dried coconut kernels (copra) by compression in a mill either by bullock or by power. It is crude, unrefined and without any additives. It has multiple uses such as edible oil, massaging oil, hair oil, cosmetic usage, medicinal and industrial purpose.

Secondly is refined coconut oil obtained by mechanically and chemically refining, bleaching and deodorizing the crude coconut oil to make it thin, colorless, odorless and without any type of particle (like protein) suspended in it, for getting only pure saturated fats. Thirdly is virgin coconut oil (VCO) derived from coconut milk extracted from freshly shredded coconut meat and not from copra by processes like fermentation, centrifugal separation and enzymatic action. It is laden with medium chain triglyceride (MCT). It has remarkable anti-oxidants and anti-microbial properties. This is the most respected and trusted variety. Lastly is organic coconut oil that has extracted from coconut palms prepared only on organic manure without involving any chemical, synthetic fertilizers or insecticides in its extraction or processing. This is another well respected variety. This form mainly used in cosmetic area ⁴.

Physico-chemical Properties: It has a long shelf life. The oil has the natural sweet taste of coconut. Coconut oil is insoluble in water. At temperature

<p>QUICK RESPONSE CODE</p> 	<p>DOI: 10.13040/IJPSR.0975-8232.7(3).1286-91</p>
<p>Article can be accessed online on: www.ijpsr.com</p>	
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.7(3).1286-91</p>	

above its melting point it is completely miscible with most of the non-hydroxylic solvents such as light petroleum, benzene, carbon tetrachloride etc. In alcohol, coconut oil is more soluble than most common fats and oils ². Among the most stable of all vegetable oil, coconut oil is slow to oxidize and thus resistant to rancidity. Unrefined coconut oil melts at 20-25°C and smokes at 170⁰ (350⁰F), while refined coconut oil has a higher smoke point of 232⁰C (450⁰F) ⁵. Coconut oil is a triglyceride consisting 92% saturated fatty acids, most of them

(about 70%) are lower chain saturated fatty acids known as medium chain fatty acids (MCFAs), 6% monounsaturated fatty acids and 2% polyunsaturated fatty acids. Out of its saturated fatty acids, 45-56% coconut oil is primarily lauric acid, 16.8% myristic acid and 8.2% palmitic acid. It's only monounsaturated fatty acid is oleic acid while it's only polyunsaturated fatty acid is linoleic acid ^{2, 5}. Physico-chemical Characteristics of unrefined, refined and virgin Coconut Oil are described in **Table 1** ².

TABLE 1: PHYSICO-CHEMICAL CHARACTERISTICS OF COCONUT OIL

Physico-chemical Characteristics	Virgin coconut oil from wet coconut	Unrefined coconut oil from copra	Refined coconut oil
Appearance	Colourless	Slight brownish	Colourless
Odour	Coconut smell	Coconut smell	Odourless
Melting point ⁰ C	24	24	24
Moisture %	<0.1	<0.1	<0.1
Iodine value (cg 12/gm.)	12-15	12-15	10-12
Peroxide value (meq.02/kg.)	0-1	0-1	0-1
Saponification value (mg KOH/g)	245-255	245-255	250-255
Phospholipids (%)	0.1	0.1	0.0
Unsaponifiable matter (%)	--	0.42	0.19
Tocopherols (mg/Kg)	150-200	150-200	4-100
Phytosterols (mg/Kg)	--	400-1200	--
Total phenolics (mg/Kg)	640	618	20
	Fatty Acid Composition Relative (%)		
Saturates	92.0	92.0	92.0
Monounsaturates	6.0	6.0	6.0
Polyunsaturates	2.0	2.0	2.0

The fatty acid composition of the oil is: α -tocopherol, 5; and β -tocopherol, 6ppm: it also contains α -, β and γ -tocotrienol, 5, 1 and 19 ppm, respectively. The fatty acid composition of the oil is: caproic, <1; caprylic, 5-9; capric, 6-10; lauric,

44-52; myristic, 13-19; palmitic, 8-11; stearic, 1-3; oleic, 5-8; and linoleic, <3%. [6] Several derivatives have been obtained from these fatty acids and their uses are given in **Table 2** ⁶.

TABLE 2: APPLICATIONS OF DERIVATIVES OF COCONUT OIL FATTY ACID

Derivative	Application
Ethyl caproate	Synthetic flavours
Amyl caproate	Synthetic flavours
Allylcaproae	Synthetic flavours
Ethyl caprylate	Perfumery intermediate
Amyl caprylate	Perfumery intermediate
Butyl caprylate	Perfumery intermediate
Barium caprylate / caprate	Stabilizer for PVC
Allyl	Sweet banana / pineapple odour
Amyl caprate	Arrachcognacodour
Amylase caprate	Dip type coatin for food products
Butyl caprate	Apricot odour
Caprylcaprate	Chemical pruning activity on plants
Lauryl alcohol	Detergent base
Sodium lauryl sulphate	Cosmetics and toiletries
Sodium lauryl ether sulphate	Shampoo
Iso propyl myristate	Cosmetics

Temperament (Mizaj):Hot 2⁰ & Wet 2⁰⁷**Therapeutic Dosage (Miqdar-e-Khurak):**2-3 Tola⁷**Method of Uses (Tarkeeb-e-Istemat):**

Roghan-e-Narjeel (coconut oil) is used for oral administration (Brah-e-Dahn) as well as topical application in the form of a paste (Zamad) and ointment (Marham)^{6,7,8}.

Pharmacological Actions (Af'aa)Dafe-e-Da-us-sadaf (Anti-psoriatic)⁹Dafe-e-Kharish (Anti-pruritic)⁹Qatil-e-Kirm-e-Shikam (Antihelminthic)⁸Muqavvi-e-Bah (Aphrodisiac)⁷Muqavvi-e-Qalb (Cardiac Tonic)^{5,10}Muqavvi-e-Mana'at (Immunomodulator)^{1,10}Musakkin (Sedative)⁷Murattib (Moisturizer)⁹Mundamil-e-Qurooh (Wounds Healer)⁹Nafey-e-Shaheeqa (Pertussis Reliever)⁸Anti-microbial activity^{1,11}Anti-bacterial activity^{1,11}Anti-viral activity^{1,11}Anti-fungal activity^{1,11}Anti-protozoal activity^{1,11}Anti-septic activity⁹**Therapeutic Uses (Mahall-e-Istemat):**

Roghan-e-Narjeel (Coconut Oil) is specially suggested in the treatment of Da-us-sadaf (Psoriasis), Nar-e-Farsi (Eczema), Hekah (Pruritis), Quba (Ring worm), Namla (Herpes), Busoor-e-Labniyya (Pimples) and Busoor-e-Jild (Skin Rashes)^{7,8,9}. It is also used in Amraz-e-Qalb (Heart Diseases), Amraz-e-Sha'ar (Hair Diseases) like hair's falling, graying and lustrelessness. It is also used in Ziabitus (Diabetes), Deedan-e-Am'aa (Intestinal worms) particularly in Habb-ul-Qara (Tape Worm) and Aujaaye Baridah (cold pains)^{7,8,10}.

Scientific Studies:

Few scientific studies are illustrated below regarding Roghan-e-Narjeel (coconut oil).

Anti-oxidant Activity:

Newinet et al (2005) studied VCO is capable of increasing antioxidant enzymes when supplemented with diets in rats¹².

Kapila et al (2009) compared the antioxidant activities of coconut oil extracted under hot and cold conditions. The coconut oil extracted under hot conditions (HECO) contained more phenolic substances than the coconut oil extracted under cold conditions (CECO). However, high temperatures used in the hot extraction of coconut oil favor the incorporation of more thermally stable phenolic antioxidants into coconut oil. Therefore, the consumption of HECO may result in improvement of antioxidant related health benefits compared with the consumption of CECO¹³.

Anti-hyperlipidemic Activity:

Newin et al (2007) showed the VCO lowered total cholesterol, triglycerides, phospholipids, low density lipoprotein (LDL), very low density lipoprotein (VLDL) and increased high density lipoprotein (HDL)¹⁴.

Vess by (1994) reported the monounsaturated fatty acids such as oleic acid are as effective in reducing serum total and low density lipoprotein cholesterol levels as polyunsaturated fatty acids such as linoleic acid⁵.

In 1989 Mendis et al reported undesirable lipid changes when young adult Sri Lankan males were changed from their normal diets by the substitution of corn oil for their customary coconut oil. Although the total serum cholesterol decreased 18.7 per cent from 179.6 to 146.0 mg/dl and the LDL cholesterol decreased 23.8 per cent from 131.6 to 100.3 mg/dl, the HDL cholesterol decreased 41.4 per cent from 43.4 to 25.4 mg/dl (putting the HDL values below the acceptable lower limit) and the LDL/ HDL ratio increased 30 per cent from 3.0 to 3.9¹⁵.

Ahrens and colleagues (1957) had shown that adding coconut oil to the diet of

hypercholesterolemics lowers serum cholesterol from 450 mg/ dl to 367 mg/dl. This is hardly a cholesterol raising effect¹⁵.

Hostmark *et al* (1980) compared the effects of diets containing 10% coconut fat and 10 % sunflower oil on lipoprotein distribution in male wistar rats. Coconut oil feeding produced significantly lower levels ($p < 0.05$) of pre-beta lipoproteins (VLDL) and significantly higher ($p = < 0.01$) alpha-lipoproteins (HDL) relative to sunflower oil¹⁵.

Anti-viral Activity:

Hierholzer and Kabara (1982) showed virucidal effects of monolaurin (a disease-fighting fatty acid derivative produced when the lauric acid in coconut oil is used by the body) on enveloped RNA and DNA viruses was done in conjunction with the Center for Disease Control of the US Public Health Service with selected prototypes or recognized representative strains of enveloped human viruses. The envelope of these viruses is a lipid membrane¹⁵.

Kabara (1978) and others have reported that certain medium-chain fatty acids such as lauric acid and their derivatives like mono-glycerides (monolaurin) can have adverse effects on various micro-organisms including bacteria, yeast, fungi, and enveloped viruses that are inactivated by disrupting the lipid membranes of the organisms (Isaacs and Thormar 1991; Isaacs *et al* 1992).

Some of the viruses inactivated by these lipids, in addition to HIV, are the measles 1), vesicular stomatitis virus (VSV), isna virus, and cytomegalovirus(CMV)¹⁵.

Anti-microbial Activity:

It is reported by Kabara (1972) that *in vitro* Lauric Acid and Capric Acid were active against all gram-positive and gram-negative organisms and Candida. Minimum Inhibitory Concentration ranges for lauric acid were 0.062–2.249 micromoles/ml, and were 1.45 and 5.8 micromoles/ml for capric acid¹⁶. Oyiet al (2010) reported that VCO in water emulsions killed *S. aureus* was by 6 hrs, *Ps. Eruginosa* by 48 hrs. *Candida* or *A. niger* by 7 days *in vitro*¹⁶.

Healing Activity:

Nevin *et al* (2010) reported that VCO-treated wounds healed much faster due to higher collagen and anti-oxidant enzymes activities. It was also observed histopathologically, that there is increase in fibroblast proliferation and neovascularization in VCO-treated wounds¹⁷.

Moisturizing activity:

Agero *et al* (2004) reported that Coconut oil is as effective and safe as mineral oil when used as a moisturizer in xerosis (dry & rough skin). A randomized double-blind controlled clinical trial was conducted on mild to moderate xerosis in 34 patients with negative patch-test reactions to the test products. These patients were randomized to apply either coconut oil or mineral oil on the legs twice a day for 2 weeks. Quantitative outcome parameters for effectivity were measured at baseline and on each visit with a Corneometer CM825 to measure skin hydration and a Sebumeter SM 810 to measure skin lipids. Both oils showed effectivity through significant improvement in skin hydration and increase in skin surface lipid levels. Subjective grading of xerosis by the investigators and visual analogue scales used by the patients showed a general trend toward better (though not statistically evident) improvement with coconut oil than with mineral oil, therefore Coconut oil is superior to mineral oil in dry skin¹⁸.

Immunomodulator Activity:

Winarsi *et al* (2008) reported that the VCO enriched with zinc increased Tc cells, Th cells, IL-2, but maintained the number of neutrophils and NK cells, while the IgG level changed from equivocal to negative in candidiasis patient¹⁹.

Anti-diabetic Activity:

It is reported by Girotti *et al* (1984) that virgin coconut oil (VCO) has been shown to possess insulin tropic effects shown in isolated perfused mouse islet with hypolipidemic effects²⁰. Further studies by Siddalingaswamy *et al* (2011) on diabetic rat with force fed with 2ml alcoholic extracts of commercial coconut oil (CCO), CEVCO and HEVCO for 21 days. The results indicated HEVCO reduced blood glucose and lipids viz. total cholesterol (TC), triglycerides (TG), High density lipoproteins (HDL), Low and Very Low Density

Lipoprotein (LDL+ VLDL) and thiobarbutyric acid reactive substances (TBARS) increased the antioxidant status by elevating activities of antioxidant enzymes such as superoxide dismutase (SOD), catalase, glutathione peroxidase (GSH-Px), glutathione (GSH) concentration and decreased lipid peroxidation in liver than CEVCO. These beneficial effects may be due to increased polyphenolic and other antioxidants content present in HEVCO²¹.

Kabra *et al* (2012) reported that *Cocosnucifera* flower extract (CNFE) and coconut oil (CO) at a dose of 300 mg/kg body weight for 15 days, suppressed the elevated blood glucose and lipid levels in alloxan induce diabetic rats. These findings indicated that the *Cocosnucifera* flower extract and coconut oil possess antihyperlipidemic effect in addition to anti-diabetic activity. The hypoglycemic activity of coconut oil seen in this study could be due to the fatty acids present in the oil²². It is reported by Maedler *et al* (2003) that monosaturated fatty acids improved β -cell secretory function by preventing β -cell apoptosis, decrease β -cell proliferation and impairment of β -cell function²³.

Anti-obesity Property:

Liau *et al* (2011) reported that VCO is safe and effective in reducing visceral adiposity in obese, though healthy men. An open label pilot study was conducted over 20 obese but healthy volunteers with intake of VCO for four weeks. Result has been concluded that only waist circumference was significantly reduced with a mean reduction of 2.86cm or 0.97% from initial measurement ($p=0.02$). Waste reduction was seen in only males ($p<0.05$)²⁴.

Another study by Assuncao *et al* (2009) showed that dietary coconut oil elevates High Density Lipoprotein (HDL) levels and reduces abdominal obesity in women. This randomized double-blind clinical trial involved 40 obese women, aged 20-40 years, presenting waist circumference(WC)>88cm (abdominal obesity). Groups received daily dietary supplements comprising 30 ml of either soybean oil (Group S; n=20) or coconut oil (Group C; n=20) over a 12- week periods. After 12- week, group C presented a higher level of HDL (48.7+/- 2.4 vs.

45+/-5.6; $P=0.01$) and a lower LDL: HDL ratio 2.41+/- 0.8 vs. 3.1 +/- 0.8; $P=0.04$). Reductions in BMI were observed in both groups but only group C exhibited a reduction in WC ($P=0.005$)²⁵.

Anti-cancer Activity:

Lim-Sylianco (1987) published a 50-year literature review showing the anti-cancer effects of coconut oil²⁶. Cohen *et al* (1986) showed that coconut oil was by far more protective than unsaturated oils in chemically induced cancers of the colon and breast, for example 32% of corn oil eaters got colon cancer whereas only 3% of coconut oil eaters got the cancer. Animals fed unsaturated oils had more tumors. This shows the thyroid-suppressive and hence, immuno-suppressive effect of unsaturated oils²⁷.

Antithrombotic Property:

VCO has significant antithrombotic effect over copra oil¹⁴. A coconut oil based diet high saturated fatty acid diet lowers postparential t-PA(tissue plasminogen activator) antigen concentration, and this may favorably affect the fibrinolytic system and the Lp-a (lipoprotein-a) concentration compared with the high mono polyunsaturated fatty acid diet. The proportions of dietary saturated fatty acids more than the percentage of saturated fat energy seem to have a beneficial influence Lp-a (lipoprotein-a) levels²⁸.

CONCLUSION: Coconut oil has many health benefits, which includes hair care, skin care, stress relief, cholesterol level maintenance, weight loss and boosted immune system. These benefits of oil can be attributed to the presence of lauric acid, capric acid and caprylic acid. The oil is known to have antioxidant, antiviral, antibacterial, antifungal antiprotozoal, antihyperlipidemic effects and excellent healing & moisturizing properties. However, more research is needed to clearly understand the many good effects of the oil.

ACKNOWLEDGEMENT: I am thankful to Director, Deputy Director, Assistant Directors, Drug Inspector, Department of Unani Medicine Rajasthan – Jaipur, India, for their encouragement and support.

REFERENCES:

- Mandal MD, Mandal S, Coconut (*Cocosnucifera* L.: *Arecaceae*): In health promotion and disease prevention. *Asian Pacific Journal of Tropical Medicine*, 2011; 241-247.
- Krishna S, Raj G, Bhatnagar AS, Kumar P, et al, Coconut Oil: Chemistry, Production and Its Applications -A Review. *Indian Coconut Journal*, 2010; 15-27.
- Fife B, Coconut cures: Preventing and treating common health problems with coconut. Piccadilly Books, Ltd., London, 2005; 184-185.
- <https://www.organicfacts.net/.../oils/health-benefits-of-coconut-oil.html>. Health Benefits of Coconut Oil: Organic Facts.
- http://en.wikipedia.org/wiki/Coconut_oil#Physical_properties. coconut oil; physical properties and health effects
- Anonymous, *The Wealth of India*, PID, CSIR New Delhi, 2001; 2:296-298.
- Kabeeruddin HM, *Makhzan-ul-Mufradat: Kitab-ul-Advia*, Faisal Publication Deoband, UP. 2000; 575.
- AliHSS, *UnaniAdvia-e-Mufrada*, 8th Edition, Qaumi Council Barai Farogh Zaban-e-Urdu, New Delhi, 1999; 270-271.
- http://www.coconut-connection.com/skin_care.htm. Virgin Coconut Oil and Your Skin.
- http://experts.about.com/e/c/co/Coconut_oil.htmhd1. coconut oil; health effects.
- A.R. Oyi, J.A. Onalapo, R.C. Obi, Formulation and Antimicrobial Studies of Coconut (*Cocosnucifera* Linn.) Oil. *Research Journal of Applied Sciences, Engineering and Technology*. 2010, 2(2): 133-137.
- Nevin KG, Rajamohan T, Virgin coconut oil increases the antioxidant status in rats. *Food Chem*. 2005; 99: 260-266.
- Seneviratne KN, Dissanayake MS, Variation of phenolic content in coconut oil extracted by two conventional methods. *International J. Food Sci. and Technol.*, 2008; 43: 597-602.
- Nevin KG, Rajamohan T, Influence of virgin coconut oil on blood coagulation factors and lipid levels and LDL oxidation in cholesterol fed Sprague-Dawley rats. *Eur e-J Clin Nutr Metabol.*, 2007; e1-e8.
- Enig MG, Health and nutritional benefits from coconut oil and its advantages over competing oils. *Indian Coconut Journal*, September 2010; 9-15.
- Elmore LK, Nance G, Singleton S, et al. Treatment of Dermal Infections with Topical Coconut Oil. *Natural Medicinal Journal*. 2014; 6(5):1-13.
- Nevin KG, Rajmohan T, Effect of topical application of virgin coconut oil on skin components and antioxidant status during dermal wound healing in young rats. *Skin Pharmacol Physio*. 2010; 23(6):290-297.
- Agero ALC, Rowell VMV, A randomized double-blind controlled trial comparing extra virgin coconut oil with mineral oil as a moisturizer for mild to moderate xerosis. *Dermatitis*. 2004; 15(3):109-16. PMID: 15724344
- Winarsi H, Purvanto A, Virgin Coconut Oil (VCO) enriched with Zn as immunostimulator for vaginal candidiasis patient. *Hayati J. Biosc*. 2008; 15(4):135-139.
- Girotti AW, Thomas IP, Damage effects of oxygen radicals on resealed erythrocyte ghosts. *The Journal of Biological Chemistry*, 1984; 259: 1744-1752.
- Siddalingaswamy M, Rayaorth A, Khanum F, Anti-diabetic effects of cold and hot extracted virgin coconut oil. *Journal of Diabetes Mellitus*, 2011; 1(4): 118-123.
- Kabra MP, Rachhadiya RM, Desai NV, et al, Hypoglycemic Effect of *Cocosnucifera* Flower Alcoholic Extract and Oil in Normal and Alloxanised Hyperglycemic Rats. *International Journal of Pharmaceutical and Phytopharmacological Research*, 2012; 2(1): 29-31.
- Maedler K, Oberholzer J, Bucher P, et al, Monounsaturated fatty acids prevent the deleterious effects of palmitate and high glucose on human pancreatic β -cell turnover and function. *Diabetes* 2003; 52:726-733.
- Liau KM, Lee YY, Chen C.K, et al, An open label pilot study to assess the efficacy and safety of virgin coconut oil in reducing visceral adiposity. *ISRN Pharmacol*. 2011; 2011: 949686. Epub 2011 March 15. PMID: 22164340.
- Assuncao ML, Ferreira HS, Santos AFD, et al, Effects of dietary coconut oil on the biochemical and anthropometric profiles of women presenting abdominal obesity. *Lipids*, 2009; 44(7): 593-601. Epub 2009 May 13. PMID: 19437058.
- Lim-Sylianco CY, Anticarcinogenic effects of coconut oil. *Phillip. J. Coconut Studies*, 1987; 12 (2): 89-102.
- Cohen LA, Thompson DO, Maeura, et al, Dietary fat and mammary cancer. I. Promoting effects of different dietary fats on N-nitrosomethylurea induced rat mammary tumorigenesis. *J. Natl. Cancer Inst.*, 1986; 77 (1): 33-42.
- Muller H, Lindman AS, Blomfeldt A, et al, A diet rich in coconut oil reduces diurnal postparental variations in circulating tissue plasminogen activator antigen and fasting lipoprotein-a compared with a diet rich in unsaturated fat in women. *J Nutr*. 2003; 133(11): 3422-3427.

How to cite this article:

Mohd. Khan S Lari QS and Mahmood Khan A: Physico-Chemical and Pharmacological Prospective of Roghane-Narjeel (Coconut Oil). *Int J Pharm Sci Res* 2016; 7(3): 1286-91. doi: 10.13040/IJPSR.0975-8232.7(3).1286-91.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)