



Received on 17 June, 2017; received in revised form, 13 January, 2018; accepted, 27 January, 2018; published 01 March, 2018

INVESTIGATION OF THE EFFECTS OF ETHANOL SEED EXTRACT OF *GARCINIA KOLA* ON HEMATOLOGICAL AND HEPATORENAL INDICES IN WISTAR ALBINO RATS

J. S. Aprioku

Department of Experimental Pharmacology and Toxicology, Faculty of Pharmaceutical Sciences, University of Port Harcourt, East-West Road, Choba, Rivers State, PMB 5323, Nigeria.

Keywords:

Bitter kola, *Garcinia kola*,
Hematological indices, Lymphocyte,
Transaminase enzymes

Correspondence to Author:

Dr. J. S. Aprioku

Associate Professor
Head Department of Experimental
Pharmacology and Toxicology,
Faculty of Pharmaceutical Sciences,
University of Port Harcourt, East-
West Road, Choba, Rivers State,
PMB 5323, Nigeria.

E-mail: sydaprio@yahoo.com

ABSTRACT: *Garcinia kola* is a known herbal plant which is widely consumed for many ethnomedicinal uses. This study was aimed at determining the effect of ethanol extract of *G. kola* seeds on levels of hematological as well as blood indices of liver (aspartate transaminase, alanine transaminase, alkaline phosphatase, total protein, and albumin) and kidney function (creatinine and urea). Extract (50, 100, 200 or 300 mg/kg) was orally administered to different groups of rats (n = 6) daily for thirty days. Control animals were given vehicle (DMSO + distilled water, 2:1). Blood levels of the parameters mentioned above were measured at the end of extract treatment. Livers and kidneys of animals were removed and weighed and organ-to-body weight ratio was obtained. Extract had no effect on serum levels of liver and kidney indices, except that it reduced (P<0.05) ALP level at 300 mg/kg. Organ-to-body weight ratio of liver and kidney were not affected by extract. Furthermore, the extract increased (P<0.001) platelet count, but had no effect on RBC and its indices (mean cell volume, mean cell haemoglobin, and mean cell haemoglobin concentration), Hb and PCV compared to control. It also caused reduction (P<0.001) in WBC and neutrophil counts, while lymphocyte was increased. The result suggests that *G. kola* seed may have no adverse influence on hematologic, hepatic or renal system, and may additionally have hepato-protective property, and enhance blood clotting and lymphocyte function.

INTRODUCTION: *Garcinia kola* is a species of flowering plant that belongs to the Family, Clusiaceae or Gutiferae. Its natural habitat is subtropical or tropical moist lowland forests. The plant grows to a height of about 15 m and produces reddish, yellowish or orange color fruit containing 2 to 4 seeds.

Garcinia kola is consumed amongst Africans, either leisurely or for medicinal purposes. The fruit, Seed, nut and bark of *Garcinia kola* have been used for centuries in alternative medicine to treat a variety of ailments like cough, fever, bronchitis, asthma, throat infections, diarrhoea, and as antidote for poisons¹⁻⁵.

Other conditions include rheumatism, headache, constipation, abdominal colic, oral hygiene, and liver disorders^{2, 4, 6}. *Garcinia kola* contains phytochemicals which have potent biological activities⁷⁻⁹, and these have been attributed to most of the plant's medicinal properties. Presently, the plant is believed to be highly efficacious and therefore used

<p>QUICK RESPONSE CODE</p> 	<p>DOI: 10.13040/IJPSR.0975-8232.9(3).975-80</p> <hr/> <p>Article can be accessed online on: www.ijpsr.com</p> <hr/> <p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.9(3).975-80</p>
---	---

as an alternative to orthodox medicines for treatment of different illnesses by many in Nigeria and other African countries. It is growing in popularity as a natural and effective remedy for weight reduction, and particularly popular for controlling cough, diabetes and diarrheal episodes. However, there is no sufficient knowledge about the safety of prolonged consumption of the plant. A study to assess the toxicity potential of prolonged use of the plant on liver and kidney functions may therefore be useful. The liver is a vital organ which functions include, detoxification of chemical compounds, synthesis of essential molecules and metabolism^{10, 11}. The kidney disposes off waste products from the body.

The objective of this study was to assess the effects of ethanol extract of *Garcinia kola* seeds on biochemical indices of liver and kidney functions in Wistar rats. Its effect on hematological indices was equally evaluated.

MATERIALS AND METHODS:

Materials:

Chemicals / Reagents: Dimethyl sulphoxide, DMSO (Zhuzhou Hansen Chemical Industry Co. Ltd., Shanghai, China), ethanol (JHD Chemicals, Guangdong, China), biochemical test kits (Randox Inc., USA)

Equipment: Analytical balance (Model CPA 3245, Sartonus AG Gohingen, Germany), spectrophotometer (Spectrum Lab., USA), oven (Model DHG - 9101 - ISA, Health Medical Equipment, England), water bath (Finlab Company, Lagos, Nigeria).

Methods:

Preparation and Extraction of Plant Material:

Fresh *Garcinia kola* seeds were purchased from a Fruit Market (Oil Mill Market), Aba Road, Port Harcourt, Nigeria in 2016. It was identified by a botanist, Mr. M. Suleiman of the Department of Pharmacognosy and Phytotherapy, Faculty of Pharmaceutical Sciences, University of Port Harcourt, Nigeria and a specimen number (DPH048) was assigned to it and deposited in the Faculty's herbarium. The seeds were air-dried for about one week and oven-dried at 40 °C until a constant weight was obtained. The seeds were then ground using an electric blender and 1000 g weight

powder was macerated in 5000 ml of 70% ethanol in a wide mouthed extraction bottle with continuous agitation for seven days. Extraction mixture was filtered with Whatman number 51 filter paper. The filtrate was evaporated using a rotary evaporator to obtain a pasty extract which was transferred into a crucible and remaining liquid was evaporated using hot water bath. The dry extract was then transferred into a clean container and stored in a refrigerator, and percentage yield was determined (19.4 %).

Phytochemical Analysis of Ethanol Extract of *Garcinia kola* Seeds: Phytochemical analysis of the extract was done using conventional methods¹².

Animals: Thirty Wistar albino male rats (180 - 200 g body weight) were obtained from the Animal House in Department of Experimental Pharmacology and Toxicology, University of Port Harcourt, Nigeria and used for the study. They were maintained at room temperature (24 ± 3 °C), under natural lighting condition, and fed with rodent's pellets and tap water (given *ad libitum*). Animal experiments were done according to approved guideline of my Institutional Research Ethics Committee (UPH/CHREC/APP/049/2016) and all animals were cared and handled according to stipulated standard guidelines.

Experimental Design: Animals were divided into five groups, each containing six rats, and given 0, 50, 100, 200 and 300 mg/kg of extract daily by oral gavage for 30 days. Vehicle used to dissolve extract was DMSO and distilled water (2:1). Control group was given 0.5 ml vehicle. Doses of extract administered were below the LD₅₀ of 5000 mg/kg as reported previously¹¹. The rats were anesthetized with diethyl ether 24 h after the last extract treatment and sacrificed by cervical dislocation. Blood was collected *via* cardiac puncture with sterile syringes and needles into EDTA and plain bottles.

Levels of hematological parameters- red blood cell (RBC), white blood cell (WBC), platelet, hemoglobin (Hb), packed cell volume (PCV), mean cell volume (MCV), mean cell hemoglobin (MCH), and mean cell hemoglobin concentration (MCHC) in whole blood in EDTA bottles were measured using an Hematology Auto-Analyzer (Sysmex 1-5-

1 Wakino-hama-Kaigandori Chuo-ku, Kobe 651-0073, Japan). Serum was separated from blood in plain bottles and the concentrations of aspartate transaminase, alanine transaminase, alkaline phosphatase, albumin, total protein, urea and creatinine were assayed using a Clinical Chemistry Auto-Analyzer (Sylectra Pro S, Model: 13-9693, Netherland). Liver and kidney were also removed and weighed and organ-to-body weight ratio for each organ was determined.

Statistical Analysis: Data obtained are presented as Mean \pm Standard Error of Mean. Data were analyzed by one-way analysis of variance (ANOVA) followed by Dunnett's Multiple Test for pair wise comparisons using GraphPad Prism 5 software. Probability values < 0.05 were considered significant.

RESULTS:

Phytochemical Analysis: Phytochemical analysis of extract showed the presence of carbohydrates, alkaloids, flavonoids, tannins, terpenoids and saponins, while arthraquinones and cardiac glycosides were absent. Quantitatively, flavonoids were most while steroids were least present in extract **Table 1**.

TABLE 1: PHYTOCHEMICAL ANALYSIS OF ETHANOL EXTRACT OF GARCINIA KOLA SEEDS

Phytochemical constituent	Observation
Alkaloids	++
Terpenoids	++
Saponins	++
Tannins	++
Flavonoids	+++
Steroids	+
Anthraquinones	-
Cardiac glycosides	-
Carbohydrates	++

- Absent

+ Slightly present

++ Moderately present

+++ Abundantly present

Hepatorenal Blood Indices: Serum levels of total protein, alanine transaminase (ALT), aspartate transaminase (AST) and albumin were not affected in all treated rats compared to control, but alkaline phosphatase (ALP) was reduced ($P = 0.02$) in rats that received 300 mg/kg (the highest dose) of the extract **Table 2**.

In addition, serum urea and creatinine levels in extract treated rats were lower but not significant ($P > 0.05$) when compared to control group **Table 3**.

TABLE 2: SERUM LEVELS OF HEPATIC INDICES IN WISTAR RATS AFTER SUBACUTE ADMINISTRATION OF ETHANOL GARCINIA KOLA SEED EXTRACT

Dose (mg/kg)	ALT (U/L)	AST (U/L)	ALP (U/L)	Total protein (g/L)	Albumin (g/dl)
Control	110.01 \pm 5.19	21.28 \pm 2.90	74.79 \pm 1.67	10.34 \pm 6.13	20.99 \pm 5.44
50	114.08 \pm 6.66	26.88 \pm 1.67	77.05 \pm 3.67	8.22 \pm 2.74	16.48 \pm 5.29
100	95.90 \pm 7.18	24.22 \pm 3.35	68.28 \pm 0.80	6.29 \pm 2.17	13.82 \pm 3.89
200	121.68 \pm 10.56	18.97 \pm 4.53	70.93 \pm 3.57	6.97 \pm 3.76	26.93 \pm 2.97
300	130.32 \pm 8.93	15.48 \pm 2.45	61.80 \pm 1.31*	7.37 \pm 2.49	17.80 \pm 4.25

Values are expressed as Mean \pm Standard Error of Mean (SEM), n = 6 rats per group.

ALT: Alanine transaminase, AST: Aspartate transaminase, ALP: Alkaline phosphatase, * Significant at $P < 0.05$

TABLE 3: SUBACUTE ORAL TREATMENT OF ETHANOL GARCINIA KOLA SEED EXTRACT DOES NOT ALTER SERUM LEVELS OF UREA AND CREATININE IN WISTAR RATS

Dose (mg/kg)	Urea (mmol/l)	Creatinine (μ mol/l)
Control	6.67 \pm 1.63	100.77 \pm 11.03
50	4.04 \pm 0.43	77.41 \pm 10.48
100	5.77 \pm 1.59	70.35 \pm 8.98
200	3.85 \pm 0.44	76.15 \pm 10.75
300	3.98 \pm 0.41	79.56 \pm 10.86

Values are expressed as Mean \pm Standard Error of Mean (SEM), n = 6 rats per group.

Haematological Indices: There was non significant reduction ($P = 0.5196$) of RBC count in extract treated rats compared with control **Table 4**. There was no significant change in Hb concentration ($P = 0.4111$) or PCV ($P = 0.5274$) in all rats that were administered extract compared to

control **Table 4**. Mean cell haemoglobin concentration (MCHC), mean cell volume (MCV), and mean cell haemoglobin (MCH) in treated rats were equally not altered when compared to control **Table 4**.

In addition, WBC count was reduced ($P = 0.0002$) dose-dependently in extract administered rats; lymphocytes were increased ($P < 0.05$), while neutrophils were reduced when compared with control **Table 5**. Platelet count was elevated in

extract treated rats in a dose-dependent fashion, but only the count in rats that received the highest dose (300 mg/kg) showed significance ($P < 0.0004$) when compared with control rats **Table 5**.

TABLE 4: EFFECT OF SUBACUTE ORAL ADMINISTRATION OF ETHANOL *GARCINIA KOLA* SEED EXTRACT ON RED BLOOD CELL (RBC) COUNT AND IT'S MARKERS IN WISTAR RATS

Dose (mg/kg)	RBC ($\times 10^{12}/L$)	Hb (g/dL)	PCV (%)	MCHC (%)	MCV (%)	MCH (%)
Control	8.16 \pm 0.25	16.60 \pm 0.49	49.60 \pm 1.63	33.62 \pm 0.29	60.52 \pm 0.15	20.35 \pm 0.18
50	7.18 \pm 0.62	14.60 \pm 0.80	45.00 \pm 2.70	32.53 \pm 0.78	63.38 \pm 2.19	20.64 \pm 0.98
100	6.80 \pm 0.59	14.52 \pm 0.90	43.60 \pm 2.69	33.44 \pm 0.13	64.98 \pm 2.73	21.63 \pm 0.90
200	7.04 \pm 0.99	14.34 \pm 1.70	43.00 \pm 5.10	33.34 \pm 0.03	62.70 \pm 3.11	20.91 \pm 1.04
300	7.83 \pm 0.12	16.03 \pm 0.24	48.00 \pm 0.71	33.38 \pm 0.02	61.36 \pm 0.71	20.36 \pm 0.21

Values are expressed as Mean \pm Standard Error of Mean (SEM), n = 6 rats per group.

TABLE 5: EFFECT OF SUBACUTE ORAL ADMINISTRATION OF ETHANOL *GARCINIA KOLA* SEED EXTRACT ON PLATELET AND WHITE BLOOD CELL (WBC) COUNTS IN WISTAR RATS

Dose (mg/kg)	Platelet ($\times 10^9/L$)	WBC ($\times 10^9/L$)	Neutrophil (%)	Lymphocyte (%)
Control	232.00 \pm 12.41	12.40 \pm 1.29	34.40 \pm 3.14	65.60 \pm 3.14
50	240.00 \pm 12.25	8.08 \pm 1.19*	27.60 \pm 2.50	59.00 \pm 2.49
100	246.00 \pm 13.64	6.60 \pm 0.67**	41.00 \pm 2.49	70.60 \pm 2.80
200	262.00 \pm 13.19	6.80 \pm 0.98**	24.40 \pm 2.80*	75.60 \pm 2.80*
300	334.00 \pm 12.08*	4.20 \pm 0.51***	22.50 \pm 1.04**	77.50 \pm 1.04**

Values are expressed as Mean \pm Standard Error of Mean (SEM), n = 6 rats per group.

* Significant value at $P < 0.05$; **Significant at $P < 0.01$; ***Significant value at $P < 0.001$

Liver and Kidney Weights: Relative weight of liver and kidney (organ-to-body weight ratio) of extract administered rats were not changed when compared with control weights **Table 6**.

TABLE 6: SUBACUTE ORAL TREATMENT OF ETHANOL *GARCINIA KOLA* SEED EXTRACT DOES NOT AFFECT LIVER AND KIDNEY WEIGHTS OF WISTAR RATS

Dose (mg/kg)	Liver		Kidney	
	Organ weight (g)	Organ-to-body weight ratio ($\times 10^2$)	Organ weight (g)	Organ-to-body weight ratio ($\times 10^2$)
Control	8.34 \pm 1.19	2.81 \pm 10.17	0.92 \pm 0.40	0.13 \pm 2.22
50	8.05 \pm 0.26	2.18 \pm 2.01	0.65 \pm 0.10	0.22 \pm 0.15
100	8.30 \pm 1.03	3.22 \pm 15.21	0.68 \pm 0.11	0.23 \pm 1.24
200	6.51 \pm 1.02	3.20 \pm 29.14	0.64 \pm 0.12	0.33 \pm 2.92
300	4.29 \pm 0.13	2.41 \pm 2.18	0.70 \pm 0.11	0.34 \pm 0.19

Values are expressed as Mean \pm Standard Error of Mean (SEM), n = 6 rats per group.

DISCUSSION: Serum levels of alanine transaminase (ALT) and aspartate transaminase (AST) are commonly measured clinically as biomarkers of liver function¹³⁻¹⁵. ALT is present mostly in liver and to a lesser extent plasma and body tissues, whereas AST is found in a number of organs including liver, heart, skeletal muscle, kidneys, brain and red blood cells¹⁶. Elevation in serum ALT and AST levels is suggestive of hepatocellular injury¹⁶.

From the results, extract treatment did not affect AST and ALT levels. Furthermore, the extract at the highest concentration reduced ALP level (which is also an indicator of liver function) but

lower doses had no effect on the enzyme level. Similar results of reduction in ALP have been reported in rabbits but effect was produced by higher dose (500 mg/kg) of ethanol extract¹⁷. Albumin is a plasma protein primarily produced by liver cells and reduction in blood level of albumin is indicative of liver dysfunction¹⁶. It was observed that both albumin and total proteins in serum were not altered by extract treatment.

The above results of the present study thus suggest that ethanol extract of *Garcinia kola* seed is not harmful to the liver over the dose range and duration used and doses higher than 300 mg/kg may have hepatoprotective activity. Additionally,

extract did not affect liver-to-body weight ratio, and this adds more evidence that *Garcinia kola* does not affect liver function^{18 - 20}. Furthermore, serum levels of urea and creatinine (standard indices of renal function) were not changed after extract treatment. Kidney-to-body weight ratio was equally not altered by extract suggesting that the plant would not be harmful to the kidney^{18 - 20}.

Hematopoietic system consists of blood, the constantly circulating fluid that serves the body with supply of nutrients and oxygen as well as removal of wastes. It also provides immunological functions, including circulation of white blood cells and detection and elimination of foreign materials. Changes in blood indices have been linked to exposure to harmful chemicals or alteration in certain physiological factors like cellular integrity and membrane permeability of cells²¹.

In this study, *Garcinia kola* treatment had no effect on haemoglobin (Hb), packed cell volume (PCV), red blood cell (RBC) count and its indices, including, mean cell volume (MCV), mean cell haemoglobin (MCH) and mean cell haemoglobin concentration (MCHC). This is in disagreement with the studies of Ahumibe and Braide²² and Uko *et al.*,²³. Ahumibe and Braide²² reported elevation of RBC, PCV and Hb after treatment with *Garcinia kola*, but at much higher doses (300 - 1200 mg/kg). Whereas, *Garcinia kola* was reported to cause reduction in Hb, PCV and erythrocyte count in young growing rats²³.

However, our finding was similar to the results of Dada and Ikuero²⁴ who demonstrated that *Garcinia kola* failed to cause change in erythrocyte count and PCV over a dose range of 250 - 2000 mg/kg. In addition, the extract in our study increased platelet count, which was consistent with the findings of Atsukwei *et al.*,²⁵ but in disagreement with Ahumibe *et al.*,²² where platelet count was reported to be unaffected by extract treatment.

Furthermore, the extract did not cause any change in total white blood cell count after treatment, but increased lymphocyte level, while neutrophil was decreased at the high doses levels of 200 - 300 mg/kg. Increase in number of platelets by the extract from our result indicates that the plant may enhance blood clotting activity in animals.

On the other hand, lymphocytes play essential roles in immunological response and their elevation in this study suggests that the plant may enhance immune function. This justifies its use as an immune booster.

The phytochemical analysis of the extract revealed the presence of carbohydrates, alkaloids, flavonoids, tannins, saponins, steroids, terpenoids, while cardiac glycosides and arthraquinones were absent. These phytochemicals may play vital role in the results that were obtained in this study.

CONCLUSION: Subacute treatment of ethanol *Garcinia kola* seed extract causes no effect on circulating markers of kidney and liver, and may have hepato-protective effect. In addition, the plant produces no effect on red blood cell count but increases lymphocyte and platelet levels, and may therefore possess blood clotting and immune boosting properties.

ACKNOWLEDGEMENT: Author thanks laboratory staff of Department of Experimental Pharmacology and Toxicology, University of Port Harcourt for providing technical assistance.

CONFLICT OF INTEREST: Nil

REFERENCES:

1. Ibiblio JO: Some Medicinal Plants of Nigeria. Ibadan Publishers Nigeria Ltd., Nigeria 1983; 132-133.
2. Ekene EN and Erhirhie EO: *Garcinia kola*: A review of its ethnomedicinal, chemical and pharmacological properties. International Journal of Current Research and Review 2014; 6(11): 01-07.
3. Falang KD, Uguru MO and Nnamonu NL: Anti-pyretic activity of *Garcinia kola* seed extract. European Journal of Medicinal Plants 2014; 4(5): 511-521.
4. Tcheghebe OT, Signe M, Seukey AJ and Tatong FN: Review on traditional uses, phytochemical and pharmacological profiles of *Garcinia kola* Heckel. Merit Research Journal of Medicine and Medical Sciences 2016; 4(11): 480-489.
5. Mensah JK, Anamoah C, Johnson KB and Akuoko Y: Assessment of the antimicrobial and antioxidant activities of the extracts of *Garcinia kola*, *Garcinia epunctata* and *Acacia kamerunensis* plants used for oral hygiene in Ghana. Current Science Perspectives 2017; 3(1): 13-20.
6. Agbor MA and Naidoo S: Ethnomedicinal plants used by traditional healers to treat oral health problems in Cameroon. Evidence - Based Complementary and Alternative Medicine 2015.
7. Sanda KA, Auwal MS, Sanda FA, Majama YB and Gambo BG: Phytochemical and *in vitro* antimicrobial properties of n-butanol seed extract of *Garcinia kola*. Journal of Molecular Pathophysiology 2013; 2(1): 11-15.

8. Alaje DO, Owolabi KT, Olakunle TP, Oluoti OJ and Adetuberu IA: Nutritional, minerals and phytochemicals composition of *Garcinia kola* [Bitter cola] and *Aframomum melegueta* [Alligator pepper]. IOSR Journal of Environmental Science, Toxicology and Food Technology 2014; 8(1): 86-91.
9. Durand D, Farid B, Adolphe A, Haziz S, Pacôme AN, Hubert A, Nafan D and Christine N: Phytochemical screening and biological activities of *Garcinia kola* (bark, leaves and seeds) collected in Benin. African Journal of Microbiology Research 2015; 9(28): 1716-1727.
10. PubMed Health, U.S. National Library of Medicine. How does the liver work? 2016. <https://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0072577/>
11. Cimeno A, French BM, Powell JM, Phelps C, Ayares D, O'Neill NA, Laird CT, Pierson RN, Azimzadeh AM, Barth RN and LaMattina JC: Synthetic liver function is detectable in transgenic porcine livers perfused with human blood. Xenotransplantation 2017; [Epub ahead of print]; doi: 10.1111/xen.12361.
12. Trease GE and Evans MC: Text Book of Pharmacognosy. Bailliere Tindal, London, Sixteenth Edition 2012.
13. Chen X, Zhang Y, Zhu Z, Liu H, Guo H, Xiong C, Xie K, Zhang X and Su S: Protective effect of berberine on doxorubicin induced acute hepatorenal toxicity in rats. Molecular Medicine Reports 2016; 13(5): 3953-3960.
14. Cachón AU, Quintal-Novelo C, Medina-Escobedo G, Castro-Aguilar G and Moo-Puc RE: Hepatoprotective effect of low doses of caffeine on CCl₄-induced liver damage in rats. Journal of Dietary Supplements 2017; 14(2): 158-172.
15. Milošević N, Milić N, Živanović-Bosić D, Bajkin I, Perčić I, Abenavoli L and Medić-Stojanoska M: Potential influence of the phthalates on normal liver function and cardio metabolic risk in males. Environmental Monitoring and Assessment 2017; 190(1): 17.
16. Singh A, Bhat TK and Sharma OP: Clinical biochemistry of hepatotoxicity. Journal of Clinical Toxicology 2011; S4: 001.
17. Adejoke EM, Akinrinade DI, Erommonselle E and Afodun AM: Hepatoprotective effects of *Garcinia kola* (Bitter kola) against paracetamol-induced oxidative damage and glycogen degranulation in hepatocytes of adult male Wistar rats. Journal of Advances in Biology and Biotechnology 2015; 3(3): 110-116.
18. Michael B, Yano B, Sellers RS, Perry R, Morton D, Roome N, Johnson JK and Schafer K: Evaluation of organ weights for rodent and non-rodent toxicity studies: A review of regulatory guidelines and a survey of current practices. Toxicologic Pathology 2007; 35: 742-750.
19. Lin JF, Lin YH, Liao PC, Lin YC, Tsai TF, Chou KY, Chen HE, Tsai SC and Hwang TI: Induction of testicular damage by daily methamphetamine administration in rats. The Chinese Journal of Physiology 2014; 57(1): 19-30.
20. Awounfack CF, Ateba SB, Zingue S, Mouchili OR and Njamen D: Safety evaluation (acute and sub-acute studies) of the aqueous extract of the leaves of *Myrianthus arboreus* P. Beauv. (Cecropiaceae) in Wistar rats. Journal of Ethnopharmacology 2016; 194: 169-178.
21. Hoffbrand VA and Moss PAH: Essential of Haematology. Wiley Blackwell Science, USA, Seventh Edition 2015.
22. Ahumibe AA and Braide VB: Effect of gavage treatment with pulverised *Garcinia kola* seeds on erythrocyte membrane integrity and selected haematological indices in male albino wistar rats. Nigerian Journal of Physiological Sciences 2009; 24(1): 47-52.
23. Uko OJ, Usman A and Ataja AM: Some biological activities of *Garcinia kola* in growing rats. Veterinarski Arhiv 2001; 71: 287-297.
24. Dada AA and Ikuerowo M: Effects of ethanolic extracts of *Garcinia kola* seeds on growth and haematology of catfish (*Clarias gariepinus*) broodstock. African Journal of Agricultural Research 2009; 4(4): 344-347.
25. Atsukwei D, Daniel EE, Odeh SO, Adams MD, Malgwi IS and Olih PC: Efficacy of *Garcinia kola* seed ethanolic extract on haematological parameters in male Wistar rats. Journal of Advances in Medical and Pharmaceutical Sciences 2015; 3(1): 10-13.

How to cite this article:

Aprioku JS: Investigation of the effects of ethanol seed extract of *Garcinia kola* on hematological and hepatorenal indices in wistar albino rats. Int J Pharm Sci & Res 2018; 9(3): 975-80. doi: 10.13040/IJPSR.0975-8232.9(3).975-80.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)