



Received on 28 July, 2017; received in revised form, 13 September, 2017; accepted, 17 September, 2017; published 01 May, 2018

CHARACTERISTIC AND OPTIMIZED USE OF BIOACTIVE COMPOUNDS FROM *GLORIOSA SUPERBA* AND *ALBIZIA AMARA* WITH APOPTOTIC EFFECT ON HEPATIC AND SQUAMOUS SKIN CARCINOMA

Fairen Angelin Jayakumar^{*}, Samson Eugin Simon, Yi Wei Yan and Pui Woon Yap

Department of Biosciences, School of Science and Engineering, Malaysia, University of Science and Technology (MUST) 12, Jalan PJU 5/5, Kota Damansara, 47810 Petaling Jaya, Selangor, Malaysia.

Keywords:

Antioxidants, Anticancer, *Gloriosa superba* Linn, *Albizia amara* (Roxb.) Boiv, Hepatic carcinoma, Squamous skin cell carcinoma

Correspondence to Author:

Fairen Angelin Jayakumar

Student, Master of Science in Biotechnology, Department of Biosciences, School of Science and Engineering, Malaysia University of Science and Technology(MUST) 12, Jalan PJU 5/5, Kota Damansara, 47810 Petaling Jaya, Selangor, Malaysia.

E-mail: fairenangelin.2806@gmail.com

ABSTRACT: Background: Cancer is still a dreadful disease, the treatment ranges with difficulties like strong side effects, shortage of donor and organs. In relation, plants and its chemical properties will provide an efficient source in the innocuous treatment for cancers. **Objectives:** To identify and isolate bioactive compounds from *Gloriosa superba* tubers and *Albizia amara* leaves with anti-cancer property that has traditionally been suggested as an anti-proliferative agent in ethnomedicine. **Design:** *G. superba* tubers and *A. amara* leaves are extracted and screened qualitatively. Extracts were then tested for antioxidant by DPPH assay. Highly potential extract with antioxidant are preceded to GCMS. The compounds determined from GCMS were selected according to its non-reported bioactive information on PubChem, and preceded against Hep-G2 and A431 cancer cell by MTT assay. **Result:** Ethanolic extract showed high antioxidant activity in both the plant extracts, GC-MS analysis was performed with ethanolic extracts revealing ample of phytoconstituent. Six non-reported bioactive compounds were isolated and identified using GC-MS. MTT assay was performed using isolated compounds such as 3-Hydroxy-4-methoxymandelic acid, 1-Butanone,1-(2, 4, 5 trihydroxy phenyl), 2H-1-Benzopyran, 3, 5, 6, 8 tetrahydro, β -Amyrin trimethylsilyl ether, Undecane, 2, 8-dimethyl and Octadecanoic, 2-oxo methyl ester on Hep-G2 and A431 cell lines. The optimized use of these compounds reveals apoptotic growth inhibition of 50.58% in A431 cell line and 53.42% in Hep-G2 cell line. **Conclusion:** The anti-cancer property of *G. superba* tubers and *A. amara* leaves as mentioned in various ethnopharmacological records, concurrently emphasize the possible bioactive compounds to treat the hepatic and squamous skin carcinoma

INTRODUCTION: Cancer is detected by medical imaging or biopsy and it is often treated with help of radiation, surgery and chemotherapy¹. Many complementary treatments have been introduced and its use has improved dramatically around the World².

People find that certain complementary methods are very useful to help control toxicity and improve the quality of their lives³. Hepatocellular carcinoma is a common type of primary liver cancer, commonly caused by cirrhosis and skin cancer is a destructive malignant cancer growth on skin^{4,5}. Chemo drug therapy were used to prevent the growth and metastasis of hepatic and skin cancers⁶, but the recovery rate stays low as these synthetic drugs cause side effects and metabolic suppression⁷. Many medicinal plants have used to treat the cancers and tumors in early days in Indian and Chinese tradition⁸.

QUICK RESPONSE CODE 	DOI: 10.13040/IJPSR.0975-8232.9(5).1769-78
	Article can be accessed online on: www.ijpsr.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.9(5).1769-78	

Based on ethnomedicine surveys, *Gloriosa superba* and *Albizia amara* is used to treat liver and skin cancer^{9,10}. Cancer growth is inhibited by the event of caspase-mediated cell death and it is also known as apoptosis¹¹. Apoptosis is a highly regulated traumatic cell death process caused by sudden cellular injury¹². The alternative products from plants provide us apoptotic induced cancer suppression with natural nontoxic metabolic activity¹³.

Gloriosa superba and *Albizia amara* are flowering plant and are rich in alkaloids and other essential compounds^{14,15}. These plants been used in the

traditional therapy for various ailments^{16,17} such as piles, ulcer, snake bite, acne, pimple, cancer, leprosy and also used as antibacterial, antifungal, astringent, antidiarrheal, anti-inflammatory, emetic and anti-cancer agent^{18,19}. *G. superba* tubers and *A. amara* leaves possess bio-active compounds involves extrinsic apoptotic pathway in cell signalling by ligand binding on the death receptors²⁰. Compounds obtained from *G. superba* tubers and *A. amara* leaves were focused in this study, as antiproliferative agents to prevent cancer cell growth and metastasis by activating caspases mediated apoptosis in squamous skin carcinoma and hepatic carcinoma cell lines²¹.




FIG. 1: RESEARCH DESIGN INVOLVED IN THE OPTIMIZED USE OF ANTIPROLIFERATIVE AGENTS FROM GLORIOSA SUPERBA AND ALBIZIA AMARA ACTING AS HEPATIC AND SQUAMOUS CELL SKIN CARCINOMA SUPPRESSOR

MATERIALS AND METHODS: *Albizia amara* leaves and *Gloriosa superba* tubers were collected from Ariyalur district, India. The collected samples were shade dried and pulverized individually. Soxhlet apparatus was setup manually and the thimble was filled with 150 gm of powdered samples. Solvents like ethanol (ET), aqueous (AQ) and benzene (BE) were used in extraction process. The solvent extracts were collected and stored in a standard method²². 1 gram of each extract was dissolved in 100 ml of respective solvents and used

further. The preliminary screening of phytochemicals was carried out using procedures, commonly employed with precipitation and coloration reactions to identify the occurrence of major secondary metabolites.

Antioxidant activity was determined for *G. superba* tubers (G) and *A. amara* leaves (A) extracts by DPPH assay²³, different concentration of 50 μ l, 100 μ l, 150 μ l, 200 μ l and 250 μ l samples were used respectively.

The sample concentrations were incubated with 3 ml of 0.1 mM DDPH for 30 minutes in dark. After incubation, the test concentrations were observed at 517 nm in triplicate values. The extracts with highest range of antioxidant activity on comparing with other solvent extracts were subjected to Gas chromatography Mass Spectrometry, GC-MS was carried out with Perkin Elmer Turbo Mass Spectrophotometer equipment's. Perkin Elmer Elite - 5 capillary columns was used, measures about 30 m × 0.25 mm with the thickness of 0.25 mm film. The carrier gas Helium was used, with the flow rate of 0.5 ml/min. The sample injection volume was 1ml. At first, the ethanolic extract of two different plants were loaded and programmed as 110 °C for 4 min, increases to 280 °C. The total run time was 90 min for each sample. The outcome were analyzed and could able to identify a six non-reported compounds from the *G. superba* tubers sample and *A. amara* leaves sample in a specific range of retention timing (RT). PUB-CHEM library was used to identify the bioactivity of the components.

Measurement of peak areas and data processing were carried out by Turbo-Mass OCPTVS-Demo SPL software. The selected compounds from the GC-MS analysis were separated particularly by flash chromatography and mixed together. The mixed samples were run again to confirm the presence of identified six non-reported compounds. The apoptotic effect of collected sample from *G. superba* tubers and *A. amara* leaves on Hep-G2 and A431 cell lines were determined by the MTT cytotoxicity assay²⁵. Hep-G2 cell lines [HEPG2] (ATCC® HB-8065™) and A431 cell lines (ATCC® CRL-1555™) were purchased from ATCC, USA and the sub culturing method was followed according to ATCC procedures. Cell lines were incubated and after 48 hrs incubation, the cell reaches the confluence. Cells (1 × 10⁵ / well) were plated in 1 ml of medium / well in 24 well plates along with samples of various concentrations²⁶. Then, the cells were incubated at 37 °C in the presence of various 5, 10, 25, 50 and 100 µg concentrations of bioactive isolates in 0.1% DMSO for 48 hrs. 200 µl (5 mg/ml) of 0.5% 3-(4, 5-dimethyl-2- thiazolyl)- 2, 5- diphenyl- tetrazolium bromide (MTT), phosphate-buffered saline solution was added to each well and incubated for 4 hrs. The viable cells were determined by the absorbance

at 570 nm. Observances were recorded and the concentration required for 50% inhibition of viability (GI₅₀) was determined graphically. DMSO was used as blank and the wells without sample containing cells were used as control.

The effect of the samples on the proliferation of Hep-G2 was expressed as the % cell viability, using the standard formula. The results were expressed as Mean ± standard deviation (SD) and statistically compared with control group or within the groups using triplicate values. For every assay, data were evaluated for separate three independent experiments. Dose-response curves were fitted using a nonlinear-regression sigmoidal dose-response curve model provided in the Graphpad Prism software (Graphpad software, Inc., San Diego, USA). IC₅₀ and GI₅₀ values were derived from fitted curves for each single experiment. Finally IC - values were calculated as average of three independent experiments with the standard deviation of the mean values. Data from the different assays were statistically compared using one way ANOVA analysis with Tukey's test (n = 5) as post-ANOVA analysis (p < 0.05).

RESULT: *G. superba* tubers and *A. amara* leaves were studied for the bioactives with antiproliferative potentiality. According to ethnomedicine survey, the leaves of *A. amara* and tubers of *G. superba* were focused in anticancer study. *A. amara* leaves and *G. Superba* tuber were extracted and qualitative occurrence of various phytochemical was performed by standard preliminary phytochemical screening procedure²⁷. Occurrence of various phytochemicals is mentioned in below **Table 1**.

Further, each extract were analyzed for antioxidant test by DPPH antiscavenging assay. The inhibition percentage of *A. amara* leaves and *G. superba* tubers exhibits triplicate value of absorbance using UV-spectrophotometry at 517 nm for 50, 100, 150, 200 and 250 µl concentration from both plant solvent extract as mentioned in **Table 2**. Values of DPPH assay was determined from standard sustained calculation using control value 1.341 ± 0.003 for *A. amara* leaves and 1.128 ± 0.001 for *G. superba* tubers. Among the three extract, AET (87.44 ± 0.172) and GET (92.19 ± 0.236) has the highest inhibition rate for 250 µl/ml.

TABLE 1: PRELIMINARY PHYTOCHEMICAL SCREENING OF *G. SUPERBA* TUBERS AND *A. AMARA* LEAVES

S. no.	Phytochemicals	Test Name	AAQ	AET	ABE	GAQ	GET	GBE
1	Alkaloids	Mayer's Test	-	++	++	++	++	++
2	Flavonoids	Shinoda Test	-	-	+	+	++	-
3	Glycosides	Keller-Killani Test	++	-	++	++	-	-
4	Saponins	Forth Test	++	+	-	++	-	++
5	Steroids	LB test	-	++	++	-	-	++
6	Phenols	Lead Acetate Test	++	++	++	-	++	++
7	Tannins	Ferric Chloride Test	++	++	+	++	+	++
8	Terpenoids	Salkowski's Test	++	++	++	-	++	++
9	Carbohydrates	Molisch's test	-	-	++	+	++	++
10	Proteins	Ninhydrin Test	++	-	++	++	-	++

Phytochemical screening for aqueous, ethanol and benzene solvent extract of *G. superba* tubers and *A. amara* leaves. ++ indicates presence, + indicates faint, - indicates absence


TABLE 2: ANTI - SCAVENGING ASSAY FOR *A. AMARA* LEAVES EXTRACT AND *G. SUPERBA* TUBERS EXTRACT

Concentration (µl/ml)	<i>A. amara</i> leaves extracts (Mean ± SD)			<i>G. superba</i> tubers extracts (Mean ± SD)		
	AAQ	AET	ABE	GAQ	GET	GBE
50µL	27.293±0.268	65.542±0.366	45.563±1.125	27.629±0.823	62.234±0.089	37.294±0.456
100 µL	48.941±0.598	84.514±0.282	57.917±0.973	44.002±0.144	89.953±0.285	46.068±0.545
150 µL	57.023±0.715	85.732±0.155	67.636±2.170	62.472±0.672	91.431±0.103	54.137±0.446
200 µL	66.293±0.596	86.742±0.052	74.969±1.528	65.575±0.183	91.786±0.050	55.939±0.154
250 µL	71.812±0.785	87.447±0.172	83.545±1.875	78.604±0.455	92.198±0.236	58.779±0.668
IC ₅₀ (µg/ml)	97.134	35.666	49.850	149.836	42.161	59.710


Anti-scavenging activity of aqueous, ethanol and benzene extracts of *A. amara* leaves and *G. superba* tubers were determined from triplicate absorbance value. Control OD value of 1.341 ± 0.003 was obtained for *A. amara* leaves and 1.128 ± 0.001 for *G. superba* tubers extracts


To understand the IC₅₀ of antioxidant, one way ANOVA non-linear line regression curve was made using 3D-PRISM software as showed in **Graph 1** and **2**. The AET showed IC₅₀ value of $35.666 \pm 1.473 \mu\text{g/ml}$ with hill coefficient of 12.690%, followed by ABE $49.850 \pm 0.450 \mu\text{g/ml}$ with the hill coefficient of 47.260% and for AAQ $97.134 \pm$

$1.271 \mu\text{g/ml}$ with the hill coefficient of 20.710%. IC₅₀ of GET as $42.161 \pm 0.350 \mu\text{g/ml}$ with the hill coefficient of 48.850% followed by GBE of $59.710 \pm 0.742 \mu\text{g/ml}$ with the hill coefficient of 44.850% and for GAQ $149.836 \pm 0.954 \mu\text{g/ml}$ with the hill coefficient of 50.320%.




GRAPH 1: IC₅₀ VALUES FOR AQUEOUS $97.134 \pm 1.271 \mu\text{g/ml}$ (*), ETHANOL $35.666 \pm 1.473 \mu\text{g/ml}$ (x) AND BENZENE $49.850 \pm 0.450 \mu\text{g/ml}$ (+) EXTRACT OF *A. AMARA* LEAVES





GRAPH 2: IC₅₀ VALUES FOR AQUEOUS 149.836 ± 0.954 µg/ml (*), ETHANOL 42.161 ± 0.350 µg/ml (x) AND BENZENE 59.710 ± 0.742 µg/ml (+) EXTRACT OF *G. SUPERBA* TUBERS



From the higher indication of antioxidant activity in the AET and GET, GC-MS analysis was done to identify the specific bio-active agents which perform antioxidants. The ethanol extract of both the plants were carried out for GC-MS and ample of compounds were obtained in TIC chromatogram. In a specific RT, the compounds were separated and collected. The bioactivity of the compounds was determined from the molecular database of PubChem.

The non-reported compounds were chosen to perform anticancer assay as mentioned in **Table 3**. Six compounds such as 3-Hydroxy-4-methoxy-mandelic acid (C1), 1-Butanone,1-(2,4,5 trihydroxy phenyl) (C2), 2H-1-Benzopyran, 3,5,6,8 tetrahydro (C3), β-Amyrin trimethylsilyl ether (C4), Undecane, 2, 8-dimethyl (C5) and Octadecanoic, 2-oxo methyl ester (C6) from ethanolic extract of both plant samples was isolated and obtained from collecting tube.

TABLE 3: GC-MS OF SIX NON - REPORTED BIOACTIVE COMPOUNDS

S. no.	Plant	RT	Name of the compound	Molecular formula	MW
1	<i>Gloriosa superba</i> and	11.038	3-Hydroxy-4-methoxymandelic acid	C ₉ H ₁₀ O ₅	198.17
2	<i>Albizia amara</i>	23.080	1-Butanone,1-(2,4,5 trihydroxy phenyl)	C ₁₀ H ₂₂ O	196.19
3		23.443	2H-1-Benzopyran, 3,5,6,8 tetrahydro	C ₁₃ H ₂₀ O	192.29
4		31.017	β-Amyrin trimethylsilyl ether	C ₃₃ H ₅₈ OSi	498.89
5		25.39	Undecane,2,8-dimethyl	C ₁₃ H ₂₈	184
6		26.77	Octadecanoic acid, 2-oxo methyl ester	C ₁₉ H ₃₆ O ₃	312

Gas Chromatography identification of six bioactive compounds obtained from analyzing the ethanolic extracts of *G. superba* tubers and *A. amara* leaves

MTT assay on Hepatic Carcinoma (Hep-G2) and Squamous Skin Carcinoma (A431) cell line was performed using C1, C2, C3, C4, C5 and C6 compounds as shown in **Fig. 2**. The cytotoxicity effect was observed as 5.568%, 9.049%, 29.775%,

40.448% and 50.58% in A431 cell line and 12.474%, 22.362%, 32.739%, 45.073% and 53.424% in Hep-G2 cell line for the respective concentration of 5, 10, 25, 50 and 100 µg/ml sample **Table 4**.

TABLE 4: CELL VIABILITY PERCENTAGE MTT ASSAY FOR A431 AND HEP-G2

Concentrations (µg/ml)	A431		Hep-G2	
	% of viability (Mean ± SD)	Death rate %	% of viability (Mean ± SD)	Death rate %
5	94.432±0.307	5.568	87.526±0.105	12.474
10	90.951±0.418	9.049	77.638±0.061	22.362
25	70.225±0.292	29.775	67.261±0.368	32.739
50	59.552±0.177	40.448	54.927±0.300	45.073
100	49.420±0.532	50.580	46.576±0.495	53.424
GI ₅₀	-	14.821	-	36.578

Cell viability and cell death percentage of six bioactive compounds against A431 cell line and Hep-G2 cell line; Mean values were obtained from triplicate absorbance reading, control value 0.862 OD observances was determined for A431 cell line and 0.954 OD observance for Hep-G2







FIG. 2: THE MTT ASSAY FOR SIX BIOACTIVE COMPOUNDS OBTAINED FROM *G. SUPERBA* TUBERS AND *A. AMARA* LEAVES AGAINST A431 AND Hep-G2 CELL LINE. CONTROL DETERMINES THE CANCER CELL PROLIFERATION. Hep-G2 AND A431 CELL LINE SHOWS APOPTOTIC BEHAVIOR IN DIFFERENT CONCENTRATION FROM 5, 10, 25, 50 AND 100 µg. AFTER THE INCUBATION OF 48 HOURS. THE HIGHEST CELL DEATH PERCENTAGE WAS OBSERVED IN 100 µg CONCENTRATION IN BOTH THE CELL LINES

GI₅₀ value for anti-cancer activity of six non-reported bioactive compounds was understood by using the 3D-PRISM software to draw one way ANOVA non-linear line regression curve $n = 5$, (p


< 0.05). GI₅₀ of 14.821 µg/ml in A431 cell line and 36.578 µg/ml in Hep-G2 cell line was determined as shown in **Graph 3** and **4**.



GRAPH 3: GI₅₀ VALUES 14.821 µg/ml FOR THE SIX BIOACTIVE COMPOUNDS AGAINST THE A431 CELL LINE



GRAPH 4: GI₅₀ VALUES 36.578 µg/ml FOR THE SIX BIOACTIVE COMPOUNDS AGAINST THE HEP-G2 CELL LINE



DISCUSSION: The preliminary phytochemical screening of *G. superba* tubers and *A. amara* leaves revealed bioactive rich compounds like alkaloids, flavonoids, phenols, saponins and glycosides. Therapeutic effects of the bioactives include prevention of cardiovascular diseases and anti-inflammatory, antimicrobial, and anticancer activity. The screened bio actives also known as primary antioxidants compounds, which has the ability to protect against the damage caused by oxidative stress induced by free radical²⁸. DPPH scavenging activity was performed as it is a widely used to measure the antioxidant property of the extracts.

In radical scavenging activity, the maximum inhibition occurs in the ethanolic extract of both the plants as the ethanol absorbs the large amount of phytochemicals during extraction process. Thus we understand *G. superba* tubers and *A. amara* leaves have a good potential of antioxidant property^{29, 30}. The DPPH scavenging activity is visualized as the purple solution turns yellow, the antioxidant

compounds from the plants extract reacts with unstable scavenging DPPH by donating extra electron to stabilize the medium. The stabilization causes the yellow colour indication shown in **Fig. 3**.

The GET and AET has reliable multiple bioactive compounds which is crucial to analyze for cancer preventing agent. Antioxidant is a precursor to prevent cancers and heart disease, compounds with good antioxidant can be used for GC-MS analysis. The result of GC-MS analysis reveals the presence of many bioactive components, these ample of compounds were studied precisely in various researches and the biological activities of those compounds were recorded in library.

Recorded actives of compounds provides their potentiality to prevent proliferation of cancer cell line by apoptosis, but some compounds were not yet tested and holds no record. Compounds with no report were chosen for anti-proliferative analysis of A431 skin cancer SCC cell line and Hep-G2 hepatic carcinoma cell line by MTT assay. On

analyzing the GC-MS result, we could able to collect the six bioactive compounds in particular

retention time obtained from TIC chromatogram Fig. 4.






FIG. 3: OBSERVATION OF SCAVENGING ACTIVITY OF EXTRACTS IN DIFFERENT CONCENTRATIONS 50 µL, 100 µL, 150 µL, 200 µL AND 250 µL. EXTRACTS WERE REPRESENTED AS GET(A), GAT (B), GBE (C), AET (D), AAQ (E) AND ABE (F)



A



B

FIG. 4: (A) TIC CHROMATOGRAM OF ETHANOLIC EXTRACT OF *G. SUPERBA* TUBERS AND *A. AMARA* LEAVES. (B) QUALITATIVE ANALYSIS OF CHROMATOGRAM OF SIX BIOACTIVE COMPOUNDS

MTT assay involves 24 well plate in which cancer cell line were cultivated and supplied with the six bioactive compounds shows cytotoxic activity against both cell lines. Bioactive compounds can be a novel product for apoptotic mechanism by stimulating cancer cell signalling pathways, this

study doesn't provides any signalling results, but yet it proves an isolated compound can provides us an anti-proliferative agent for cancer treatment³¹. Isolated compounds from *G. superba* tubers and *A. amara* leaves are identified and described as compound C1, C2, C3, C4, C5, C6 in Fig. 4.




FIG. 4: ISOLATED COMPOUNDS WERE DENOTED AS C1, C2, C3, C4, C5 AND C6

A431 and Hep-G2 cell lines lead apoptotic death visualized under microscopic view. Cell death was reported as the cells fails to retain it growth in media. Further studies can be assisted in cancer cell signaling pathways which provide dynamics of drug mechanism in apoptosis of hepatic carcinoma and squamous skin carcinoma cells.

CONCLUSION: According to ethno-pharmacological records, the ethanolic extract of *G. superba* tubers and *A. amara* leaves showed antioxidant property which can prevent the cancer by stabilizing the cells. Six isolated non-reported compounds such as 3-Hydroxy-4-methoxymandelic acid, 2H-1-Benzopyran, 3, 5, 6, 8 tetrahydro, β-

Amyrin trimethylsilyl ether, 3- Hydroxy- 4-methoxymandelic acid Undecane, 2, 8- dimethyl, Octadecanoic acid and 2-oxo methyl ester obtained from *G. superba* tubers and *A. amara* leaves possess anticancer property on Hep-G2 and A431 cell lines. Thus *G. superba* and *A. amara* can provide an alternative drug to treat the cancer as it has the potentiality to suppress the cancer cell growth.

ACKNOWLEDGEMENT: We would like to express our special thanks of gratitude to people who helped us and to Dr. Suresh M. Pharm., Ph.D Greens Med laboratory, India; for allowing us to pursue our cancer research.

CONFLICTS OF INTEREST: No competing financial interests exist.

REFERENCES:

1. Visvader JE and Lindeman GJ: Cancer stem cells in solid tumours: accumulating evidence and unresolved questions. *Nat Rev Cancer* 2008; 8: 755–68.
2. Langeveld NE, Stam H and Grootenhuis MA: Quality of life in young adult survivors of childhood cancer. *Support Care Cancer* 2002; 10(8): 579–600.
3. Mariotto AB, Rowland JH, Yabroff KR *et al.*: Long-term survivors of childhood cancers in the United States. *Cancer Epidemiol Biomarkers Prev.* 2009; 18(4): 1033–1040.
4. Anand P, Kunnumakara AB, Sundaram C, Harikumar KB, Tharakan ST, Lai OS and Aggarwal BB: Cancer is a Preventable Disease that Requires Major Lifestyle Changes. *Pharmaceutical Research* 2008; 25(9): 2097–2116.
5. Michael RS, Peter JC and Futreal PA: The cancer genome. *Nature* 2009; 458: 719-724.
6. Tulin E, Mehmet O and Cetin-Atalay R: Molecular Biology of Liver Cancer. *Reviews in Cell Biology and Molecular Medicine* 2015; 1: 206–243.
7. Simoes MC, Sousa JJ and Pais AA: Skin cancer and new treatment perspectives: a review. *Cancer Lett* 2015; 357(1): 8-42.
8. Darvesh AS, Aggarwal BB and Bishayee A: Curcumin and liver cancer: a review. *Curr Pharm Biotechnol* 2012; 13(1): 218-28.
9. Greenwell M and Rahman PKSM: Medicinal Plants: Their Use in Anticancer Treatment. *International journal of pharmaceutical sciences and research* 2015; 6(10): 4103-4112.
10. Lisa H: Golden age of melanoma therapy. *Nature Reviews Clinical Oncology* 2015; 12: 1.
11. Shigekazu N and Masato T: Programmed cell death and immune system. *Nature Reviews Immunology* 2017; 17: 333–340.
12. Susan E: Apoptosis: A Review of Programmed Cell Death. *Toxicologic Pathology* 2007; 35(4): 495–516.
13. Zhao H: Extrinsic and Intrinsic Apoptosis Signal Pathway Review, *Apoptosis and Medicine*, Dr. Tobias Ntuli (Ed.), In Tech 2012.
14. Padmapriya S, Rajamani K, Sathiyamurthy VA: Glory Lily (*Gloriosa superba* L.) - A Review. *International Journal of Current Pharmaceutical Review and Research* 2015; 7(1): 43-49.
15. Kumar SK and Sagar VG: Phytochemical and pharmacological evaluation of *Gloriosa superba*. *Journal of Drug Delivery and Therapeutics* 2015; 5(4): 27-42.
16. Kokila K, Priyadarshini SD and Sujatha V: Phytopharmacological Properties of *Albizia Species*: A Review. *Int J Pharm Pharm Sci* 2013; 5(3): 70-73.
17. Thippeswamy S, Mohana DC, Abhishek RU and Manjunath K: Inhibitory effect of alkaloids of *Albizia amara* and *Albizia saman* on growth and fumonisin B1 production by *Fusarium verticillioides*. *International Food Research Journal* 2015; 21(3): 947-952.
18. Devi NN and Femina W: GC-MS analysis of *Gloriosa superba* medicinal plant of Tamilnadu. *Journal of Pharmacy Research* 2012; 5(1): 343.
19. Elisha S, Michal L, Sarah S, Helena P, Elaine S and Lorberboum-Galski H: Evaluating Medicinal Plants for Anticancer Activity. *The Scientific World Journal* 2014; 721402: 12.
20. Hosseini A and Ghorbani A: Cancer therapy with phytochemicals: evidence from clinical studies. *Avicenna Journal of Phytomedicine* 2015; 5(2): 84–97.
21. Khalil DN, Smith EL, Brentjens RJ and Wolchok JD: The future of cancer treatment: immunomodulation, CARs and combination immunotherapy. *Nature Reviews Clinical Oncology* 2016; 13: 273–290.
22. Tiwari P, Kumar B, Kaur M, Kaur G and Kaur H: Phytochemical screening and Extraction: A Review. *Internationale Pharmaceutica Scientia* 2011; 1(1): 98-106.
23. Al-Rimawi F, Rishmawi S, Ariqat SH, Khalid MF, Warad I and Salah Z: Anticancer Activity, Antioxidant Activity, and Phenolic and Flavonoids Content of Wild *Tragopogon porrifolius* Plant Extracts. Evidence - Based Complementary and Alternative Medicine 2016.
24. Ezhilan BP and Neelamegam R: GC-MS analysis of phytocomponents in the ethanol extract of *Polygonum chinense* L. *Pharmacognosy Research* 2012; 4(1): 11-14.
25. Mosmann T: Rapid Colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983; 65: 55-63.
26. Sathishkumar J, Manoharan MS and Illanchezian S: *In-vitro* Antimicrobial and Antitumor Activities of *Stevia Rebaudiana* (Asteraceae) Leaf Extracts. *Tropical Journal of Pharmaceutical Research*, December 2008; 7(4): 1143-1149.
27. Mariappan SK: Phytochemical Screening of *Gloriosa superba* Linn. from different geographical positions. *International Journal of Scientific and Research Publications* 2013; 3(1): 1-5.
28. Sudarmani G: Identification of bioactive constituents of *Albizia amara* using FT-IR, HPTLC and GC-MS. *International Journal of Chemical and Pharmaceutical Analysis* 2016; 3: 1.
29. Manoharan S: "Antibacterial activity of 50 medicinal plants used in folk medicine." *Int. J. of Biosciences* 2013; 3(4): 104-121.
30. Saikat S *et al.*: "Free radicals, antioxidants, diseases and phytomedicines: Current status and future prospect." *International Journal of Pharmaceutical Sciences Review and Research* 2010; 3(1): 91-100.
31. Hillegass JM, Shukla A, Lathrop SA, MacPherson MB, Fukagawa NK and Mossman BT: Assessing nanotoxicity in cells *in vitro*. *Wiley interdisciplinary reviews Nanomedicine and nanobiotechnology* 2010; 2(3): 219-31. doi:10.1002/wnan.54.

How to cite this article:

Jayakumar FA, Simon SE, Wei YY and Woon YP: Characteristic and optimized use of bioactive compounds from *Gloriosa superba* and *Albizia amara* with apoptotic effect on hepatic and squamous skin carcinoma. *Int J Pharm Sci Res* 2018; 9(5): 1769-78. doi: 10.13040/IJPSR.0975-8232. 9(5):1769-78.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)