



Received on 18 September, 2017; received in revised form, 30 December, 2017; accepted, 05 January, 2018; published 01 June, 2018

CHILLING STRESS: HOW IT AFFECTS THE PLANTS AND ITS ALLEVIATION STRATEGIES

Kalpna Bhandari

Department of Botany, Panjab University, Chandigarh - 160014, Punjab, India.

Keywords:

Low temperature, Chilling sensitivity, Plant response, Cryoprotectants, Transgenic approach

Correspondence to Author:

Kalpna Bhandari

Research Fellow,
Department of Botany,
Panjab University, Chandigarh -
160014, Punjab, India.

E-mail: kalpna.bhandari@gmail.com

ABSTRACT: Low temperatures have been reported to be having far reaching effects on all plant species. The effects, in turn depend on the low temperature sensitivity of each and every plant species. In nature, the effects of various stresses may be seen overlapping thus the effects may be specific or non-specific. The effects in response to low temperature can be well ameliorated by the exogenous application of various protective molecules *viz.*, proline, glycine betaine, trehalose. Various cold tolerant transgenic crops over-expressing these molecules have also been developed.

INTRODUCTION: Every plant species has its own of temperature range within which it shows maximum growth and development and any deviation from this range, above or below it, hampers its overall performance. Sub-optimum temperatures have been proved to be having appreciably negative effects on various agricultural and horticultural crops resulting in heavy yield losses. Since cold stress invokes multiple effects and numerous complex mechanisms **Fig. 1** come into play in response to injuries and adaptations in cold-stressed plants¹³ thus, the overall effects have been rightly termed as a complex syndrome than a single response

Effects on Plants: The effects of cold stress, in turn depend on the low temperature sensitivity of each and every plant species.

Crops of tropical and sub-tropical origin, in particular, are more chilling sensitive *e.g.* rice (*Oryza sativa*), maize (*Zea mays*), chickpea (*Cicer sp.*). Other economically important crops suffering cold induced huge yield losses include soybean (*Glycine max* Linn.), cucurbits (*Cucurbita sp.*) and various cereal crops²⁵.

Though, the primary site of cold injury and hence electrolyte leakage is plasma membrane¹ in all the plants, however, in nature the effects in response to various stresses may be specific or non-specific *i.e.* they may overlap *e.g.* effects of cold and draught stress may overlap as shown in **Fig. 2**. To alleviate the harmful effects of cold stress, plants may accumulate various cryoprotectants. This article summarises the role of some of these protective molecules.

Proline: Proline is a well establish protective molecule and has been reported to impart protection against cold stress to many plants by not only maintaining its osmolarity but also acting as molecular chaperon thereby stabilizing the structure and hence functioning of crucial proteins and enzymes².

QUICK RESPONSE CODE 	DOI: 10.13040/IJPSR.0975-8232.9(6).2197-00
	Article can be accessed online on: www.ijpsr.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.9(6).2197-00	

It also protects the plant by maintaining the double stranded structure of genetic material *i.e.* DNA¹⁶ and by up-regulating the anti-oxidative stress machinery²³. Thus, if exogenously supplied, proline can well mitigate the harmful effects of cold stress as reported by Kumar¹⁴ in cold stressed chickpea.

Proline supplemented chickpea plants exhibited improved pollen viability, pollen germination and pollen tube growth along with better flower and pod retention. Similarly, proline conferred appreciable cold tolerance to rapeseed plants as well¹⁰.




FIG. 1: THE COLD STRESS SYNDROME OF PLANTS




FIG. 2: STRAINS PRODUCED BY COLD AND DROUGHT IN PLANT TISSUES

Glycine Betaine: Glicine betaine (GB) is a quaternary ammonium compound and has been proposed to be an effective cryoprotectant as it protects the cold stressed plant plants from photoinhibition induced damage to photosynthetic machinery⁴ by protecting the various components of PSII¹⁵ and also as a molecular chaperone. Exogenously supplied GB improved the performance of tomato plants by¹². Similarly, GB supplemented sweet pepper plants also performed better than untreated one under cold stress²⁸ which corroborated similar reportson wheat³¹ and other grain crops²⁵.

Trehalose: Trehalose, an alpha linked disaccharide, has been proposed to be playing protective roles under various abiotic stresses including cold and freezing⁹. There are numerous reports of trehalose levels going up under low temperatures rice in Arabidopsis²⁶ and rice²⁰. Thus, exogenously

supplied trehalose thus can well protect not only cold stressed plants as proved in Arabidopsis⁶ but also yeast⁷ and even of *Cryptococcus laurentii* and *Rhodotorula glutinis*¹⁷.

The protective effects of trehalose have been credited to its membrane stabilizing ability and depression in phase transition temperature of biomembranes and remains amorphous even under completely dehydrated conditions²⁴.

Transgenic Approach: The protective molecules, discussed above don't seem to accumulate in sufficient amounts so the exogenous treatment has been proposed or the plants may be even genetically modified to over-accumulate specific molecule which in turn can well impart cold tolerance to the targeted plant. Numerous such transgenic plants have been raised and summarized in the **Table 1** below.

TABLE 1: LIST OF TYRANGENIC PLANTS OVER-EXPRESSIG PROTECTIVE BIOMOLECULES

Transgenic plant	Biomolecule accumulated	References
<i>Arabidopsis thaliana</i>	Proline	(Zhu <i>et al.</i> , 2012) ³²
<i>Oryza sativa</i>	Proline	(Yang <i>et al.</i> , 2012) ²⁹
<i>Nicotiana tabacum</i>	Proline	(Pan <i>et al.</i> , 2012) ²¹
<i>Brassica napus</i>	Proline	(Gomaa <i>et al.</i> , 2012) ⁵
<i>Brassica campestris</i>	Glycine Betaine	(Wang <i>et al.</i> , 2010) ²⁷
<i>Triticum aestivum</i>	Glycine Betaine	(Zhang <i>et al.</i> , 2010) ³¹
<i>Ipomea batatas</i>	Glycine Betaine	(Fan <i>et al.</i> , 2012) ³
<i>Xenopus laevis</i>	Trehalose	(Li <i>et al.</i> , 2009) ¹⁸
<i>Oryza sativa</i>	Trehalose	(Hossain <i>et al.</i> , 2010) ⁸
<i>Nicotiana tabacum</i>	Trehalose	(Yang <i>et al.</i> , 2010) ³⁰
<i>Hordeum vulgare</i>	Trehalose	(Ligaba <i>et al.</i> , 2011) ¹⁹
<i>Arabidopsis thaliana</i>	Trehalose	(Kang <i>et al.</i> , 2011) ¹¹
<i>Oryza sativa</i>	Trehalose	(Li <i>et al.</i> , 2011) ²⁰

CONCLUSION: Cold stress affects plants at all levels and the effects of cold stress are multiple and far reaching. This review thus focuses on the use of some of promising cryoprotetants in this regard. These biomolecules not only maintain the osmotic concentration of the cell sap but also protect and maintain the structure and function of various crucial enzymes and proteins besides preserving the integrity of various biomembranes.

ACKNOWLEDGEMENT: The technical support and access to e-journals provided by the Department of Botany, Panjab University, Chandigarh is highly acknowledged.

CONFLICT OF INTEREST: Corresponding author is the sole author hence no conflict of interest.

REFERENCES:

- Bhandari K and Nayyar H: Low temperature stress in plants: an overview of roles of cryoprotectants in defense. Springer, New York, Edition 3rd, Vol 1, 2014: 193-265.
- Diehl RC, Guinn EJ, Capp MW, Tsodikov OV and Record Jr. MT: Quantifying additive interactions of the osmolyte proline with individual functional groups of proteins: comparisons with urea and glycine betaine, interpretation of m-values. *Biochemistry* 2013; 52(35): 5997-6010.
- Fan W, Zhang M, Zhang H and Zhang P: Improved tolerance to various abiotic stresses in transgenic sweet potato (*Ipomoea batatas*) expressing spinach betaine aldehyde dehydrogenase. *PLoS One* 2012; 7: e37344.
- Giri J: Glycine betaine and abiotic stress tolerance in plants. *Plant Signal Behaviour* 2011; 6: 1746-1751.
- Gomaa AM, Raldugina GN, Burmistrova NA, Radionov NV, Kuznetsov VV: Response of transgenic rape plants bearing the Osmyb4 gene from rice encoding a trans-factor to low above-zero temperature. *Russian Journal of Plant Physiol* 2012; 59: 105-114.

6. Hanhong B, Eliot H and Richard S: Exogenous trehalose promotes non-structural carbohydrate accumulation and induces chemical detoxification and stress response proteins in *Arabidopsis thaliana* grown in liquid culture. *Plant Science* 2005; 168: 1293-1301.
7. Hirasawa R, Yokoigawa K, Isobe Y and Kawai H: Improving the freeze tolerance of baker's yeast by loading with trehalose. *Bioscience Biotechnology and Biochemistry* 2001; 65: 522-526.
8. Hossain MA, Cho JI, Han M, Ahn CH, Jeon JS, An G and Park PB: The ABRE-binding bZIP transcription factor OsABF2 is a positive regulator of abiotic stress and ABA signaling in rice. *Journal of Plant Physiology* 2010; 167: 1512-1520.
9. Iordachescu M and Imai R: Trehalose and abiotic stress in biological systems. In *Tech, Croatia*, 2011: 215-234.
10. Jonytiene V, Burbulis N, Kupriene R and Blinstrubiene A: Effect of exogenous proline and de-acclimation treatment on cold tolerance in *Brassica napus* shoots cultured *in vitro*. *Journal of Food Agriculture and Environment* 2012; 10(1): 327-330.
11. Kang HG, Kim J, Kim B, Jeong H, Choi SH, Kim EK, Lee HY and Lim PO: Over-expression of FTL1/DDF1, an AP2 transcription factor, enhances tolerance to cold, drought, and heat stresses in *Arabidopsis thaliana*. *Plant Science* 2011; 180: 634-641.
12. Karabudak T, Bor M, Özdemir F and Türkan I: Glycine betaine protects tomato (*Solanum lycopersicum*) plants at low temperature by inducing fatty acid desaturase7 and lipoxygenase gene expression. *Molecular biology reports*, 2014; 41(3): 1401-1410.
13. Kazan K: Diverse roles of jasmonates and ethylene in abiotic stress tolerance. *Trends in plant science*, 2015; 20(4): 219-229.
14. Kumar S, Malik J, Thakur P, Kaistha S, Sharma KD, Upadhyaya HD, Berger JD and Nayyar H: Growth and metabolic responses of contrasting chickpea (*Cicer arietinum* L.) genotypes to chilling stress at reproductive phase. *Acta Physiologiae Plantarum* 2011; 33: 779-787.
15. Kurepin LV, Ivanov A, Zaman M, Pharis RP, Hurry V and Hüner NP: Interaction of glycine betaine and plant hormones: protection of the photosynthetic apparatus during abiotic stress. Springer International Publishing, Cham, 2017: 185-202.
16. Law JA and Jacobsen SE: Establishing, maintaining and modifying DNA methylation patterns in plants and animals. *Nature Reviews Genetics* 2010; 11(3): 204-220.
17. Li BQ, Zhou ZW and Tian SP: Combined effects of endo- and exogenous trehalose on stress tolerance and biocontrol efficacy of two antagonistic yeasts. *Biological Control* 2008; 46: 187-193.
18. Li DD, Tai FJ, Zhang ZT, Li Y, Zheng Y, Wu YF and Li XB: cotton gene encodes a tonoplast aquaporin that is involved in cell tolerance to cold stress. *Gene* 2009; 438: 26-32.
19. Ligaba A, Katsuhara M, Shibasaka M and Djira G: Abiotic stresses modulate expression of major intrinsic proteins in barley (*Hordeum vulgare*). *Comptes Rendus Biologies* 2011; 334: 127-139
20. Li HW, Zang BS, Deng XW and Wang XP: Over-expression of the trehalose-6-phosphate synthase gene OsTPS1 enhances abiotic stress tolerance in rice. *Planta* 2011; 234: 1007-1018.
21. Pan J, Zhang M, Kong X, Xing X, Liu Y, Zhou Y, *et al.*: ZmMPK17, a novel maize group D MAP kinase gene, is involved in multiple stress responses. *Planta* 2012; 235: 661-676.
22. Pramanik MHR and Imai R: Functional identification of a trehalose 6-phosphate phosphatase gene that is involved in transient induction of trehalose biosynthesis during chilling stress in rice. *Plant Molecular Biology* 2005; 58: 751-762.
23. Rejeb KB, Abdelly C and Savaouré A: How reactive oxygen species and proline face stress together. *Plant Physiology and Biochemistry* 2014; 80: 278-284.
24. Sundaramurthi P, Patapoff TW and Suryanarayanan R: Crystallization of trehalose in frozen solutions and its phase behavior during drying. *Pharmaceutical Research* 2010; 27: 2374-2383.
25. Thakur P, Kumar S, Malik JA, Berger JD and Nayyar H: Cold stress effects on reproductive development in grain crops: An overview. *Environmental and Experimental Botany* 2010; 67: 429-443.
26. Veyres N, Aono M, Sangwan-Norreel BS and Sangwan RS: The *Arabidopsis* sweetie mutant is affected in carbohydrate metabolism and defective in the control of growth, development and senescence. *Plant Journal* 2008; 55: 665-686.
27. Wang QB, Xu W, Xue QZ and Su WA: Transgenic *Brassica chinensis* plants expressing a bacterial codA gene exhibit enhanced tolerance to extreme temperature and high salinity. *Journal of Zhejiang University Science B* 2010; 11: 851-861.
28. Wang Q, Ding T, Zuo J, Gao L and Fan L: Amelioration of postharvest chilling injury in sweet pepper by glycine betaine. *Postharvest Biology and Technology* 2016; 112: 114-120.
29. Yang A, Dai XY and Zhang WH: A R2R3-type MYB gene, OsMYB2, is involved in salt, cold, and dehydration tolerance in rice. *Journal of Experimental Botany* 2012; 63: 2541-2556.
30. Yang JS, Wang R, Meng JJ, Bi YP, Xu PL, Guo F, Wan SB, He QH and Li XG: Over-expression of *Arabidopsis* CBF1 gene in transgenic tobacco alleviates photo-inhibition of PSII and PSI during chilling stress under low irradiance. *Journal of Plant Physiology* 2010; 167: 534-539
31. Zhang XY, Liang C, Wang GP, Luo Y and Wang W: The protection of wheat plasma membrane under cold stress by glycine betaine overproduction. *Biologia Plantarum*, 2010; 54(1): 83-88.
32. Zhu B, Peng RH, Xiong AS, Xu J, Fu XY, Zhao W *et al.*: Transformation with a gene for myo-inositol O-methyl transferase enhances the cold tolerance of *Arabidopsis thaliana*. *Biologia Plantarum* 2012; 56: 135-139.

How to cite this article:

Bhandari K: Chilling stress: how it affects the plants and its alleviation strategies. *Int J Pharm Sci Res* 2018; 9(6): 2197-00. doi: 10.13040/IJPSR.0975-8232.9(6).2197-00.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)