



Received on 19 April 2019; received in revised form, 24 August 2019; accepted, 01 September 2019; published 01 February 2020

## ANTIBIOTIC SUSCEPTIBILITY PATTERN OF ESBL PRODUCING BACTERIA FROM URINARY TRACT INFECTION (UTI)

Juhi Hattewar<sup>\*</sup>, Akshma Koul and Abhishek Mathur

NCS Green Earth Pvt. Ltd., M-7, MIDC Hingna Industrial Area, Nagpur - 440016, Maharashtra, India.

### Keywords:

ESBL producing bacteria,  
Urinary tract infection, Antibiotic  
susceptibility, Antibiotic resistance

### Correspondence to Author:

**Juhi Hattewar**

Junior Scientist (R&D),  
NCS Green Earth Pvt. Ltd., M-7,  
MIDC Hingna Industrial Area,  
Nagpur - 440016, Maharashtra, India.

**E-mail:** hattewarjuhi@gmail.com

**ABSTRACT:** Antibiotics inhibit bacterial cell wall synthesis, protein synthesis or nucleic acid replication. The antibiotic must have access to and bind to its bacterial target site. Whether antibiotic resistance is intrinsic or acquired, the genetic determinants of resistance encode specific biochemical resistance mechanism that may include enzymatic inactivation of the drug, alteration to the structure of the antibiotic target site and changes that prevent access of an adequate concentration of the antimicrobial agents to the active site. Klebsiella organisms are often resistant to multiple antibiotics. Current evidence implicates plasmids as the primary source of the resistance genes. The susceptibility pattern of different antibiotics to ESBL producing *Escherichia coli* and Klebsiella spp. isolated from urine samples and they found that most of the infection caused by *E. coli* (61.29%) found to be most common bacteria in urinary tract infection followed by Klebsiella spp. (45.16%). Isolates were highly resistance to gentamicin followed by tetracycline, ampicillin, and amikacin. gentamicin showed resistance against *E. coli* and Klebsiella spp were 89.47% and 85.71% respectively. While tetracyclin and ampicillin showed 80% and 78% resistance to *E. coli* and Klebsiella spp. The isolates were highly susceptible and least susceptible to ciprofloxacin and norfloxacin against *E. coli* and Klebsiella spp. These antibiotics are considered as appropriate antimicrobials for empirical treatment of urinary tract infections. Most of the ESBL A producing isolates were multidrug-resistant. During present study, when urine sample inoculated on selective media like EMB, CLED, MSA, PIA and CIA, the growth of isolates on the medium were obtained as *E. coli*, *Klebsiella pneumoniae*, *S. aureus*, and *Pseudomonas aeruginosa*.

**INTRODUCTION:** Antibiotics susceptibility tests including, ESBL screening and confirmation, were carried out by disc diffusion technique using Clinical Laboratory Standard Institute (CLSI) criteria. Ten different types of bacteria genera were observed from nine different clinical samples.

*E. coli* was the most frequently isolated bacteria (30.5%) followed by *Staphylococcus aureus* (21.3%). ESBL producers showed high-level resistance against the quinolones, aminoglycoside, and cotrimoxazole but were sensitive to carbapenems and levofloxacin.

Non-ESBL organisms showed increased resistance to amoxicillin-clavulanate, ceftazidime, cotrimoxazole, tetracycline and amoxicillin. The prevalence of ESBL producers was 12.8%. *E. coli*, *Klebsiella pneumoniae*, and *Pseudomonas aeruginosa* showed ESBL production of 17.3, 14.9 and 10.0%, respectively<sup>1</sup>.

	<b>QUICK RESPONSE CODE</b> DOI: 10.13040/IJPSR.0975-8232.11(2).799-10
	This article can be accessed online on <a href="http://www.ijpsr.com">www.ijpsr.com</a>
DOI link: <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.11(2).799-10">http://dx.doi.org/10.13040/IJPSR.0975-8232.11(2).799-10</a>	

Antimicrobial susceptibility pattern and ESBL prevalence in *Klebsiella pneumoniae* from urinary tract infections. The study included 92 *Klebsiella pneumoniae*, isolated from the urine of patients suffering from UTIs. Antibiotic susceptibility testing and ESBL detection were carried out according to Clinical and Laboratory Standards Institute (CLSI) criteria. Eighteen antibiotics were tested in this study. A total of 54 (58.7%) isolates were found to be ESBL producers<sup>2</sup>. The detection of ESBL producing *E. coli* isolated from clinical samples.

This study was to determine the prevalence of ESBL isolation and identification of ESBL producing *E. coli* isolation and its confirmation via different confirmative tests such as ESBLs screening test. Double disc synergy test (DDST) and double-disc diffusion test (Phenotypic confirmatory test). In antibiotic susceptibility test the antibiogram revealed that 4 (6.15%) isolates were resistance to amikacin, (4.61%) were resistance to amoxycylav, 36 (55.38%) were resistant to ampicillin, gentamycin, 8 (12.30%) were resistant to levofloxacin, 5 (7.69%) were resistance to nalidixic acid, 25 (38.46%) were resistant to norfloxacin, 10 (15.38%) isolates were resistant to ofloxacin, 8 (12.30%) isolates were resistant to piperacillin. The ESBL percentage was found to be more in female urine samples<sup>3</sup>.

In the present investigation, antibiotic susceptibility pattern of ESBL producing bacteria from urinary tract infection.

## MATERIALS AND METHODS:

**Collection of Microorganism:** The urine sample is collected from the pathological laboratory Nagpur.

### Isolation and Identification of Bacteria:

- Urine samples are stricked on different selective media's.
- Plates were incubated at 37 °C for 24 h.
- After 24 h plates were observed for colonies.

### Morphological Characteristics:

- **Gram Character Examination:** For the study of gram character a well-isolated colony was selected by picking the colony with inoculating

loop smear was prepared on clean slide. A staining technique was performed as per standard procedure. Shape, size, gram character was then observed.

- **Motility:** Bacterial motility is an important part of making final identification. Motility of the given isolates was observed by using hanging drop method.
- **Biochemical Characteristics:** Various biochemical tests unique to each organism were performed by inoculating small portion of the well isolated colony into series of media. This include following tests.
- **IMViC Reactions:** IMViC reactions are a set of four useful reactions that are commonly employed in the identification of members of family *Enterobacteriaceae*. The four reactions are Indole test, methyl red test, Voges proskauer test, and citrate utilization test.

**Indole Test:** Bacterium to be tested is inoculated in peptone water, which contains amino acid tryptophan and incubated overnight at 37 °C. Following incubation few drops of Kovac's reagent are added. Kovac's reagent consists of para-dimethyl amino benzaldehyde, isoamyl alcohol, and con. HCl. Ehrlich's reagent is more sensitive in detecting indole production in anaerobes and non-fermenters.

Formation of a red or pink colored ring at the top is taken as positive. Example: *Escherichia coli*: positive; *Klebsiella pneumoniae*: negative.

### Methyl Red Test:

**Procedure:** The bacterium to be tested is inoculated into glucose phosphate broth, which contains glucose and a phosphate buffer and incubated at 37 °C for 48 h. Over the 48 h the mixed-acid producing organism must produce sufficient acid to overcome the phosphate buffer and remain acid.

The pH of the medium is tested by the addition of 5 drops of MR reagent. Development of red color is taken as positive. MR negative organism produces a yellow color. Example: *Escherichia coli*: positive; *Klebsiella pneumoniae*: negative.

**Voges Proskauer (VP) Test:**

**Procedure:** Bacterium to be tested is inoculated into glucose phosphate broth and incubated for at least 48 h. 0.6 ml of alpha-naphthol is added to the test broth and shaken. 0.2 ml of 40% KOH is added to the broth and shaken. The tube is allowed to stand for 15 min. The appearance of the red color is taken as a positive test.

The negative tubes must be held for one hour since maximum color development occurs within one hour after addition of reagents. Examples: *Escherichia coli*: negative; *Klebsiella pneumoniae*: positive.

**Citrate Utilization Test:**

**Procedure:** Bacterial colonies are picked up from a straight wire and inoculated into the slope of Simmon's citrate agar and incubated overnight at 37 °C. If the organism has examples: *Escherichia coli*: negative; *Klebsiella pneumoniae*: positive.

**Triple Sugar Iron Test:** The peptone mixture and the Beef and Yeast extracts provide the nutrients essential for growth. Sodium chloride maintains the osmotic balance of the medium. The Bacteriological agar is the solidifying agent.

**Procedure:**

1. Sterilize the inoculating needle in the blue flame of the bunsen burner till red hot and then allowed to cool.
2. From the rack, take the Trypticase soy broth tube containing the 24-48 h culture, remove the cap and flame the neck of the tube.
3. Using aseptic technique, take the culture of the organism from the TSB (tryptic soy broth) tube with the needle.
4. Again flame the neck of the tube and replace the tube in the test tube rack.
5. Take a sterile TSI slant tube from the rack, remove the cap and flame the neck of the tube.
6. Stab the needle containing the pure culture into the medium, up to the butt of the TSI tube, and then streak the needle back and forth along the surface of the slant.

7. Again flame the neck of the TSI tube, cap it and place it in the test tube rack.

8. Incubate at 37 °C for 18 to 24 h.

**Catalase Test:**

1. Take 3% of H<sub>2</sub>O<sub>2</sub> in a clean test tube.
2. Take loopful of culture and insert it in the test tube.
3. Formation of bubbles within 10 sec indicates a positive test.

**Urease Test:**

1. Prepare Christensen's urea agar slants. Streak the slant with the inoculum.
2. Incubate the test tubes at 37 °C for 24 h.
3. After incubation observe the tubes. A pink color indicates positive test reaction.

**Oxidase Test:**

1. Soak a small piece of filter paper in 1% Kovacs oxidase reagent and let dry.
2. Use a loop and pick a well-isolated colony from a fresh (18- to 24-h culture) bacterial plate and rub onto treated filter paper.
3. Observe for color changes.

**Antibiotic Susceptibility Test:** Once a culture is established, there are two possible ways to get an antibiogram:

- A semi-quantitative way based on diffusion (Kirby-Bauer method); small discs containing different antibiotics, or impregnated paper discs, are dropped in different zones of the culture on an agar plate, which is a nutrient-rich environment in which bacteria can grow. The antibiotic will diffuse in the area surrounding each tablet, and a disc of bacterial lysis will become visible. Since, the concentration of the antibiotic was the highest at the center, and the lowest at the edge of this zone, the diameter is suggestive for the Minimum Inhibitory Concentration, or MIC, (conversion of the diameter in millimeter to

the MIC, in  $\mu\text{g/ml}$ , is based on known linear regression curves).

- A quantitative way based on dilution: a dilution series of antibiotics is established (this is a series of reaction vials with progressively lower concentrations of antibiotic substance). The last vial in which no bacteria grow contains the antibiotic at the Minimal Inhibiting Concentration.

**Kirby Diffusion Method:** The agar diffusion test (Kirby–Bauer antibiotic testing, KB testing, or disc diffusion antibiotic sensitivity testing) is a test of the antibiotic sensitivity of bacteria. It uses antibiotic-impregnated wafers to test the extent to which bacteria are affected by those antibiotics. In this test, wafers containing antibiotics are placed on an agar plate where bacteria have been placed, and the plate is left to incubate. If an antibiotic stops the bacteria from growing or kills the bacteria, there will be an area around the wafer where the bacteria have not grown enough to be visible. This is called a zone of inhibition.

The size of this zone depends on how effective the antibiotic is at stopping the growth of the bacterium. A stronger antibiotic will create a larger zone because a lower concentration of the antibiotic is enough to stop growth. The bacteria in question are swabbed uniformly across a culture plate. A filter-paper disk, impregnated with the compound to be tested, is then placed on the surface of the agar. The compound diffuses from the filter paper into the agar. The concentration of the compound will be highest next to the disk and will decrease as distance from the disk increases. If the compound is effective against bacteria at a certain concentration, no colonies will grow where the concentration in the agar is greater than or equal to the effective concentration. This is the zone of inhibition. These along with the rate of antibiotic diffusion are used to estimate the bacteria's sensitivity to that particular antibiotic. In general, larger zones correlate with smaller minimum inhibitory concentration (MIC) of antibiotics for that bacterium. Inhibition produced by the test is compared with that produced by known concentration of a reference compound. This information can be used to choose appropriate antibiotics to combat a particular infection<sup>4-8</sup>.

**Procedure:** All aspects of the Kirby-Bauer procedure are standardized to ensure consistent and accurate results. Because of this, a laboratory must adhere to these standards. The media used in Kirby-Bauer testing must be Mueller-Hinton agar at only 4 mm deep, poured into either 100 mm or 150 mm Petri dishes. The pH level of the agar must be between 7.2 and 7.4. A close-up looks at the results of an agar diffusion test. Inoculation is made with a broth culture diluted to match a 0.5 McFarland turbidity standard, which is roughly equivalent to 150 million cells per ml<sup>9-13</sup>.

#### Incubation Procedure:

1. Using an aseptic technique, place a sterile swab into the broth culture of a specific organism and then gently remove the excess liquid by gently pressing or rotating the swab against the inside of the tube.
2. Using the swab, streak the Mueller-Hinton agar plate to form a bacterial lawn.
  - To obtain uniform growth, streak the plate with the swab in one direction, rotate the plate 90° and streak the plate again in that direction.
  - Repeat this rotation 3 times.
  - Allow the plate to dry for approximately 5 minutes.
1. Use an Antibiotic Disc Dispenser to dispense discs containing specific antibiotics onto the plate.
2. Using a flame-sterilized forceps, gently press each disc to the agar to ensure that the disc is attached to the agar.
3. Plates should be incubated overnight at an incubation temperature of 37 °C (98.6 °F)<sup>14-16</sup>.

**E–Strip Method:** The E - test principle was developed to standardize the disc diffusion method and to improve its reproducibility and reliability for clinical susceptibility predictions. The inhibition zone sizes from disc test results were compared to Minimum Inhibitory Concentration (MIC) values based on the reference agar dilution procedure.

The correlation between zone sizes and MIC values was then assessed using regression analysis and regression lines were used for extrapolating zone interpretive limits that corresponded to the MIC breakpoint values that defined susceptible, intermediate and resistant categorical results<sup>17-28</sup>.

## RESULTS:

**Isolation and Characterization of Microbes from Clinical Specimens:** After inoculation on EMB agar plates were observed for growth forms green metallic sheen. Similarly on MSA agar plates

were observed for growth observed as golden yellow, smooth colonies along with the production of acid which turns the media color yellow. On PIA plates showed bluish-green, pigmented colonies were observed. On Mac Conkey agar bluish color colonies observed. The results are shown in **Table 1-13; Fig 1-14**.

**Characterization of Isolates:** The clinical isolates were characterized on the basis of microscopic and biochemical test various test performed were shown in tables.

**Microscopic and Biochemical Characterization of Clinical Isolates:** Total no. of samples (n) = 8.

**TABLE 1: MORPHOLOGICAL AND BIOCHEMICAL CHARACTERISTICS OF URINE SAMPLE 1**

S. no.	Bacterial Isolates	Sugar Fermentation				IMVIC				Identified Bacteria
		lactose	Glucose	Mannitol	Sucrose	Indole	MR	VP	C	
1	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	+ve	+ve	-ve	-ve	<i>E. coli</i>
2	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	-ve	-ve	+ve	-ve	<i>Klebsiella pneumonia</i>
3	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	-ve	-ve	-ve	+ve	<i>S. aureus</i>
4	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	+ve	-ve	+ve	+ve	<i>Pseudomonas aeruginosa</i>

On the basis of the above results, it can be said that the clinical isolates were found to be *Escherichia coli*, *Klebsiella pneumonia*, *S. aureus*, and *Pseudomonas aeruginosa*.

**TABLE 2: MORPHOLOGICAL AND BIOCHEMICAL CHARACTERISTICS OF URINE SAMPLE 2**

S. no.	Bacterial Isolates	Sugar Fermentation				IMVIC				Identified Bacteria
		lactose	Glucose	Mannitol	Sucrose	Indole	MR	VP	C	
1	US -2	Acid +gas	Acid +gas	Acid +gas	Acid +gas	-ve	-ve	+ve	-ve	<i>Klebsiella Pneumonia</i>
2	US -2	Acid +gas	Acid +gas	Acid +gas	Acid +gas	-ve	-ve	-ve	+ve	<i>S. aureus</i>
3	US -2	Acid +gas	Acid +gas	Acid +gas	Acid +gas	-ve	-ve	-ve	+ve	<i>S. aureus</i>
4	US -2	Acid +gas	Acid +gas	Acid +gas	Acid +gas	+ve	-ve	+ve	+ve	<i>Pseudomonas aeruginosa</i>

On the basis of above result, it can be said that, the clinical isolates were found to be *Klebsiella pneumonia*, *S. aureus* and *Pseudomonas aeruginosa*.

**TABLE 3: MORPHOLOGICAL AND BIOCHEMICAL CHARACTERISTICS OF URINE SAMPLE 3**

S. no.	Bacterial Isolates	Sugar Fermentation				IMVIC				Identified Bacteria
		lactose	Glucose	Mannitol	Sucrose	Indole	MR	VP	C	
1	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	+ve	+ve	-ve	-ve	<i>E. coli</i>

On the basis of above result, it can be said that the clinical isolate was found to be *Escherichia coli*.

**TABLE 4: MORPHOLOGICAL AND BIOCHEMICAL CHARACTERISTICS OF URINE SAMPLE 4**

S. no.	Bacterial Isolates	Sugar Fermentation				IMVIC				Identified Bacteria
		lactose	Glucose	Mannitol	Sucrose	Indole	MR	VP	C	
1	US -2	Acid +gas	Acid +gas	Acid +gas	Acid +gas	-ve	-ve	-ve	+ve	<i>S. aureus</i>
2	US -2	Acid +gas	Acid +gas	Acid +gas	Acid +gas	+ve	-ve	+ve	+ve	<i>Pseudomonas aeruginosa</i>

On the basis of above result, it can be said that the clinical isolates were found to be *S. aureus* and *Pseudomonas aeruginosa*.

**TABLE 5: MORPHOLOGICAL AND BIOCHEMICAL CHARACTERISTICS OF URINE SAMPLE 5**

S. no.	Bacterial Isolates	Sugar Fermentation				IMVIC				Identified Bacteria
		lactose	Glucose	Mannitol	Sucrose	Indole	MR	VP	C	
1	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	+ve	+ve	-ve	-ve	<i>E. coli</i>
2	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	-ve	-ve	+ve	-ve	<i>Klebsiella pneumoniae</i>
3	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	-ve	-ve	-ve	+ve	<i>S. aureus</i>
4	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	+ve	-ve	+ve	+ve	<i>Pseudomonas aeruginosa</i>

On the basis of the above results, it can be said that the clinical isolates were found to be *Escherichia coli*, *Klebsiella pneumoniae*, *S. aureus*, and *Pseudomonas aeruginosa*.

**TABLE 6: MORPHOLOGICAL AND BIOCHEMICAL CHARACTERISTICS OF URINE SAMPLE 6**

S. no.	Bacterial Isolates	Sugar Fermentation				IMVIC				Identified Bacteria
		lactose	Glucose	Mannitol	Sucrose	Indole	MR	VP	C	
1	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	+ve	-ve	+ve	+ve	<i>Pseudomonas aeruginosa</i>

On the basis of the above results, it can be said that the clinical isolates were found to be *Pseudomonas aeruginosa*.

**TABLE 7: MORPHOLOGICAL AND BIOCHEMICAL CHARACTERISTICS OF URINE SAMPLE 7**

S. no.	Bacterial Isolates	Sugar Fermentation				IMVIC				Identified Bacteria
		lactose	Glucose	Mannitol	Sucrose	Indole	MR	VP	C	
1	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	-ve	-ve	-ve	+ve	<i>S. aureus</i>

On the basis of the above results, it can be said that the clinical isolates were found to be *S. aureus*.

**TABLE 8: MORPHOLOGICAL AND BIOCHEMICAL CHARACTERISTICS OF URINE SAMPLE 8**

S. no.	Bacterial Isolates	Sugar Fermentation				IMVIC				Identified Bacteria
		lactose	Glucose	Mannitol	Sucrose	Indole	MR	VP	C	
1	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	+ve	+ve	-ve	-ve	<i>E. coli</i>
2	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	+ve	+ve	-ve	-ve	<i>E. coli</i>
3	US -1	Acid +gas	Acid +gas	Acid +gas	Acid +gas	-ve	-ve	+ve	-ve	<i>Klebsiella pneumoniae</i>

On the basis of the above results, it can be said that the clinical isolates were found to be *Escherichia coli* and *Klebsiella pneumoniae*.

**TABLE 9: COLLECTION OF ISOLATES FROM URINE SAMPLES**

S. no.	Clinical samples	Total no. of isolates collected	<i>E. coli</i>	<i>Klebsiella pneumoniae</i>	<i>S. aureus</i>	<i>Pseudomonas aeruginosa</i>
1	US-1	4	1	1	1	1
2	US-2	4	-	1	2	1
3	US-3	1	1	-	-	-
4	US-4	2	-	-	1	1
5	US-5	4	1	1	1	1
6	US-6	1	-	-	-	1
7	US-7	1	-	-	1	-
8	US-8	3	2	1	-	-

**TABLE 10: ANTIBIOTIC SUSCEPTIBILITY OF *E. COLI***

Samples	Antibiotics	Concentration of antibiotics	Diameter of zones (mm)	Interpretation
US-1	Norfloxacin	10 mcg	17 mm	Resistant
US-1	Amikacin	30 mcg	17 mm	Sensitive
US-2	Gentamycin	10 mcg	13 mm	Resistant
US-2	Erythromycin	15 mcg	NZ	Resistant
US-3	Cefepime	10 mcg	NZ	Resistant
US-3	Amoxicillin	30 mcg	NZ	Resistant
US-4	Morfloxacin	5 mcg	28 mm	Sensitive
US-4	Ofloxacin	5 mcg	28 mm	Sensitive
US-5	Tetracycline	30 mcg	NZ	Resistant
US-5	Streptomycin	10 mcg	NZ	Resistant
US-6	Methicillin	5 mcg	NZ	Resistant
US-6	Clindamycin	20 mcg	NZ	Resistant
US-7	Penicillin	10 units	NZ	Resistant

US-7	Rifamicine	5 mcg	16 mm	Resistant
US-8	Lomefloxacin	10 mcg	17 mm	Resistant
US-8	Cotrimoxazole	25 mcg	23 mm	Sensitive
US-8	Tobramicine	10 mcg	15 mm	Resistant

\*Note: Control = antibiotics without Tris, S = Sensitive, NZ = NO zone of inhibition, I = Intermediate, R = Resistant

**TABLE 11: ANTIBIOTIC SUSCEPTIBILITY OF *KLEBSIELLA PNEUMONIAE***

Samples	Antibiotics	Concentration of antibiotics	Diameter of zones (mm)	Interpretation
US-1	Norfloxacin	10 mcg	22 mm	Sensitive
US-1	Amikacin	30 mcg	18 mm	Sensitive
US-2	Gentamycin	10 mcg	16 mm	Resistant
US-2	Erythromycin	15 mcg	NZ	Resistant
US-3	Tobramicine	10 mcg	13 mm	Resistant
US-3	Amoxicillin	30 mcg	NZ	Resistant
US-4	Morfloxacin	5 mcg	18 mm	Resistant
US-4	Ofloxacin	5 mcg	21 mm	Sensitive
US-5	Teicoplanin	30 mcg	19 mm	Sensitive
US-5	Tetracycline	10 mcg	13 mm	Resistant
US-6	Methicillin	5 mcg	NZ	Resistant
US-6	Linezolid	30 mcg	NZ	Resistant
US-7	Penicillin	10 units	NZ	Resistant
US-7	Amphicillin	30mcg	16 mm	Intermediate
US-8	Lomefloxacin	10 mcg	16 mm	Resistant
US-8	Tobramicine	10 mcg	17 mm	Resistant

\*Note: Control = antibiotics without Tris, S = Sensitive, NZ = NO zone of inhibition, I = Intermediate, R = Resistant

**TABLE 12: ANTIBIOTIC SUSCEPTIBILITY OF *STAPHYLOCOCCUS AUREUS***

Samples	Antibiotics	Concentration of antibiotics	Diameter of zones (mm)	Interpretation
US-1	Vancomycin	10 mcg	17 mm	Sensitive
US-1	Amikacin	30 mcg	15 mm	Sensitive
US-1	Gentamycin	10 mcg	15 mm	Resistant
US-2	Erythromycin	15 mcg	31 mm	Sensitive
US-2	Norfloxacin	10 mcg	21 mm	Sensitive
US-3	Amphicillin	30 mcg	26 mm	Sensitive
US-3	Morfloxacin	5 mcg	18 mm	Resistant
US-4	Ofloxacin	5 mcg	21 mm	Sensitive
US-4	Tetracycline	30 mcg	NZ	Resistant
US-5	Streptomycin	10 mcg	NZ	Resistant
US-5	Methicillin	5 mcg	NZ	Resistant
US-6	Clindamycin	20 mcg	NZ	Resistant
US-6	Penicillin	10 units	NZ	Resistant
US-7	Rifampicin	5 mcg	16 mm	Sensitive
US-7	Lomefloxacin	10 mcg	21 mm	Resistant
US-8	Cotrimoxazole	25 mcg	NZ	Resistant
US-8	Levofloxacin	10 mcg	15 mm	Resistant

\*Note: Control = antibiotics without Tris, S = Sensitive, NZ = NO zone of inhibition, I = Intermediate, R = Resistant

**TABLE 13: ANTIBIOTIC SUSCEPTIBILITY OF *PSEUDOMONAS AERUGINOSA***

Samples	Antibiotics	Concentration of antibiotics	Diameter of zones (mm)	Interpretation
US-1	Norfloxacin	10 mcg	17 mm	Intermediate
US-1	Amikacin	30 mcg	20 mm	Sensitive
US-2	Gentamycin	10 mcg	20 mm	Sensitive
US-2	Erythromycin	15 mcg	NZ	Resistant
US-3	Ceprofloxacin	10 mcg	18 mm	Sensitive
US-3	Meropenem	10 mcg	15 mm	Intermediate
US-4	Morfloxacin	5 mcg	28 mm	Sensitive
US-4	Cefepime	30 mcg	NZ	Resistant
US-5	Tetracycline	30 mcg	13 mm	Resistant
US-5	Streptomycin	10 mcg	NZ	Resistant
US-6	Methicillin	5 mcg	13 mm	Intermediate
US-6	Clindamycin	2 mcg	NZ	Resistant
US-7	Penicillin	10 units	NZ	Resistant
US-7	Rifamicine	5 mcg	16 mm	Resistant
US-7	Lomefloxacin	10 mcg	16 mm	Resistant
US-8	Cotrimoxazole	25 mcg	NZ	Resistant
US-8	Tobramicine	10 mcg	32 mm	Sensitive

\*Note: Control = antibiotics without Tris, S = Sensitive, NZ = NO zone of inhibition, I = Intermediate, R = Resistant




FIG. 1: COLLECTION OF URINE SAMPLES





FIG. 2: GROWTH OF ISOLATES ON EMB




FIG. 3: GROWTH OF ISOLATES ON CLED




FIG. 4: GROWTH OF ISOLATES ON MSA




FIG. 5: GROWTH OF ISOLATES ON PIA



FIG. 6: RESULTS OF SUGAR FERMENTATION AND IMVIC FOR *E. COLI*



FIG. 7: RESULTS OF SUGAR FERMENTATION AND IMVIC TESTS FOR *KLEBSIELLA PNEUMONIAE*






FIG. 8: RESULTS OF SUGAR FERMENTATION AND IMVIC TEST FOR *S. AUREUS*




FIG. 9: RESULTS OF SUGAR FERMENTATION AND IMVIC TEST FOR *PSEUDOMONAS AERUGINOSA*




FIG. 10: ANTIBIOTIC SUSCEPTIBILITY TEST AGAINST *E. COLI*





FIG. 11: ANTIBIOTIC SUSCEPTIBILITY TEST AGAINST *KLEBSIELLA PNEUMONIAE*




FIG. 12: ANTIBIOTIC SUSCEPTIBILITY TEST AGAINST *S. AUREUS*




FIG. 13: ANTIBIOTIC SUSCEPTIBILITY TEST AGAINST *PSEUDOMONAS AERUGINOSA*



FIG. 14: RESULTS OF MIC TEST BY E- STRIP METHOD

**DISCUSSION AND CONCLUSION:** During the study, the urine sample was inoculated on selective media like EMB, CLED, MSA, PIA, CIA medium followed with biochemical tests. The results showed the isolation of *E. coli*, *K. pneumoniae*, *S. aureus*, and *Pseudomonas aeruginosa*.

The results of antibiotic susceptibility showed that norfloxacin, gentamycin, rifamycin, lomeflo-xacine and tobramycin, penicillin, clindamycin, methicillin, streptomycin, tetracycline, amoxicilline, ciprofloxacin and erythromycin are resistant to *E. coli* while amikacin, moxifloxacin, ofloxacin and cotrimoxazole are found to be sensitive against the same. It was observed that norfloxacin, ofloxacin, and teicoplanin are the sensitive antibiotics for klebsiella while gentamycin, tobramycine, moxifloxacin, tetracycline, lomefloxacin-cine, and tobramycine are resistant antibiotics. Ampicillin shows intermediate state and methicillin, linzoids, penicillin, amoxicillin, erythromycin showed resistance against Klebsiella. Norfloxacin, amikacin, ofloxacin, vancomycin, erythromycin, ampicillin and rifamycin are sensitive antibiotics. Gentamycin, moxifloxacin, lomefloxacin, levo-

floxacin, penicillin, clinda-mycine, methicillin, streptomycin, tetracycline were found resistant against *Staphylococcus aureus* while amikacin, gentamycin, ciprofloxacin and tobramycin were found to be sensitive antibiotics against the same. Tetracycline, rifamycin, lome-floxacine, erythromycin, cefepime, streptomycin, penicillin, clindamycin were found to be resistant against *P. aeruginosa*. Antibiotics like norfloxacin, meropenem, and methicillin were found to be intermediate antibiotics.

ESBL producing organisms, being commonest nosocomial pathogens, and thus it is much essential to detect and treat them as early as possible The study thus suggests that the present study will provide the basis for routine screening of antibiotics against ESBL producing bacteria.

The study will provide the basis for optimizing the different lines of antibiotics against specific bacterial cultures. Thus, the results will help in exploring the specific treatment against such pathogens borne infections.

**ACKNOWLEDGEMENT:** The authors acknowledge the research staff and colleagues of R&D project on the concerned topic.

**CONFLICTS OF INTEREST:** The authors declare no conflicts of interest.

## REFERENCES:

- Burcu BE, Açık L and Sultan N: Phenotypic and molecular characterization of SHV, TEM, CTX-M and extended-spectrum beta lactamase produced by *Escherichia coli*, *Acinetobacter baumannii* and *Klebsiella* isolates in a Turkish hospital. *African Journal of Microbiology Research* 2010; 4(8): 650-54.
- Mandell GL, Bennett JE and Dolin R: Principles and practice of infectious diseases. Churchill Livingstone 2005; 881-82.
- Ronald A: The etiology of urinary tract infection: Traditional and emerging pathogens. *Am J Med.* 2002; 113(S1A): 14S-9S.
- Drieux L, Brossier F, Sougakoff W and Jarlier V: Phenotypic detection of extended-spectrum  $\beta$ -lactamase production in Enterobacteriaceae: Review and bench guide. *Clin Microbiol* 2008; 14 (S1): 90-103.
- Gazin M, Paasch F, Goosens H and Malhotra-Kumar S: Current trends in culture-based and molecular detection of extended-spectrum  $\beta$ -lactamase-harboring and carbapenem-resistant Enterobacteriaceae. *J Clin Microbiol* 2012; 50(4): 1140-46.
- Schwaber MJ, Navon-Venezia S, Kaye KS, Ben-Ami R, Schwartz D and Carmeli Y: Clinical and economic impact of bacteremia with extended-spectrum  $\beta$  lactamase-producing Enterobacteriaceae. *Antimicrob. Agents Chemother* 2006; 50: 1257-62.
- Podschun R and Ullmann U: *Klebsiella* spp. as nosocomial pathogens: epidemiology, taxonomy, typing methods, and pathogenicity factors. *Clin Microbiol Rev* 1998; 11(4): 589-03.
- Kim YK and Pai H: Bloodstream Infections by Extended-Spectrum {beta}-Lactamase-Producing *Escherichia coli* and *Klebsiella pneumoniae* in Children: Epidemiology and Clinical Outcome. *Antimicrob. Agents Chemother.* 2002; 46(5): 1481-91.
- Nwankwo EO, Magaji NS and Tijjani J: Antibiotic susceptibility pattern of Extended Spectrum Betalactamase (ESBL) Producers and other Bacterial Pathogens. *Tropical Journal of Pharmaceutical Research* 2015; 14(7): 1273-78.
- Ullah F, Malik SA and Ahmed J: Antimicrobial susceptibility pattern and ESBL prevalence in *Klebsiella pneumoniae* from urinary tract infections. *African Journal of Microbiology Research* 2009; 3(11): 676-80.
- Ejaz H, Haq IU, Zafar A, Mahmood S and Javed MM: Urinary tract infections caused by extended-spectrum  $\beta$ -lactamase (ESBL) producing *Escherichia coli* and *Klebsiella pneumoniae*. *African Journal of Biotechnology* 2011; 10(73): 16661-66.
- Taha S, Youssef N, Elkazaz A and Ramadan H: Detection of extended-spectrum beta-lactamases (ESBLs) in clinical isolates of *Klebsiella pneumoniae* using the ESBL NDP test and flow cytometric assay in comparison to the standard disc diffusion. *African Journal of Microbiology Research Science* 2016; 10(8): 238-44.
- Tissera S and Lee SM: Isolation of Extended Spectrum  $\beta$ -lactamase (ESBL) producing bacteria from urban surface waters in Malaysia. *Malays J Med Sci* 2013; 20(3).
- Ray J, Paul R, Haldar A and Mondol S: A study on antibiotic resistance pattern of *Escherichia coli* isolated from urine specimens in Eastern India. *International Journal of Medical Sciences and Public Health* 2015; 3(3): 1670-74.
- Wasnik DD and Tumane PM: Detection of extended spectrum beta - lactamase producing *E. coli* & *Klebsiella species* causing urinary tract infection. *Journal of Biomedical and Pharmaceutical Res* 2013; 2(3): 122-25.
- Vinodhini R, Moorthy K, Palanivel P, Puniitha T, Saranya S, Bhuvaneshwari M and Kanimozhi C: Detection and antimicrobial susceptibility pattern of ESBL producing gram negative bacteria. *Asian Journal of Pharmacy and Clinical Research* 2014; 7(1): 243-47.
- Dugal S and Purohit H: Antimicrobial susceptibility profile and detection of extended spectrum beta-lactamase production by gram negative uropathogens. *International Journal of Pharmacy and Pharmaceutical Sciences* 2013; 5(4): 434-38.
- Varkey DR, Balaji V and Abraham J: Molecular characterisation of extended spectrum beta lactamase producing strains from blood sample. *Int Jou of Pharmacy and Pharmaceutical Sciences* 2014; 6(3): 276-78.
- Yadav RR and Chauhan PB: The detection of extended spectrum Beta Lactamases (ESBLs) producing *Escherichia coli* isolated from clinical sample. *International Journal of Advance Research in Biological Science* 2016; 3(5): 9-1
- Motmedifar M, Mansury D, Sarvari J, Shirazi B and Khaledi A: Antibiotic susceptibility pattern and identification of extended spectrum of Beta lactamases (ESBLs) in clinical isolates of *Klebsiella pneumoniae* from Shiraz, Iran. *Iranian Jou of Microbiol* 2016; 8(1): 55-61.
- Ahmad M, Hassan M, Khalid A, Tariq I, Asad MHHB, Samad A, Mahmood Q and Murtaza G: Prevalence of extended spectrum Beta lactamase and antimicrobial susceptibility pattern of clinical isolates of *Pseudomonas* from patients of Khyber Pakhtunkhwa, Pakistan. *Biomed Research International* 2016; 1-10.
- Abayneh M, Tesfaw G and Abdissa A: Isolation of extended-spectrum Beta-lactamase producing *E. coli* and *Klebsiella pneumoniae* from patients with community onset UTI infections in Jimma University specialized hospital, south west Ethiopia. *Canadian Journal of Infectious diseases and Medical Microbiology* 2018; 1-8.
- Marguilex KR, Srijan A, Ruekit S, Nobthai P, Poramathikul K, Pandey P, Serichantalergs O and Shreshtha SK: Extended-spectrum Beta lactamases prevalence and virulence factor characterization of enterotoxigenic *E. coli* responsible for acute diarrhoea in Nepal from 2001 to 2016. *Antimicrobial resistance and Infection control* 2018; 87.
- Vila J, Saez- Lopez E, Johnson JR, Romling U, Dobrindt U, Canton R, Giske CG, Naas T, Carattoli A and Martinez-Medina M: *Eisearchia coli*: An old friend with new tidings. *Frontiers FMS Microbiol. Rev* 2016; 40: 437-63.
- Bajaj P, Singh MS and Viridi JS: *E. coli* Beta lactamases: What really matters? *Front Microbiol* 2016.
- Kumar P, Bag S, Ghosh TS, Dety P, Dayal M, Saha B, Verma J, Pant A, Saxena S and Desigamani A: Molecular insights into antimicrobial resistance traits of multi-drug resistant enteric pathogens isolated from India. *Science Reporter* 2017; 7: 14468.
- Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing- 28<sup>th</sup> edition. Wayne: Clinical and Laboratory Standards Institute 2017.

28. Parajuli NOP, Mharjan P, Parajuli H, Joshi G, Paudel D, Sayami S and Khanal PR: High rates of multi-drug resistance among uropathogenic *E. coli* in children and

analyses of ESBL producers from Nepal. Antimicrob Resist Infect Control 2017; 6: 9.

**How to cite this article:**

Hattewar J, Koul A and Mathur A: Antibiotic susceptibility pattern of ESBL producing bacteria from urinary tract infection (UTI). Int J Pharm Sci & Res 2020; 11(2): 799-10. doi: 10.13040/IJPSR.0975-8232.11(2).799-10.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)